

Type 2 diabetes in adults: management

NICE guideline

Published: 2 December 2015

www.nice.org.uk/guidance/ng28

Your responsibility

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals and practitioners are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or the people using their service. It is not mandatory to apply the recommendations, and the guideline does not override the responsibility to make decisions appropriate to the circumstances of the individual, in consultation with them and their families and carers or guardian.

Local commissioners and providers of healthcare have a responsibility to enable the guideline to be applied when individual professionals and people using services wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with complying with those duties.

Commissioners and providers have a responsibility to promote an environmentally sustainable health and care system and should assess and reduce the environmental impact of implementing NICE recommendations wherever possible.

Contents

Overview	5
Who is it for?	5
Introduction	6
Reasons for the update	6
Medicines	7
Patient-centred care	8
Key priorities for implementation	9
Patient education	9
Dietary advice	9
Blood pressure management	9
Blood glucose management.....	10
Drug treatment.....	10
1 Recommendations	12
Terms used in this guideline	12
1.1 Individualised care	12
1.2 Patient education.....	12
1.3 Dietary advice and bariatric surgery	14
1.4 Blood pressure management	15
1.5 Antiplatelet therapy	17
1.6 Blood glucose management.....	17
1.7 Managing complications	27
2 Research recommendations	34
2.1 The effects of stopping and/or switching drug treatments to control blood glucose levels	34
2.2 Non-metformin-based drug treatment combinations to control blood glucose levels.....	34
2.3 Drug treatment (third intensification) for when blood glucose levels are inadequately controlled by 3 oral antidiabetic drugs and/or insulin combinations.....	35
2.4 Long-term outcomes associated with blood glucose lowering agents.....	35

2.5 Self-monitoring of blood glucose levels..... 35

More information..... 36

Update information.....37

Recommendations from NICE clinical guideline 87 (2009) that have been amended..... 38

This guideline replaces CG87, TA203, TA248 and CG66.

This guideline is the basis of QS6.

Overview

This guideline covers the care and management of type 2 diabetes in adults (aged 18 and over). It focuses on patient education, dietary advice, managing cardiovascular risk, managing blood glucose levels, and identifying and managing long-term complications.

Who is it for?

- Healthcare professionals that care for adults with diabetes
- Commissioners and providers of diabetes services
- Adults with type 2 diabetes, and their families and carers

Introduction

This guidance updates and replaces 'Type 2 diabetes' (NICE guideline CG87). The recommendations are labelled according to when they were originally published (see [update information](#) for details).

Type 2 diabetes is a chronic metabolic condition characterised by insulin resistance (that is, the body's inability to effectively use insulin) and insufficient pancreatic insulin production, resulting in high blood glucose levels (hyperglycaemia). Type 2 diabetes is commonly associated with obesity, physical inactivity, raised blood pressure, disturbed blood lipid levels and a tendency to develop thrombosis, and therefore is recognised to have an increased cardiovascular risk. It is associated with long-term microvascular and macrovascular complications, together with reduced quality of life and life expectancy.

In 2013, over 3.2 million adults were diagnosed with diabetes, with prevalence rates of 6% and 6.7% in England and Wales respectively. It is estimated that about 90% of adults currently diagnosed with diabetes have type 2 diabetes. Type 2 diabetes is more common in people of African, African-Caribbean and South Asian family origin. It can occur in all age groups and is increasingly being diagnosed in children.

Multiple vascular risk factors and wide-ranging complications make diabetes care complex and time-consuming, and many areas of healthcare services must be involved for optimal management. Necessary lifestyle changes, the complexities and possible side effects of therapy make patient education and self-management important aspects of diabetes care. Diabetes care is estimated to account for at least 5% of UK healthcare expenditure, and up to 10% of NHS expenditure.

This guideline contains recommendations for managing type 2 diabetes in adults, and focuses on patient education, dietary advice, managing cardiovascular risk, managing blood glucose levels, and identifying and managing long-term complications. The guideline does not cover diagnosis, secondary diabetes, type 1 diabetes in adults, diabetes in pregnancy and diabetes in children and young people.

Reasons for the update

Since the publication of the 2009 guideline, availability of new evidence and several key developments have prompted an update in the following areas: managing blood glucose levels, antiplatelet therapy and erectile dysfunction. In particular, reasons included safety concerns surrounding some blood glucose lowering medicines, new evidence on new dipeptidyl peptidase-4

(DPP-4) inhibitors and glucagon-like peptide-1 (GLP-1) receptor agonists, new indications and licensed combinations for licensed class members and the potential impact of drugs coming off patent on health-economic issues. In addition, new evidence and safety issues relating to the off-label use of antiplatelet therapy (aspirin and clopidogrel) in the primary prevention of cardiovascular disease motivated an update of this review.

Medicines

The guideline will assume that prescribers will use a medicine's summary of product characteristics to inform decisions made with individual patients.

This guideline recommends some medicines for indications for which they do not have a UK marketing authorisation at the date of publication, if there is good evidence to support that use. The prescriber should follow relevant professional guidance, taking full responsibility for the decision. The patient (or those with authority to give consent on their behalf) should provide informed consent, which should be documented. See the General Medical Council's [Prescribing guidance: prescribing unlicensed medicines](#) for further information. Where recommendations have been made for the use of medicines outside their licensed indications ('off-label use'), these medicines are marked with a footnote in the recommendations.

Patient-centred care

This guideline offers best practice advice on the care of adults with type 2 diabetes.

When caring for older adults with type 2 diabetes, particular consideration should be given to their broader health and social care needs. Older people are more likely to have co-existing conditions and to be on a greater number of medicines. Their ability to benefit from risk-reduction interventions in the longer term may also be reduced.

Much of the evidence base used to inform this guideline has been generated from studies involving younger adults (study mean ages ranged from 45 to 68 years). While the Guideline Development Group thought that the recommendations are applicable to a wider age group, they highlighted that there needs to be flexibility, to ensure that the care of older people with diabetes also addresses their broader health and social care needs.

Patients and healthcare professionals have rights and responsibilities as set out in the [NHS Constitution for England](#) – all NICE guidance is written to reflect these. Treatment and care should take into account individual needs and preferences. Patients should have the opportunity to make informed decisions about their care and treatment, in partnership with their healthcare professionals. If the patient is under 16, their family or carers should also be given information and support to help the child or young person to make decisions about their treatment. If it is clear that the child or young person fully understands the treatment and does not want their family or carers to be involved, they can give their own consent. Healthcare professionals should follow the [Department of Health's advice on consent](#). If someone does not have capacity to make decisions, healthcare professionals should follow the [code of practice that accompanies the Mental Capacity Act](#) and the supplementary [code of practice on deprivation of liberty safeguards](#).

NICE has produced guidance on the components of good patient experience in adult NHS services. All healthcare professionals should follow the recommendations in [patient experience in adult NHS services](#).

Key priorities for implementation

The following recommendations have been identified as priorities for implementation. The full list of recommendations is in [section 1](#).

Patient education

- Offer structured education to adults with type 2 diabetes and/or their family members or carers (as appropriate) at and around the time of diagnosis, with annual reinforcement and review. Explain to people and their carers that structured education is an integral part of diabetes care. [2009]
- Ensure that any structured education programme for adults with type 2 diabetes includes the following components:
 - It is evidence-based, and suits the needs of the person.
 - It has specific aims and learning objectives, and supports the person and their family members and carers in developing attitudes, beliefs, knowledge and skills to self-manage diabetes.
 - It has a structured curriculum that is theory-driven, evidence-based and resource-effective, has supporting materials, and is written down.
 - It is delivered by trained educators who have an understanding of educational theory appropriate to the age and needs of the person, and who are trained and competent to deliver the principles and content of the programme.
 - It is quality assured, and reviewed by trained, competent, independent assessors who measure it against criteria that ensure consistency.
 - The outcomes are audited regularly. [2015]

Dietary advice

- Integrate dietary advice with a personalised diabetes management plan, including other aspects of lifestyle modification, such as increasing physical activity and losing weight. [2009]

Blood pressure management

- Add medications if lifestyle advice does not reduce blood pressure to below 140/80 mmHg