

Mappings valued in the Wasserstein space

Hugo Lavenant^a


May 14th, 2020

Università Commerciale Luigi Bocconi

^aDepartment of Mathematics, University of British Columbia

The Wasserstein¹ space


D convex and compact domain of \mathbb{R}^d .



¹and Monge, Lévy, Fréchet, Kantorovich, Rubinstein, etc.

The Wasserstein¹ space

D convex and compact domain of \mathbb{R}^d .




$\mathcal{P}(D)$ space of probability measures on D .


The **Wasserstein space** is the space $\mathcal{P}(D)$ endowed with the Wasserstein distance.

¹and Monge, Lévy, Fréchet, Kantorovich, Rubinstein, etc.









In this presentation

1. A quick introduction to the Wasserstein space
2. Harmonic mappings valued in the Wasserstein space
3. An application in nonlinear elasticity

1. A quick introduction to the Wasserstein space

The metric tensor in the Wasserstein space




The metric tensor in the Wasserstein space




The metric tensor in the Wasserstein space

Vertical derivative




The metric tensor in the Wasserstein space

Vertical derivative




Horizontal derivative




A particle located at x moves to $x + h\nu(x)$

The metric tensor in the Wasserstein space

Vertical derivative




Horizontal derivative



$$\partial_t \mu = -\nabla \cdot (\mu \mathbf{v})$$

A particle located at x moves to $x + h\mathbf{v}(x)$

The metric tensor in the Wasserstein space



- Quadratic Optimal Transport: the square of the norm of the speed is

$$\min_{\mathbf{v}: D \rightarrow \mathbb{R}^d} \left\{ \int_D |\mathbf{v}(y)|^2 \mu(dy) : -\nabla \cdot (\mu \mathbf{v}) = \partial_t \mu \right\}.$$

Action and geodesics

If $\mu : [0, 1] \rightarrow \mathcal{P}(D)$ is given, its **action** is

$$\mathcal{A}(\mu) := \min_{\mathbf{v}} \left\{ \frac{1}{2} \int_0^1 \int_D |\mathbf{v}_t|^2 \, d\mu_t \, dt : \partial_t \mu_t + \nabla \cdot (\mu_t \mathbf{v}_t) = 0 \right\}.$$

Action and geodesics

If $\mu : [0, 1] \rightarrow \mathcal{P}(D)$ is given, its **action** is

$$\mathcal{A}(\mu) := \min_{\mathbf{v}} \left\{ \frac{1}{2} \int_0^1 \int_D |\mathbf{v}_t|^2 \, d\mu_t \, dt : \partial_t \mu_t + \nabla \cdot (\mu_t \mathbf{v}_t) = 0 \right\}.$$

The Wasserstein distance W_2 is

$$\frac{1}{2} W_2^2(\rho, \nu) = \min_{\mu} \{ \mathcal{A}(\mu) : \mu_0 = \rho, \mu_1 = \nu \},$$


Action and geodesics

If $\mu : [0, 1] \rightarrow \mathcal{P}(D)$ is given, its **action** is

$$\mathcal{A}(\mu) := \min_{\mathbf{v}} \left\{ \frac{1}{2} \int_0^1 \int_D |\mathbf{v}_t|^2 \, d\mu_t \, dt : \partial_t \mu_t + \nabla \cdot (\mu_t \mathbf{v}_t) = 0 \right\}.$$

The Wasserstein distance W_2 is

$$\frac{1}{2} W_2^2(\rho, \nu) = \min_{\mu} \{ \mathcal{A}(\mu) : \mu_0 = \rho, \mu_1 = \nu \},$$



and the minimizers are the constant-speed geodesics.

Wasserstein spaces on manifolds

The definition can be extended for D Riemannian manifold, and similar numerical methods can be used².



²H. Lavenant, S. Clatici, E. Chien and J. Solomon. *Dynamical Optimal Transport on Discrete Surfaces*. 2018.

Wasserstein spaces on manifolds

The definition can be extended for D Riemannian manifold, and similar numerical methods can be used².



²H. Lavenant, S. Claiç, E. Chien and J. Solomon. *Dynamical Optimal Transport on Discrete Surfaces*. 2018.

Why use Wasserstein distances?

Finding matching between distributions of mass is an ubiquitous task.

³Y. Brenier. *The least action principle and the related concept of generalized flows for incompressible perfect fluids*. 1989.

⁴R. Jordan, D. Kinderlehrer, and F. Otto. *The variational formulation of the Fokker–Planck equation* 1998.

⁵J.-D. Benamou, G. Carlier, and F. Santambrogio. *Variational mean field games*. 2017.

⁶G. Peyré and M. Cuturi. *Computational optimal transport*. 2019.

Why use Wasserstein distances?

Finding matching between distributions of mass is an ubiquitous task. Some use of the Wasserstein distance:

- **Least action principle** in fluid mechanics³.
- **Gradient flows** in Wasserstein space yield parabolic PDEs⁴.

³Y. Brenier. *The least action principle and the related concept of generalized flows for incompressible perfect fluids*. 1989.

⁴R. Jordan, D. Kinderlehrer, and F. Otto. *The variational formulation of the Fokker–Planck equation* 1998.

⁵J.-D. Benamou, G. Carlier, and F. Santambrogio. *Variational mean field games*. 2017.

⁶G. Peyré and M. Cuturi. *Computational optimal transport*. 2019.

Why use Wasserstein distances?

Finding matching between distributions of mass is an ubiquitous task. Some use of the Wasserstein distance:

- **Least action principle** in fluid mechanics³.
- **Gradient flows** in Wasserstein space yield parabolic PDEs⁴.
- **Nash equilibrium** in variational Mean Field Games⁵.

³Y. Brenier. *The least action principle and the related concept of generalized flows for incompressible perfect fluids*. 1989.

⁴R. Jordan, D. Kinderlehrer, and F. Otto. *The variational formulation of the Fokker–Planck equation* 1998.

⁵J.-D. Benamou, G. Carlier, and F. Santambrogio. *Variational mean field games*. 2017.

⁶G. Peyré and M. Cuturi. *Computational optimal transport*. 2019.

Why use Wasserstein distances?

Finding matching between distributions of mass is an ubiquitous task. Some use of the Wasserstein distance:

- **Least action principle** in fluid mechanics³.
- **Gradient flows** in Wasserstein space yield parabolic PDEs⁴.
- **Nash equilibrium** in variational Mean Field Games⁵.
- **Data fitting** in machine learning⁶.


³Y. Brenier. *The least action principle and the related concept of generalized flows for incompressible perfect fluids*. 1989.

⁴R. Jordan, D. Kinderlehrer, and F. Otto. *The variational formulation of the Fokker–Planck equation* 1998.

⁵J.-D. Benamou, G. Carlier, and F. Santambrogio. *Variational mean field games*. 2017.

⁶G. Peyré and M. Cuturi. *Computational optimal transport*. 2019.


Regularizer in machine learning⁷



- Stochastic process (X_t), access to *independent samples* at different times.

⁷B. Schmitzer, K. P. Schäfers, and B. Wirth. *Dynamic Cell Imaging in PET with Optimal Transport Regularization*. 2019.


Regularizer in machine learning⁷



- Stochastic process (X_t), access to *independent samples* at different times.

⁷B. Schmitzer, K. P. Schäfers, and B. Wirth. *Dynamic Cell Imaging in PET with Optimal Transport Regularization*. 2019.

Regularizer in machine learning⁷




- Stochastic process (X_t), access to *independent samples* at different times.
- Reconstruction of the process?

$$\min_{\rho} \left\{ \sum_{t_i} \text{Loss}(\rho_{t_i}, \text{data}_{t_i}) + \lambda \underbrace{\mathcal{A}(\rho)}_{\text{regularization}} \right\}$$

⁷B. Schmitzer, K. P. Schäfers, and B. Wirth. *Dynamic Cell Imaging in PET with Optimal Transport Regularization*. 2019.

Regularizer in machine learning⁷



- Stochastic process (X_t), access to *independent samples* at different times.
- Reconstruction of the process?
- Presence of noise?
Handling birth and death of cells?

$$\min_{\rho} \left\{ \sum_{t_i} \text{Loss}(\rho_{t_i}, \text{data}_{t_i}) + \lambda \underbrace{\mathcal{A}(\rho)}_{\text{regularization}} \right\}$$


⁷B. Schmitzer, K. P. Schäfers, and B. Wirth. *Dynamic Cell Imaging in PET with Optimal Transport Regularization*. 2019.

2. Harmonic mappings valued in the Wasserstein space

Measure-valued mappings⁸

Ω bounded set of \mathbb{R}^n with Lipschitz boundary


We study $\mu : \Omega \rightarrow \mathcal{P}(D)$.



Measure-valued mappings⁸

Ω bounded set of \mathbb{R}^n with Lipschitz boundary (before $\Omega = [0, 1] \subset \mathbb{R}$).

We study $\mu : \Omega \rightarrow \mathcal{P}(D)$.




Definition of $\text{Dir}(\mu) = \frac{1}{2} \int_{\Omega} |\nabla \mu|^2$ the **Dirichlet energy** generalizing \mathcal{A} .

⁸H. Lavenant. *Harmonic mappings valued in the Wasserstein space*. 2019.

Measure-valued mappings⁸

Ω bounded set of \mathbb{R}^n with Lipschitz boundary (before $\Omega = [0, 1] \subset \mathbb{R}$).

We study $\mu : \Omega \rightarrow \mathcal{P}(D)$.



Definition of $\text{Dir}(\mu) = \frac{1}{2} \int_{\Omega} |\nabla \mu|^2$ the **Dirichlet energy** generalizing \mathcal{A} .


Minimizers of Dir are called harmonic mappings (valued in the Wasserstein space).

⁸H. Lavenant. *Harmonic mappings valued in the Wasserstein space*. 2019.

Measure-valued mappings⁸

Ω bounded set of \mathbb{R}^n with Lipschitz boundary (before $\Omega = [0, 1] \subset \mathbb{R}$).

We study $\mu : \Omega \rightarrow \mathcal{P}(D)$.



Definition of $\text{Dir}(\mu) = \frac{1}{2} \int_{\Omega} |\nabla \mu|^2$ the **Dirichlet energy** generalizing \mathcal{A} .

Minimizers of Dir are called harmonic mappings (valued in the Wasserstein space).

If $u : \Omega \rightarrow D$ and $\mu(x) := \delta_{u(x)}$ then $\text{Dir}(\mu) = \frac{1}{2} \int_{\Omega} |\nabla u|^2$.

⁸H. Lavenant. *Harmonic mappings valued in the Wasserstein space*. 2019.

The Dirichlet energy⁹

Definition

If $\mu : \Omega \rightarrow \mathcal{P}(D)$ is given,

$$\text{Dir}(\mu) := \min_{\mathbf{v}} \left\{ \frac{1}{2} \int_{\Omega} \int_D |\mathbf{v}|^2 d\mu : \nabla_{\Omega} \mu + \nabla_D \cdot (\mu \mathbf{v}) = 0 \right\},$$

where $\mathbf{v} : \Omega \times D \rightarrow \mathbb{R}^{nd}$ “density of Jacobian matrix”.

If $\Omega = [0, 1]$ it coincides with \mathcal{A} .

⁹Brenier. *Extended Monge-Kantorovich theory*. 2003.

Equivalence with a metric definition¹⁰

$$\frac{W_2^2(\mu(x), \mu(x'))}{\varepsilon^2}$$

¹⁰Korevaar and Schoen. *Sobolev spaces and harmonic maps for metric space targets*. 1993.

Equivalence with a metric definition¹⁰

$$\frac{1}{\varepsilon^n} \int_{\Omega} \frac{W_2^2(\mu(x), \mu(x'))}{\varepsilon^2} \mathbb{1}_{|x-x'| \leq \varepsilon} dx'$$

¹⁰Korevaar and Schoen. *Sobolev spaces and harmonic maps for metric space targets*. 1993.

Equivalence with a metric definition¹⁰

$$\mathrm{Dir}_\varepsilon(\mu) := \frac{C_n}{2} \int_\Omega \frac{1}{\varepsilon^n} \int_\Omega \frac{W_2^2(\mu(x), \mu(x'))}{\varepsilon^2} \mathbb{1}_{|x-x'| \leq \varepsilon} \, dx' \, dx$$

Proposed by Korevaar, Schoen (and Jost) for mappings valued in metric spaces.

¹⁰Korevaar and Schoen. *Sobolev spaces and harmonic maps for metric space targets*. 1993.

Equivalence with a metric definition¹⁰

$$\mathrm{Dir}_\varepsilon(\mu) := \frac{C_n}{2} \int_\Omega \frac{1}{\varepsilon^n} \int_\Omega \frac{W_2^2(\mu(x), \mu(x'))}{\varepsilon^2} \mathbb{1}_{|x-x'| \leq \varepsilon} \, dx' \, dx$$

Proposed by Korevaar, Schoen (and Jost) for mappings valued in metric spaces.

Theorem

There holds

$$\lim_{\varepsilon \rightarrow 0} \mathrm{Dir}_\varepsilon = \mathrm{Dir},$$

and the convergence holds pointwisely and in the sense of Γ -convergence along the sequence $\varepsilon_m = 2^{-m}$.

¹⁰Korevaar and Schoen. *Sobolev spaces and harmonic maps for metric space targets*. 1993.


Curvature and convexity

If $\mu, \nu \in \mathcal{P}(D)$, two ways to interpolate.



Curvature and convexity

If $\mu, \nu \in \mathcal{P}(D)$, two ways to interpolate.



The **displacement** interpolation



- Midpoint of the geodesic in the Wasserstein space.
- The space $(\mathcal{P}(D), W_2)$ is a **positively curved space**: no convexity of W_2^2 nor Dir.

Curvature and convexity

If $\mu, \nu \in \mathcal{P}(D)$, two ways to interpolate.




The **displacement** interpolation



- Midpoint of the geodesic in the Wasserstein space.
- The space $(\mathcal{P}(D), W_2)$ is a **positively curved space**: no convexity of W_2^2 nor Dir.

The **Linear** interpolation



- The Wasserstein distance square W_2^2 and the Dirichlet energy are convex.
- Tools from convex analysis.

The Dirichlet problem

The Dirichlet problem

We choose $\mu_b : \partial\Omega \rightarrow \mathcal{P}(D)$ the boundary data.

Definition

The Dirichlet problem is

$$\min_{\mu} \{ \text{Dir}(\mu) : \mu = \mu_b \text{ on } \partial\Omega \}.$$

The solutions of the Dirichlet problem are called harmonic mappings (valued in the Wasserstein space).

The Dirichlet problem

We choose $\mu_b : \partial\Omega \rightarrow \mathcal{P}(D)$ the boundary data.

Definition

The Dirichlet problem is

$$\min_{\mu} \{ \text{Dir}(\mu) : \mu = \mu_b \text{ on } \partial\Omega \}.$$


The solutions of the Dirichlet problem are called harmonic mappings (valued in the Wasserstein space).

Theorem


Assume $\mu_b : \partial\Omega \rightarrow (\mathcal{P}(D), W_2)$ is a Lipschitz mapping. Then there exists at least one solution to the Dirichlet problem.

Uniqueness is an open question.

Numerics: example



Numerics: example



Numerics: adaptation of Benamou and Brenier¹¹

The Dirichlet problem is a convex optimization problem.

¹¹J.-D. Benamou, and Y. Brenier. *A computational fluid mechanics solution to the Monge-Kantorovich mass transfer problem*. 2000.

Numerics: adaptation of Benamou and Brenier¹¹

The Dirichlet problem is a convex optimization problem.

Unknowns ($\mathbf{m} = \mu \mathbf{v}$ is the momentum):

$$\mu : \Omega \times D \rightarrow \mathbb{R}_+$$

$$\mathbf{m} : \Omega \times D \rightarrow \mathbb{R}^{nd}$$

¹¹J.-D. Benamou, and Y. Brenier. *A computational fluid mechanics solution to the Monge-Kantorovich mass transfer problem*. 2000.

Numerics: adaptation of Benamou and Brenier¹¹

The Dirichlet problem is a convex optimization problem.

Unknowns ($\mathbf{m} = \mu \mathbf{v}$ is the momentum):

$$\mu : \Omega \times D \rightarrow \mathbb{R}_+$$

$$\mathbf{m} : \Omega \times D \rightarrow \mathbb{R}^{nd}$$

Objective:

$$\min_{\mu, \mathbf{m}} \left\{ \iint_{\Omega \times D} \frac{|\mathbf{m}|^2}{2\mu} \right\}$$

under the constraints:

$$\begin{cases} \nabla_{\Omega} \mu + \nabla_D \cdot \mathbf{m} = 0, \\ \mu = \mu_b \text{ on } \partial\Omega. \end{cases}$$

¹¹J.-D. Benamou, and Y. Brenier. *A computational fluid mechanics solution to the Monge-Kantorovich mass transfer problem*. 2000.

Numerics: convergence? (for geodesics $\Omega = [0, 1]$)

In practice: finite-dimensional “approximation” with two convex optimization problems in duality, then **ADMM**.

¹²N. Papadakis, G. Peyré, and E. Oudet. *Optimal transport with proximal splitting*. 2014.

¹³R. Hug, E. Maitre, and N. Papadakis. *On the convergence of augmented Lagrangian method for optimal transport between nonnegative densities*. 2020

¹⁴H. Lavenant. *Unconditional convergence for discretizations of dynamical optimal transport*. 2020.

Numerics: convergence? (for geodesics $\Omega = [0, 1]$)

In practice: finite-dimensional “approximation” with two convex optimization problems in duality, then **ADMM**.

Two kinds of convergence:

- Convergence of the **convex optimization algorithm** to solve the discretized problem^{12 13}.

¹²N. Papadakis, G. Peyré, and E. Oudet. *Optimal transport with proximal splitting*. 2014.

¹³R. Hug, E. Maitre, and N. Papadakis. *On the convergence of augmented Lagrangian method for optimal transport between nonnegative densities*. 2020

¹⁴H. Lavenant. *Unconditional convergence for discretizations of dynamical optimal transport*. 2020.

Numerics: convergence? (for geodesics $\Omega = [0, 1]$)

In practice: finite-dimensional “approximation” with two convex optimization problems in duality, then **ADMM**.

Two kinds of convergence:

- Convergence of the **convex optimization algorithm** to solve the discretized problem^{12 13}.
- Convergence of the solutions of the **discretized problem to the continuous one**¹⁴.

¹²N. Papadakis, G. Peyré, and E. Oudet. *Optimal transport with proximal splitting*. 2014.

¹³R. Hug, E. Maitre, and N. Papadakis. *On the convergence of augmented Lagrangian method for optimal transport between nonnegative densities*. 2020

¹⁴H. Lavenant. *Unconditional convergence for discretizations of dynamical optimal transport*. 2020.

Maximum principle

Maximum principle

Some functionals $F : \mathcal{P}(D) \rightarrow \mathbb{R} \cup \{+\infty\}$ are convex along geodesics, e.g.

$$\mu \rightarrow \int_D \mu(y) \ln(\mu(y)) \, dy.$$

Maximum principle

Some functionals $F : \mathcal{P}(D) \rightarrow \mathbb{R} \cup \{+\infty\}$ are convex along geodesics, e.g.

$$\mu \rightarrow \int_D \mu(y) \ln(\mu(y)) \, dy.$$

Theorem

Take $F : \mathcal{P}(D) \rightarrow \mathbb{R} \cup \{+\infty\}$ convex along generalized geodesics (and few additional regularity property) and a boundary condition $\mu_b : \partial\Omega \rightarrow \mathcal{P}(D)$ such that $\sup_{\partial\Omega} (F \circ \mu_b) < +\infty$.

Maximum principle

Some functionals $F : \mathcal{P}(D) \rightarrow \mathbb{R} \cup \{+\infty\}$ are convex along geodesics, e.g.

$$\mu \rightarrow \int_D \mu(y) \ln(\mu(y)) \, dy.$$

Theorem

Take $F : \mathcal{P}(D) \rightarrow \mathbb{R} \cup \{+\infty\}$ convex along generalized geodesics (and few additional regularity property) and a boundary condition $\mu_b : \partial\Omega \rightarrow \mathcal{P}(D)$ such that $\sup_{\partial\Omega} (F \circ \mu_b) < +\infty$.

Then there exists at least one solution μ of the Dirichlet problem with boundary conditions μ_b such that

$$\Delta(F \circ \mu) \geq 0 \quad \text{and} \quad \operatorname{ess\,sup}_{\Omega} (F \circ \mu) \leq \sup_{\partial\Omega} (F \circ \mu_b).$$

Maximum principle

Some functionals $F : \mathcal{P}(D) \rightarrow \mathbb{R} \cup \{+\infty\}$ are convex along geodesics, e.g.

$$\mu \rightarrow \int_D \mu(y) \ln(\mu(y)) \, dy.$$

Theorem

Take $F : \mathcal{P}(D) \rightarrow \mathbb{R} \cup \{+\infty\}$ convex along generalized geodesics (and few additional regularity property) and a boundary condition $\mu_b : \partial\Omega \rightarrow \mathcal{P}(D)$ such that $\sup_{\partial\Omega} (F \circ \mu_b) < +\infty$.

Then there exists at least one solution μ of the Dirichlet problem with boundary conditions μ_b such that

$$\Delta(F \circ \mu) \geq 0 \quad \text{and} \quad \operatorname{ess\,sup}_{\Omega} (F \circ \mu) \leq \sup_{\partial\Omega} (F \circ \mu_b).$$

Already known for harmonic mappings valued in Riemannian manifolds (Ishihara) and Non Positively Curved spaces (Sturm).

3. An application in nonlinear elasticity

A variational problem inspired from elasticity theory

\mathcal{L}_Ω and \mathcal{L}_D Lebesgue measures restricted to D and Ω respectively.

$$\min_{u:\Omega\rightarrow D} \left\{ E(u) := \int_{\Omega} \left(\frac{1}{2} |\nabla u(x)|^2 - f(x) \cdot u(x) \right) dx : u = g \text{ on } \partial\Omega \text{ and } u\#\mathcal{L}_\Omega = \mathcal{L}_D \right\}$$

- $f: \Omega \rightarrow \mathbb{R}^d$ exterior force.
- $g: \partial\Omega \rightarrow \partial D$ prescribed deformation on the boundary.
- $u\#\mathcal{L}_\Omega = \mathcal{L}_D \Leftrightarrow \forall B \subset D, \mathcal{L}_\Omega(u^{-1}(B)) = \mathcal{L}_D(B)$.

If $d = n$ and u smooth and one-to-one, it's equivalent to

$$|\det \nabla u| = 1.$$

A variational problem inspired from elasticity theory

\mathcal{L}_Ω and \mathcal{L}_D Lebesgue measures restricted to D and Ω respectively.

$$\min_{u:\Omega\rightarrow D} \left\{ E(u) := \int_{\Omega} \left(\frac{1}{2} |\nabla u(x)|^2 - f(x) \cdot u(x) \right) dx : u = g \text{ on } \partial\Omega \text{ and } u\#\mathcal{L}_\Omega = \mathcal{L}_D \right\}$$


- $f: \Omega \rightarrow \mathbb{R}^d$ exterior force.
- $g: \partial\Omega \rightarrow \partial D$ prescribed deformation on the boundary.
- $u\#\mathcal{L}_\Omega = \mathcal{L}_D \Leftrightarrow \forall B \subset D, \mathcal{L}_\Omega(u^{-1}(B)) = \mathcal{L}_D(B)$.

If $d = n$ and u smooth and one-to-one, it's equivalent to

$$|\det \nabla u| = 1.$$

Critical points satisfy $\Delta u + f = (\nabla \omega) \circ u$ in the interior of Ω , where $\omega: D \rightarrow \mathbb{R}$ is a Lagrange multiplier.

An example: pure torsion of a cylinder




$\Omega = D = B(0, 1) \times [0, 1]$. For $a > 0$,

$$u_a \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} R_{az} \begin{pmatrix} x \\ y \end{pmatrix} \\ z \end{pmatrix}$$

where $R_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ rotation by an angle θ .

An example: pure torsion of a cylinder



$\Omega = D = B(0, 1) \times [0, 1]$. For $a > 0$,

$$u_a \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} R_{az} \begin{pmatrix} x \\ y \end{pmatrix} \\ z \end{pmatrix}$$


where $R_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ rotation by an angle θ .

Result ($f \equiv 0$)


For all a , the function u_a is a critical point of the energy.

At least for small a , it is a global minimizer with boundary condition $g = u_a|_{\partial\Omega}$.

Transport plan



Relaxation




Method

$u : \Omega \rightarrow D$ is replaced by $\mu : \Omega \rightarrow \mathcal{P}(D)$.

- By disintegration/fubiniization, one can see a mapping $\mu : \Omega \rightarrow \mathcal{P}(D)$ as $\mu \in \mathcal{P}(\Omega \times D)$ whose first marginal is \mathcal{L}_Ω .

Transport plan



Method

$u : \Omega \rightarrow D$ is replaced by $\mu : \Omega \rightarrow \mathcal{P}(D)$.

- By disintegration/fubiniization, one can see a mapping $\mu : \Omega \rightarrow \mathcal{P}(D)$ as $\mu \in \mathcal{P}(\Omega \times D)$ whose first marginal is \mathcal{L}_Ω .
- The constraint $u\#\mathcal{L}_\Omega = \mathcal{L}_D$ is replaced by the second marginal of μ being \mathcal{L}_D . We write $\mu \in \Pi(\mathcal{L}_\Omega, \mathcal{L}_D)$.
- The marginal constraints are linear. For instance, for all $a \in \mathcal{C}(\Omega)$:

$$\iint_{\Omega \times D} a(x) \mu(dx, dy) = \int_{\Omega} a(x) dx$$

A convex relaxation¹⁵

$$\min_{u:\Omega\rightarrow D} \left\{ \int_{\Omega} \left(\frac{1}{2} |\nabla u(x)|^2 - f(x) \cdot u(x) \right) dx : u = g \text{ on } \partial\Omega \text{ and } u\#\mathcal{L}_{\Omega} = \mathcal{L}_D \right\}$$

\downarrow

$$\min_{\mu:\Omega\rightarrow\mathcal{P}(D)} \left\{ \text{Dir}(\mu) - \iint_{D\times\Omega} f(x) \cdot y \mu(dx, dy) : \right. \\ \left. \mu(x, \cdot) = \delta_{g(x)} \text{ for } x \in \partial\Omega \text{ and } \mu \in \Pi(\mathcal{L}_{\Omega}, \mathcal{L}_D) \right\}$$

Without Dirichlet energy, it's exactly the relaxation used by Yann Brenier in 1987 to prove polar factorization!

¹⁵N. Ghoussoub, Y.-H. Kim, H. Lavenant, A. Z. Palmer. *Hidden convexity in a problem of nonlinear elasticity*. 2020.

A convex relaxation¹⁵

$$\min_{u:\Omega\rightarrow D} \left\{ \int_{\Omega} \left(\frac{1}{2} |\nabla u(x)|^2 - f(x) \cdot u(x) \right) dx : u = g \text{ on } \partial\Omega \text{ and } u \# \mathcal{L}_{\Omega} = \mathcal{L}_D \right\}$$



$$\min_{\mu:\Omega\rightarrow\mathcal{P}(D)} \left\{ \text{Dir}(\mu) - \iint_{D\times\Omega} f(x) \cdot y \mu(dx, dy) : \right. \\ \left. \mu(x, \cdot) = \delta_{g(x)} \text{ for } x \in \partial\Omega \text{ and } \mu \in \Pi(\mathcal{L}_{\Omega}, \mathcal{L}_D) \right\}$$

Without Dirichlet energy, it's exactly the relaxation used by Yann Brenier in 1987 to prove polar factorization!

Remark

It's a convex problem!

¹⁵N. Ghoussoub, Y.-H. Kim, H. Lavenant, A. Z. Palmer. *Hidden convexity in a problem of nonlinear elasticity*. 2020.

Tightness of the relaxation

Let $\lambda_1(\Omega)$ the first eigenvalue of the Dirichlet Laplacian on Ω .

Theorem

Let $u : \Omega \rightarrow D$ a smooth function satisfying $u = g$ on $\partial\Omega$ and $\omega \in C(D)$ such that

$$\Delta u + f = (\nabla \omega) \circ u.$$

If ω can be extended on \mathbb{R}^d in a λ -**convex** function **with** $\lambda > -\lambda_1(\Omega)$ then u is the unique global minimizer global of the energy and and of the relaxed energy.

Tightness of the relaxation

Let $\lambda_1(\Omega)$ the first eigenvalue of the Dirichlet Laplacian on Ω .

Theorem

Let $u : \Omega \rightarrow D$ a smooth function satisfying $u = g$ on $\partial\Omega$ and $\omega \in C(D)$ such that

$$\Delta u + f = (\nabla \omega) \circ u.$$

If ω can be extended on \mathbb{R}^d in a λ -**convex** function **with** $\lambda > -\lambda_1(\Omega)$ then u is the unique global minimizer global of the energy and and of the relaxed energy.

In the pure torsion of the cylinder, global optimality in the case a small. For large a , there holds $\min E_{\text{relaxed}} < \min E$.

Tightness of the relaxation

Let $\lambda_1(\Omega)$ the first eigenvalue of the Dirichlet Laplacian on Ω .

Theorem

Let $u : \Omega \rightarrow D$ a smooth function satisfying $u = g$ on $\partial\Omega$ and $\omega \in C(D)$ such that

$$\Delta u + f = (\nabla \omega) \circ u.$$

If ω can be extended on \mathbb{R}^d in a λ -**convex** function **with** $\lambda > -\lambda_1(\Omega)$ then u is the unique global minimizer global of the energy and and of the relaxed energy.

In the pure torsion of the cylinder, global optimality in the case a small. For large a , there holds $\min E_{\text{relaxed}} < \min E$.

Remark

There exists a simpler proof of this consequence which does not rely on the convex relaxation.

Conclusion

- Working on optimal transport by studying curves and mappings valued in the space of probability distributions.
- Some promising directions in Data Science using optimal transport as a regularizer.
- Definition of harmonic mappings valued in the Wasserstein space, with applications in nonlinear elasticity.

Conclusion

- Working on optimal transport by studying curves and mappings valued in the space of probability distributions.
- Some promising directions in Data Science using optimal transport as a regularizer.
- Definition of harmonic mappings valued in the Wasserstein space, with applications in nonlinear elasticity.

Thank you for your attention