

[Journals A-Z](#) > [Archives of General Psychiatry](#) > [55\(8\) August 1998](#) > Nortriptyline and Cognitive-Behavioral Therapy in the Treatment of Cigarette Smoking.

### Archives of General Psychiatry

Issue: Volume 55(8), August 1998, pp 683-690

Copyright: Copyright 1998 by the American Medical Association. All Rights Reserved. Applicable FARS/DFARS

Restrictions Apply to Government Use. American Medical Association, 515 N. State St, Chicago, IL 60610.

Publication Type: [Original Article]

ISSN: 0003-990X

Accession: 00000756-199808000-00001

Keywords: Behavior Therapy, Cognitive Therapy, Depressive Disorder, Nortriptyline, Smoking Cessation

[Original Article]

## Nortriptyline and Cognitive-Behavioral Therapy in the Treatment of Cigarette Smoking

Hall, Sharon M. PhD; Reus, Victor I. MD; Munoz, Ricardo F. PhD; Sees, Karen L. DO; Humfleet, Gary PhD; Hartz, Diane T. MA; Frederick, Sydney PhD; Triffleman, Elisa MD

### Author Information

From the Department of Psychiatry University of California, San Francisco (Drs Hall, Reus, Munoz, Sees, Humfleet, Frederick, and Triffleman and Ms Hartz); and the San Francisco Veterans Affairs Medical Center Psychiatry Service, San Francisco (Drs Hall, Sees, and Triffleman). Dr Triffleman is now at the Department of Psychiatry, Yale University, New Haven, Conn.

### Abstract

**Background:** A history of major depressive disorder (MDD) predicts failure to quit smoking. We determined the effect of nortriptyline hydrochloride and cognitive-behavioral therapy on smoking treatment outcome in smokers with a history of MDD. The study also addressed the effects of diagnosis and treatment condition on dysphoria after quitting smoking and the effects of dysphoria on abstinence.

**Methods:** This was a 2 (nortriptyline vs placebo) x 2 (cognitive-behavioral therapy vs control) x 2 (history of MDD vs no history) randomized trial. The participants were 199 cigarette smokers. The outcome measures were biologically verified abstinence from cigarettes at weeks 12, 24, 38, and 64. Mood, withdrawal, and depression were measured at 3, 5, and 8 days after the smoking quit date.

**Results:** Nortriptyline produced higher abstinence rates than placebo, independent of depression history. Cognitive-behavioral therapy was more effective for participants with a history of depression. Nortriptyline alleviated a negative affect occurring after smoking cessation. Increases in the level of negative affect from baseline to 3 days after the smoking quit date predicted abstinence at later assessments for MDD history-negative smokers. There was also a sex-by-depression history interaction; MDD history-positive women were less likely to be abstinent than MDD history-negative women, but depression history did not predict abstinence for men.

**Conclusions:** Nortriptyline is a promising adjunct for smoking cessation. Smokers with a history of depression are aided by more intensive psychosocial treatments. Mood and diagnosis interact to predict relapse. Increases in negative affect after quitting smoking are attenuated by nortriptyline.

Arch Gen Psychiatry. 1998;55:683-690

THE LINK between cigarette smoking and depressed mood is well established in population studies. [1-4] There is also evidence that diagnosable depression, particularly major depressive disorder (MDD), and a diagnosis of nicotine dependence are correlated in the general population. [5] Individuals with current or historical depressive disorders seem to be overrepresented among those seeking smoking treatment. [6,7] Smokers with a history of MDD may experience more severe mood-related symptoms when they quit smoking. [8,9]

This study investigated the efficacy of antidepressant therapy as smoking treatment for MDD history-positive smokers. Antidepressants that have been studied in smoking cessation include imipramine hydrochloride, [10] doxepin hydrochloride, [11] bupropion hydrochloride, [12,13] and fluoxetine hydrochloride. [14] The results of imipramine [10] and doxepin [11] studies are mixed and difficult to interpret because of methodological problems. Bupropion has shown short-term efficacy in small trials. [12,13] Recently, a large-scale pharmaceutical industry-sponsored study was sufficiently convincing that the drug received a Federal Drug Administration indication for the treatment of cigarette smoking. [15] Results from multisite pharmaceutical industry-sponsored studies of fluoxetine have not been published, but a secondary analysis of part of that data set suggested that participants reporting depressive symptoms at study enrollment showed significantly improved cessation rates with the administration of active drug. [14]

Nortriptyline hydrochloride has adrenergic activity; therefore, it might be useful in smoking cessation. Nortriptyline is an inexpensive generic drug. Also, therapeutic serum levels, at least for the treatment of depression, are known (190-532 nmol/L), [16] thus providing initial treatment parameters.

One study by Hall and colleagues [7] found that smokers with a history of depression were more likely to be abstinent if treated with 10 sessions of cognitive-behavioral (CB) therapy than if treated with 5 sessions of health education therapy. Zelman and colleagues [17] found that dysphoric smokers were helped by a supportive treatment to a greater extent than nondysphoric smokers. Hall and colleagues, [18] using a control that was equivalent in contact time to cognitive-behavioral therapy, failed to replicate their original findings; one purpose of the proposed study was an attempt to recapture the findings of the original study. Thus, we compared the 10-session CB intervention for smoking cessation with the 5-session health education (HE) control intervention used in the original study.

The present study was a 2 x 2 x 2 factorial design in which medication (nortriptyline vs placebo) was crossed with CB vs HE intervention and with presence vs absence of a history of MDD. We hypothesized the following: (1) Participants with a history of MDD would show higher abstinence rates with the administration of nortriptyline than with the administration of placebo. No differences were hypothesized for MDD history-negative participants. (2) Participants with a history of MDD would show higher abstinence rates with the CB intervention than with the HE intervention. No differences were hypothesized for MDD history-negative smokers. (3) Major depressive disorder history-positive participants would show greater increases in dysphoria while quitting smoking than MDD history-negative participants. (4) Nortriptyline would be more effective than placebo in reducing the dysphoria occurring after quitting smoking in MDD history-positive participants only, with no difference predicted for MDD history-negative participants. (5) Independent of MDD history, participants with smaller increases in dysphoria immediately after quitting smoking would be more likely to be abstinent at later assessments than participants with greater increases in dysphoria after quitting smoking.

## PARTICIPANTS AND METHODS

### PARTICIPANTS

This study was approved by the Institutional Review Board of the University of California, San Francisco. Participants were recruited by public service and newspaper announcements. After giving written, informed consent, 248 individuals completed the Diagnostic Interview Schedule, [19] and their conditions were evaluated for study enrollment. Three withdrew for personal reasons before random assignment. We excluded 46 patients: 27 because of electrocardiographic abnormalities, 11 because of a current MDD, and 8 for other reasons. Thus, 199 participants were randomly assigned to 1 of 2 psychological treatments, CB vs HE therapy, and to 1 of 2 drug treatments, active nortriptyline vs placebo. They were men (n=89) and women (n=110), aged 21 to 65 years, who smoked 10 or more cigarettes per day, and who wanted to quit smoking.

Individuals who met the criteria for MDD within 3 months of the baseline were excluded and referred for depression treatment. Individuals who were taking prescribed psychotropic drugs or who showed evidence of alcohol or other nonnicotine drug use were excluded. Individuals who had taken a monoamine oxidase inhibitor within the last 2 weeks were excluded even if they were not currently taking that drug. A history of MDD was diagnosed in 65 (32.7%) of the participants. Fifty participants reported recurrent episodes.

## PROCEDURES

Nortriptyline was dispensed from week 1 through week 12. Psychological treatment occurred from week 4 through week 12. The smoking quit date was during week 5 so that therapeutic nortriptyline levels could be reached.

Participants were assessed at baseline on demographic variables, mood, depressive and withdrawal symptoms, and current and historical MDD. Additional assessments were held at weeks 12, 24, 38, and 64. Psychological measures were completed and smoking status was determined at each assessment. Participants were asked if they had smoked at all, even a puff, during the last week. If they had not, and their expired air carbon monoxide level was 10 ppm or less, their urine sample was analyzed for cotinine. If cotinine levels were 341 nmol/L or less, participants were coded as abstinent. In addition to serum samples for nortriptyline levels, compliance was assessed by capsule counts (capsules allotted-capsules returned). We determined side effects by a checklist administered at baseline and at weeks 1 through 4 and 6.

Participants completed the Profile of Mood States (POMS) [20] and a withdrawal scale [21] at 3, 5, and 8 days after their smoking quit date. The Beck Depression Inventory [22] was completed 8 days after the smoking quit date.

Participants were stratified on MDD history vs no MDD history and on number of cigarettes smoked and randomly assigned from within stratification blocks to the 4 experimental conditions using a computerized randomization program.

## MEASURES

Bachelor's and master's level survey workers, trained by one of us (R.F.M.), administered the computerized Diagnostic Interview Schedule. [23] Based on the Diagnostic Interview Schedule, participants were classified as MDD history positive or MDD history negative. Negative affect was assessed using the POMS total mood disturbance (TMD) score. We assessed depressive symptoms during the past week by the Beck Depression Inventory and trait anxiety by the State-Trait Anxiety Inventory Trait scale. [24]

We administered 2 smoking scales, the Shiffman Tobacco Withdrawal Scale, [21] a widely used scale that shows changes in scores as a function of time since last cigarette smoked and differentiates between partial and total abstinence, and the Fagerstrom Tolerance Questionnaire, [25,26] a measure of tobacco dependence. High scores predict smoking treatment failure and correlate with physiological smoking-induced changes.

## TREATMENT INTERVENTIONS

## Drug Treatment

Medication was placebo controlled and double blind. Placebo and active drug were identical in appearance. Participants met with study physicians at week 1 to begin the administration of medication and at weeks 2 and 3 for the administration of medication and a review of the side effects. The drug dose was titrated to therapeutic levels for depression. Participants received nortriptyline hydrochloride, 25 mg/d, for 3 days; the dose was then increased to 50 mg/d for 4 days. Serum levels were assessed at week 2; the dose was increased to 75 mg/d if a therapeutic level had not been reached. Serum levels were reassessed at week 4, and the dose increased to 100 mg/d, if necessary, and at week 6, when the dosage was titrated downward, if necessary. The modal dose was 100 mg/d. Whenever titration occurred for a participant receiving active drug, the dose was titrated for a participant receiving placebo in the same cohort. Participants continued to receive the maintenance dose of medication until week 12, and the medication was tapered during week 13.

## Psychological Treatment

**CB Treatment.** The central premise of the CB treatment, described in detail elsewhere, [7] was that thoughts, activities, and mood interact to influence smoking. Sessions focused on mood management skills to manage dysphoria and maintain nonsmoking, including methods to increase the frequency of pleasant activities and decrease relapse-related thoughts, by thought stopping, increasing pleasant thoughts, and rational-emotive techniques. Techniques for increasing positive social contacts, decreasing negative contacts, and improving relationships were developed.

Participants met for ten 2-hour group sessions (with 5-11 members) for 8 weeks. The therapist manual is available (S.M.H., R.F.M., and K. Organista, PhD, unpublished data, 1992).

**HE Intervention.** The HE intervention, also described in detail elsewhere, [7] provided health-related information and facilitated discussion. A core element was the development of a plan to quit smoking and weekly modification of that plan. Methods used included paper-and-pencil exercises, informational handouts, brief didactic presentations, homework assignments, and smoking monitoring. Participants met for five 90-minute sessions (with 5-11 members) for 8 weeks.

## PROVIDERS

The therapists were 3 doctoral-level clinical psychologists. All had experience in CB and psychoeducational interventions. All therapists treated participants in each of the 4 experimental conditions. Therapists were supervised weekly by two of us (R.F.M. and G.H.). Adherence to the protocol was monitored; if situations arose that were not in the protocol, audiotapes were reviewed and a standard response was developed. Medication was prescribed by 1 of 2 psychiatric residents, who were supervised by two of us (V.I.R. and K.L.S.).

## STATISTICAL METHODS

The principal data analysis method was a generalized linear model (GLM), a generalization of the classic linear model estimated using likelihood functions instead of least squares. [27] Generalized linear models allow use of repeated measurements when there is missing data, without excluding participants with data missing or imputing values. [28,29] We used a computer program (PROC Mixed, SAS Institute Inc, Cary, NC) to estimate and test the GLM. When abstinence was the dependent variable, we also used another computer program (GLMMIX Macro, SAS Institute Inc) [30] that interacts with the first program mentioned (PROC Mixed) to modify it so that it is appropriate for dichotomous data. To evaluate the 3 hypotheses (hypotheses 1, 2, and 5) that predicted abstinence, abstinence status at weeks 12, 24, 38, and 64 were the dependent variables. Hypotheses 1 and 2 were tested by a 2 (active vs placebo drug) by 2 (CB vs HE intervention) x 2 (MDD vs no MDD history) model. Hypothesis 5 was tested by a model that included the baseline POMS-TMD score, the day 3 POMS-TMD score, and the 3 design variables (drug, psychological treatment condition, and diagnostic status) and their interaction as independent variables.

For hypotheses 3 and 4, which predicted changes in dysphoria after the quit date, we completed 3 parallel analyses. In the first analysis, the dependent variables were the POMS-TMD scores at baseline and at days 3, 5, and 8 after the smoking quit date. In the second analysis, withdrawal scores replaced POMS scores. In the third analysis, dependent variables were Beck Depression Inventory scores at baseline and at day 8. The 3 design variables and their interaction were the independent variables.

To provide a comparison with recent antidepressant studies, [15] we present continuous abstinence rates (abstinent at each assessment) and use a chi squared test to evaluate this univariate variable.

We evaluated the effects of sex and its interaction with the 3 design variables on abstinence rates at weeks 12 through 64 using 3 GLM models computed using a macro (GLMMIX). A parallel model was used to determine therapist effects. The effects of nortriptyline levels and capsules ingested on abstinence rates at the 4 assessments were evaluated by 2 GLM analyses. For nortriptyline, the independent variables were serum level, diagnostic category, and their interaction; for capsules, the independent variables were active drug vs placebo, capsule count, and their interaction. For GLM analyses, if the overall effect was significant, post hoc contrasts were completed using a computer program (PROC Mixed) and, if appropriate, a macro (GLMMIX). Effect sizes are expressed as odds ratios (ORs) and confidence intervals (CIs).

We used Student t tests and chi squared tests to determine whether there were differences between participants with and participants without missing data at assessments on baseline demographic, smoking, mood, and psychiatric variables ( $P > .05$ , all comparisons).

Analysis of variance and chi squared tests were used to evaluate baseline differences among treatment conditions. A chi squared test was used to evaluate the rate of occurrence of side effects. The tests were 2 tailed, with  $P < .05$  for all comparisons.

## RESULTS

### PARTICIPANT CHARACTERISTICS

The demographic, smoking, and psychiatric characteristics of the participants in each experimental group are given in (Table 1).

Variables	Drug Condition/Psychological Treatment			
	Active Drug/CI Intervention (n = 51)	Active Drug/HI Intervention (n = 48)	Placebo Drug/CI Intervention (n = 52)	Placebo Drug/HI Intervention (n = 48)
<b>Demographic</b>				
Age†	41.7 (8.4)	40.7 (8.5)	40.0 (8.8)	39.4 (8.7)
Sex (female)‡	22 (43)	20 (58)	30 (58)	30 (63)
Race (white)‡	43 (86)	41 (87)	45 (87)	42 (88)
Married or with a partner (yes)‡	22 (43)	16 (34)	19 (37)	25 (52)
<b>Smoking†</b>				
No. of cigarettes smoked per day	22.1 (11.9)	24.9 (12.7)	21.1 (7.6)	22.5 (8.5)
Fagerstrom Tolerance Scale score	5.6 (2.1)	5.6 (2.2)	5.4 (2.3)	5.5 (2.1)
Years as regular smoker	23.0 (10.7)	22.2 (10.3)	21.7 (10.0)	21.7 (10.4)
Carbon monoxide level, ppm	24.0 (10.2)	25.3 (11.9)	25.6 (12.0)	24.7 (12.4)
Prior quit attempts	3.0 (1.2)	2.6 (1.4)	2.8 (1.2)	2.5 (1.5)
<b>Psychiatric</b>				
Beck Depression Inventory score‡	8.6 (7.2)	9.5 (7.5)	7.7 (6.5)	6.7 (6.8)
FCMS-TMD‡	30.7 (28.1)	31.4 (32.5)	22.6 (30.5)	32.4 (32.2)
Shiffman Withdrawal Scale score‡	50.4 (11.1)	55.5 (10.0)	55.5 (11.5)	56.0 (12.3)
Spiegelberger State Anxiety Score†	40.7 (11.5)	39.3 (11.3)	39.0 (10.5)	41.9 (10.5)
Confidence Scale total score‡	47.1 (14.8)	48.8 (15.5)	43.8 (14.9)	46.9 (16.6)
History of MDD‡	17 (33.3)	15 (31.3)	17 (32.7)	15 (31.3)

\* There were no significant differences among treatment conditions, except that psychological treatment conditions differed in the presence or absence of previous quit attempts ( $\chi^2$ ,  $df = 1$ ,  $P = .007$ ) and drug condition on baseline withdrawal scores ( $t_{1,90} = 1.9$ ,  $P = .03$ ). Neither potential confounding variable correlated significantly with abstinence rates at any assessment. CI indicates cognitive-behavioral HI, health education; FCMS, Profile of Mood States; TMD, total mood disturbance; and MDD, major depressive disorder.

† Data are given as the mean (SD).

‡ Data are given as the number (percentage) of subjects.

Table 1. Data for Variables by Psychological Treatment and Drug Condition\*

The demographic, smoking, and psychiatric characteristics of MDD history-positive and MDD history-negative participants are given in (Table 2).



Variables	MDD History–Negative Participants (n = 134)	MDD History–Positive Participants (n = 65)
Demographic		
Age*	40.5 (9.5)	41.1 (9.9)
Sex (female)†	68 (49)	42 (64)
Race (white)†	114 (87)	59 (89)
Married or with a partner (yes)†‡	66 (50)	16 (24)
Smoking		
No. of cigarettes smoked per day*	23.0 (10.2)	21.8 (10.4)
Fagerstrom Tolerance Scale score*	5.5 (2.1)	5.5 (2.2)
Years a regular smoker*	22.0 (9.7)	22.2 (11.4)
Carbon monoxide level, ppm*	24.2 (10.6)	26.0 (13.7)
Prior quit attempts†	121 (92)	60 (92)
Psychiatric*		
Beck Depression Inventory score‡	7.2 (5.6)	12.1 (8.3)
Profile of Mood States total mood disturbance	21.4 (29.0)	45.9 (36.8)
Shiffman Withdrawal Scale score	54.0 (11.6)	56.1 (11.7)
Speilberger State Anxiety score	38.7 (10.4)	39.3 (8.4)
Confidence Scale total score	46.6 (15.2)	46.4 (15.5)

\*Data are given as the mean (SD).

†Data are given as the number (percentage) of subjects.

‡There were no significant differences, except that MDD history–positive participants were less likely to be married or have a partner than MDD history–negative participants ( $\chi^2$ , [N = 197] = 11.6,  $P < .001$ ) and to have higher Profile of Mood States total mood disturbance ( $t_{195} = 4.90$ ,  $P < .001$ ) and Beck Depression Inventory ( $t_{190} = 5.03$ ,  $P < .001$ ) scores.

Table 2. Baseline Mood- and Withdrawal-Related Variables for Participants With and Without a History of Major Depressive Disorder (MDD)\*

## ATTRITION

Forty-seven (24%) of the participants reported that they did not want to complete treatment. Of these participants, 23 dropped out because of personal reasons, 5 dropped out because of medication side effects (4 in the active drug group and 1 in the placebo group), and 11 dropped out because of program issues (eg, dislike of the group format); 1 participant moved, 1 had an unrelated medical problem, and 6 gave no reason for dropping out. There were no differences between psychological interventions ( $\chi^2_1 [N=199]=1.51, P=.22$ ) or between diagnostic categories ( $\chi^2_1 [N=199]=.63$ ) in dropout rates, but participants in the placebo group had a higher overall dropout rate (30%) than did participants in the active drug group (17%) ( $\chi^2_1 [N=199]=4.54, P=.04$ ; OR, 2.07; 95% CI, 1.05-4.06).

The no-show rates at each assessment were as follow: week 12, 19%; week 24, 17%; week 38, 20%; and week 64, 17%. There were no significant differences in the number of assessments missed for any of the design variables (for the psychological treatment condition,  $t_{197} < .42$ ; for the drug treatment condition,  $t_{197} = 1.63, P=.25$ ; and for the diagnosis,  $t_{197} = .72$ ).

## ABSTINENCE RATES

The main effect for the drug was significant ( $\chi^2_1 [N=199]=4.34, P=.04$ ; OR, 2.42; 95% CI, 1.75-3.35), as was the treatment condition x diagnosis interaction ( $\chi^2_1 [N=199]=3.84, P=.05$ ). The main effect of assessment (at weeks 12, 24, 38, and 64) was also significant ( $\chi^2_3 [N=199]=10.33, P=.02$ ; OR, 0.78; 95% CI, 0.68-0.90), indicating a decline in abstinence rates for both conditions from week 12 to week 64. There were no other significant effects ( $P < .05$ ), including the hypothesized diagnosis x drug interaction ( $\chi^2_1 [N=199] < .92$ ), the psychological treatment condition x drug interaction ( $\chi^2_1 [N=199] < .98$ ), or the diagnosis x psychological treatment condition x drug interaction ( $\chi^2_1 [N=199] < .59$ ). The abstinence rates at each assessment are shown in [\(Figure 1\)](#).






Figure 1. Abstinence rates across time for participants receiving active nortriptyline hydrochloride (n=99) and placebo (n=100). The effect of drug is significant across weeks 12 through 64 whether the dependent variable is abstinence at each assessment ( $P=.04$ ) or continuous abstinence for the year ( $P=.02$ ). There were no significant main or interaction effects for diagnosis ( $P>.05$ , all comparisons).

In the active drug condition, 24% of the participants achieved continuous abstinence; in the placebo drug condition, 12% achieved continuous abstinence ( $\chi^2_1 [N=199]=5.03$ ,  $P=.02$ ; OR, 2.3; 95% CI, 1.1-5.0). We concluded that there was evidence of superiority of nortriptyline vs placebo for cigarette smokers, for MDD history-negative and MDD history-positive smokers.

The abstinence rates by psychological treatment condition and diagnostic grouping are shown in (Figure 2). As (Figure 2) indicates, MDD history-positive participants assigned to the CB intervention achieved abstinence rates comparable with the MDD history-negative participants, but MDD history-positive subjects assigned to the HE intervention were less likely to be abstinent than those assigned to the CB intervention.




Figure 2. Abstinence rates across time for those in the cognitive-behavioral (CB) (n=100) vs health education (HE) (n=99) intervention by major depressive disorder (MDD) history-negative (n=134) vs MDD history-positive (n=65) diagnosis. The significant interaction effect ( $P=.05$ ) reflects a significantly poorer abstinence rate for MDD history-positive subjects (n=31) than for MDD history-negative subjects (n=68) in the HE intervention group ( $\chi^2_{1,68}=6.44$ ,  $P=.01$ ; OR, 1.70; 95% CI, 1.00-2.98) and no significant difference ( $P=.91$ ) between MDD history-positive (n=34) and MDD history-negative (n=66) participants in the CB intervention group.

We concluded that we achieved support for the second hypothesis, that participants with a history of MDD would be more likely to benefit from CB therapy than would participants without a history of MDD. Abstinence rates by diagnosis, psychological treatment condition, and active vs placebo drug are given in (Table 3).

Characteristics	Week			
	12	24	38	64
MDD history—negative participants				
Health education intervention group				
Receiving placebo (n = 32)	<b>36</b> (31) 28	<b>30</b> (25) 27	<b>33</b> (25) 24	<b>27</b> (22) 26
Receiving active drug (n = 33)	<b>74</b> (61) 27	<b>46</b> (39) 28	<b>48</b> (42) 29	<b>44</b> (36) 27
Cognitive behavioral intervention group				
Receiving placebo (n = 35)	<b>30</b> (20) 23	<b>31</b> (23) 26	<b>25</b> (20) 28	<b>24</b> (20) 29
Receiving active drug (n = 34)	<b>66</b> (56) 29	<b>61</b> (50) 28	<b>39</b> (32) 28	<b>41</b> (35) 29
MDD history—positive participants				
Health education intervention group				
Receiving placebo (n = 16)	<b>21</b> (19) 14	<b>15</b> (13) 13	<b>20</b> (13) 10	<b>15</b> (13) 13
Receiving active drug (n = 15)	<b>47</b> (47) 15	<b>33</b> (33) 15	<b>46</b> (40) 13	<b>23</b> (20) 13
Cognitive behavioral intervention group				
Receiving placebo (n = 17)	<b>54</b> (41) 13	<b>21</b> (18) 14	<b>27</b> (24) 15	<b>36</b> (29) 14
Receiving active drug (n = 17)	<b>62</b> (47) 13	<b>50</b> (41) 14	<b>46</b> (35) 13	<b>29</b> (24) 14

*\*Abstinence rates computed with missing data omitted are in boldface. Abstinence rates with missing data coded or indicative of smoking are in parentheses. The sample size in the first column indicates the number of participants originally assigned to the condition; the columns for weeks 12, 24, 38, and 64 are the number of subjects assessed for the reference cells at that point. MDD indicates major depressive disorder.*

Table 3. Abstinence Rates at All Assessments by Baseline Diagnosis, Psychological Treatment, and Drug Dose\*

Abstinence rates did not differ because of therapist, the interaction of therapist with drug, psychological treatment, or drug x psychological treatment condition ( $P > .05$ , all comparisons).

## PREDICTION OF ABSTINENCE BY NORTRIPTYLINE SERUM LEVELS AND NET CAPSULE COUNT

Nortriptyline serum levels for participants who were assigned to receive active nortriptyline and who were abstinent the week before the blood draw predicted abstinence at weeks 12, 24, 38, and 64 ( $\chi^2_1$  [N=86]=14.28,  $P < .001$ ; OR, 1.02; 95% CI, 1.01-1.03). Abstinent participants were selected because smoking is known to result in lowered nortriptyline levels. [31] Of the participants who received nortriptyline for whom we were able to obtain serum samples at week 6, 44 (51%) achieved serum levels considered therapeutic for the treatment of MDD. Mean week 6 nortriptyline serum levels for abstinent and smoking subjects by week are given in (Table 4).

Partici- pant Status	Week			
	12	24	38	64
Relapsed	132.2 (101.5)	151.6 (120.8)	162.3 (107.5)	166.1 (116.7)
Abstinent	209.8 (116.3)	218.9 (108.3)	216.6 (112.1)	219.6 (117.8)

*\* Data are given as the mean (SD) level of nortriptyline (given as nanomoles per liter). The maintenance nortriptyline serum level predicts abstinence at later assessments ( $P < .001$ ). Nortriptyline was given as nortriptyline hydrochloride.*

Table 4. Maintenance Nortriptyline Serum Levels by Abstinence Status at Each Assessment for 86 Participants Receiving Active Nortriptyline Who Provided Blood Samples at Week 6\*

Capsules ingested did not differ between conditions ( $\chi^2_1$  [N=199]=1.43,  $P = .23$ ) nor did capsules interact with drug ( $\chi^2_1$  [N=199]<.68). Controlling for these 2 variables, drug (abstinent vs placebo) remained significant ( $\chi^2_1$  [N=199]=4.32,  $P = .04$ ; OR, 3.03; 95% CI, 1.07-8.57).

## SIDE EFFECTS

Of the 14 potential side effects (dry mouth, rash, weight gain, lightheadedness, shaky hands, constipation, blurry vision, sexual problems, difficulty in urinating, racing heart, swollen legs, chest pain or pressure, shortness of breath, and "other"), postbaseline endorsement rates differed between drug condition for 4: (1) dry mouth (active=78%; placebo=33%;  $\chi^2_1$  [N=198]=39.6,  $P < .001$ ; OR, 7.00; 95% CI, 3.73-13.17); (2) lightheadedness (active, 49%; placebo, 22%;  $\chi^2_1$  [N=198]=16.01,  $P < .001$ ; OR, 2.42; 95% CI, 1.85-6.35); (3) shaky hands (active=23%; placebo=11%;  $\chi^2_1$  [N=198]=5.11,  $P = .02$ ; OR, 2.42; 95% CI, 1.11-5.29); and (4) blurry vision (active=16%; placebo=6%;  $\chi^2_1$  [N=198]=5.11,  $P = .02$ ; OR, 3.00; 95% CI, 1.12-7.99).

## SEX EFFECTS

Sex interacted with diagnostic category ( $\chi^2_1$  [N=199]=9.08,  $P=.002$ ), not with drug ( $\chi^2_1$  [N=199]=3.33,  $P=.07$ ) or psychological treatment condition ( $\chi^2_1$  [N=199]<1,  $P=.93$ ). Post hoc analyses indicated poorer abstinence rates for MDD history-positive women than for MDD history-negative women ( $\chi^2_{1,110}=9.82$ ,  $P=.002$ ; OR, 2.05; 95% CI, 1.32-3.23) but not for MDD history-positive men ( $\chi^2_{1,89}=1.66$ ,  $P=.20$ ). For women, the abstinence rates were as follow: week 12, MDD history positive=38%, MDD history negative=53%; week 24, MDD history positive=22%, MDD history negative=47%; week 38, MDD history positive=29%, MDD history negative=39%; and week 64, MDD history positive=20%, MDD history negative=37%. For men, the abstinence rates were as follow: week 12, MDD history positive=61%, MDD history negative=52%; week 24, MDD history positive=45%, MDD history negative=37%; week 38, MDD history positive=47%, MDD history negative=36%; and week 64, MDD history positive=37%, MDD history negative=31%.

## CHANGES IN NEGATIVE AFFECTIVE STATES AS A FUNCTION OF DIAGNOSTIC AND DRUG CONDITION

For POMS-TMD score, there was a significant drug (active vs placebo) by assessment (baseline and days 3, 5, and 8) interaction effect ( $F_{3,359}=3.30$ ,  $P=.02$ ). As (Figure 3) shows, POMS-TMD scores for participants given placebo increased from baseline to day 3, while the scores for those given active drug decreased. There was also a significant main effect for assessment ( $F_{3,359}=6.77$ ,  $P=.001$ .) The hypothesized diagnostic status by assessment interaction failed to meet traditional levels of significance ( $F_{3,359}=2.47$ ,  $P=.06$ ). The POMS-TMD scores tended to decrease over time for MDD history-positive participants from baseline to day 8 (baseline mean=45.93, SD=36.83; day 3 mean=38.36, SD=36.08; day 5 mean=24.71, SD=26.85; and day 8 mean=22.72, SD=33.52). For MDD history-negative participants, the POMS-TMD scores tended to increase from baseline to day 3 (baseline mean=21.37, SD=29.03; day 3 mean=28.11, SD=33.37) and decrease thereafter (day 5 mean=17.67, SD=28.03; day 8 mean=15.53, SD=27.39). There were no significant effects for Beck Depression Inventory ( $P>.05$  for all) or total withdrawal symptoms ( $P>.05$  for all).




Figure 3. Profile of Mood States Total Mood Disturbance scores for participants receiving active nortriptyline hydrochloride and placebo at baseline ( $n=192$ ) and at days 3 ( $n=139$ ), 5 ( $n=119$ ), and 8 ( $n=130$ ) after the smoking quit date. The significant drug  $\times$  assessment interaction ( $P=.02$ ) reflects a downward slope for participants receiving active drug ( $F_{3,181}=8.38$ ,  $P=.001$ ) and an upward slope for participants receiving placebo ( $F_{3,178}=2.55$ ,  $P=.06$ ).

### PREDICTIONS OF ABSTINENCE BY POSTQUIT MOOD CHANGES

We hypothesized that increases in the POMS-TMD score occurring from baseline to the smoking quit date would predict lower rates of abstinence. The day 3 score was used because it showed the greatest change from baseline. Controlling for the baseline POMS-TMD score, the day 3 score interacted with diagnostic history ( $\chi^2_1 [N=199]=5.26$ ,  $P=.02$ ). The POMS-TMD scores at baseline and at day 3 by diagnostic condition for subjects smoking and abstinent at weeks 12 and 64 are given in (Table 5). Only 2 weeks were selected for clarity of presentation. Data at weeks 24 and 38 show similar patterns and magnitude of change.



Major Depressive Disorder History	Abstinent Participants	Smoking Participants
Negative (n = 134)		
Week 12	(n = 47)	(n = 38)
Baseline	21.4 (30.2)	19.5 (26.3)
Day 3	26.3 (31.4)	28.4 (32.3)
Week 64	(n = 31)	(n = 53)
Baseline	19.2 (29.4)	20.9 (28.0)
Day 3	24.1 (26.4)	30.6 (37.4)
Positive (n = 65)		
Week 12	(n = 19)	(n = 23)
Baseline	42.4 (33.8)	50.8 (38.2)
Day 3	32.3 (40.2)	45.8 (32.5)
Week 64	(n = 10)	(n = 31)
Baseline	58.6 (46.0)	42.3 (34.4)
Day 3	20.0 (34.3)	43.5 (32.3)

*\*Data are given as the mean (SD) at baseline and day 3. POMS indicates Profile of Mood States; TMD, total mood disturbance. Major depressive disorder history-negative participants showed increased POMS-TMD scores from baseline to day 3, and the magnitude of this increase predicted smoking at weeks 12 through 64 ( $\chi^2_{1,134} = 5.48$ ,  $P = .02$ ; OR, 1.01; 95% confidence interval, 1.00-1.01). Major depressive disorder history-positive participants showed decreased POMS-TMD scores for baseline to day 3, but the magnitude of the decrease was not a significant predictor of smoking status ( $\chi^2_{1,65} = 2.26$ ,  $P = .13$ ).*

Table 5. POMS-TMD Scores for Participants With and Without a History of Major Depressive Disorder, Smoking vs Abstinent, Weeks 12 and 64\*

As (Table 5) indicates, the postquit increase in poor mood predicted abstinence for MDD history-negative participants only.

## COMMENT

The outcome data suggest that nortriptyline is a useful adjunct to smoking cessation efforts. This is supported by the correlation between serum levels and abstinence status. That it reflects more than compliance is indicated by the sustained significance of nortriptyline dose when net capsules ingested for active and placebo treatment were entered into the statistical model. Although there were differences in side effects between the active and placebo groups, they were not perceived as sufficiently troublesome to cause a high dropout rate. In fact, the dropout rate for participants in the active drug group was less than that for participants in the the placebo group.

Industry-sponsored studies have garnered a Federal Drug Administration indication for bupropion. Yet nortriptyline, a generic drug, is less expensive. Contrary to hypotheses, there was not a differential abstinence effect for smokers with a history of MDD using nortriptyline. These findings parallel those reported by others using bupropion. [13,14] These 2 antidepressant drugs have similar effects on smoking cessation; both have effects on noradrenergic transmission, although effects on other neurotransmitter systems may differ between the 2 drugs. Independent of diagnosis, nortriptyline seems to have alleviated the increases in poor mood that occurred the first few days after smoking cessation. This finding has implications for the use of antidepressants to ameliorate the negative mood that follows smoking cessation in non-clinically depressed patients, as well as for use of the drug for instances of poor mood that are not part of a diagnosable disorder. It could be argued that differences in rates of smoking cessation among the 4 experimental cells could have confounded the observed differences in mood between the active and placebo drug conditions. We did not collect smoking data on days 3, 5, and 98, so we cannot definitely refute this argument. However, the effects observed are not easily explainable by possible differences in cessation rates. Dysphoria decreased from baseline to day 3 in participants in the nortriptyline treatment group and increased in participants in the placebo group. Increases in dysphoria are frequently noted after quitting smoking [8,9]; therefore, the observed differences between the drug conditions are not likely to be due to confounding by smoking quit rates.

We replicated the original findings that MDD history-positive participants are differentially helped by a 10-session CB intervention when it is compared with a 5-session HE intervention. [7] These results contrast with a lack of significant enhancement of outcome for MDD history-positive participants when therapeutic contact time was held equivalent in the control condition. [17] Thus, increased psychotherapeutic contact may differentially benefit smokers with a depressive history. This interpretation of the data is further supported by the failure of the psychotherapeutic condition to differentially alter cessation-induced changes in mood.

A history of MDD has predicted poor treatment outcome in some studies [6] but not others. [7,17] Only a few reported sex differences, and lack of analysis for such differences may explain the inconsistencies in the literature. [6,32]

The results do not support the hypothesis that MDD history-positive participants would have a greater increase in negative affect after quitting smoking than MDD history-negative smokers. Indeed, while MDD history-negative smokers did show an increase in poor mood after quitting smoking, MDD history-positive smokers showed a nonsignificant decrease. These data are not consistent with earlier investigations. These earlier investigations are difficult to compare with the current study, because of lack of baseline data, [8] failure to include subjects who successfully quit, [33] or small sample size. [9] However, the failure of current data to agree with the existing literature, even though the literature is imperfect, is troubling, and the current findings should be considered tentative until replicated.

Similarly, increase in the POMS-TMD scores from baseline to 3 days after the quit date predicted failure to maintain abstinence for MDD history-negative participants but not for MDD history-positive participants. For the latter participants, a greater decrease in poor mood from baseline to day 3 was related to abstinence but not significantly so.

There are limitations to the study. First, the outcomes reported do not represent those that could be expected were nortriptyline to be used in a typical medical setting. Psychological intervention in this study provided more contact time and intensity than could be provided by most physicians. Second, this is a volunteer sample of participants who sought treatment and who were more motivated to change than the smokers encountered in physicians' practices. Third, most participants, independent of condition, were smoking at the 64-week follow-up.

Accepted for publication March 24, 1998.

This study was supported by grants R01 DA 23652, P50 DA 09253, and T32 DA 07250 from the National Institute on Drug Abuse, Rockville, Md; and by a Career Research Scientist Award from the Veterans Administration, Washington, DC (Dr Hall).

We thank Scott Walters and Chris Hobart for their help in manuscript preparation and Kevin Delucchi for statistical consultation.

Reprints: Sharon M. Hall, PhD, University of California, San Francisco, Box 0984TRC, Treatment Research Center, 401 Parnassus Ave, San Francisco, CA 94143-0984.

This article is also available on our Web site: [www.ama-assn.org/psych](http://www.ama-assn.org/psych).

## REFERENCES

1. Anda RF, Williamson DF, Escobedo LG, Mast EE, Giovino GA, Remington PL. Depression and the dynamics of smoking: a national perspective. *JAMA*. 1990;264:1541-1545. [SFX](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)
2. Kandel D, Davies M. Adult sequelae of adolescent depressive symptoms. *Arch Gen Psychiatry*. 1986;43:255-262. [SFX](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)
3. Covey LS, Tam D. Depressive mood, the single-parent home, and adolescent cigarette smoking. *Am J Public Health*. 1990;80:1330-1333. [SFX](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)
4. Perez-Stable EJ, Marin G, Marin BV, Katz MH. Depressive symptoms and cigarette smoking among Latinos in San Francisco. *Am J Public Health*. 1990;80:1500-1502. [SFX](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)
5. Breslau N, Kilbey M, Andreski P. Nicotine dependence, major depression, and anxiety in young adults. *Arch Gen Psychiatry*. 1991;48:1069-1074. [SFX](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)
6. Glassman AH, Stetner F, Walsh BT, Raizman PS, Fleiss JL, Cooper TB, Covey LS. Heavy smokers, smoking cessation, and clonidine: results of a double-blind, randomized trial. *JAMA*. 1988;259:2863-2866. [\[Context Link\]](#)
7. Hall SM, Munoz RF, Reus VI. Cognitive-behavioral intervention increases abstinence rates for depressive-history smokers. *J Consult Clin Psychol*. 1994;62:141-146. [\[Context Link\]](#)
8. Covey LS, Glassman AH, Stetner F. Depression and depressive symptoms in smoking cessation. *Compr Psychiatry*. 1990;31:350-354. [SFX](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)
9. Ginsberg D, Hall SM, Reus VI, Munoz RF. Mood and depression diagnosis in smoking cessation. *Exp Clin Psychopharmacol*. 1995;3:389-395. [SFX](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)
10. Jacobs MA, Spilken AZ, Norman MM, Wohlberg GW, Knapp PH. Interaction of personality and treatment conditions associated with success in smoking control program. *Psychosom Med*. 1971;33:545-556. [SFX](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)
11. Edwards NB, Murphy JK, Downs AD, Ackerman BJ, Rosenthal TL. Doxepin as an adjunct to smoking

cessation: a double-blind pilot study. *Am J Psychiatry*. 1989;146:373-376. [SFX](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)

12. Ferry LH, Robbins AS, Scariati PD, Masterson A, Abbey DE, Burchette RJ. Enhancement of smoking cessation using the antidepressant bupropion hydrochloride [abstract]. *Circulation*. 1992;86(suppl):167. [SFX](#) | [The Knowledge Network](#) | [\[Context Link\]](#)

13. Ferry LH, Burchette RJ. Evaluation of bupropion versus placebo for treatment of nicotine dependence. Paper presented at: American Psychiatric Association Scientific Session; May 26, 1994; Philadelphia, Pa. Presentation NR 554. [\[Context Link\]](#)

14. Niaura R, Goldstein MG, Depue J, Keuthen N, Kristeller J, Abrams D. Fluoxetine, symptoms of depression, and smoking cessation. *Ann Behav Med*. 1995;(suppl):S061. [\[Context Link\]](#)

15. Hurt RD, Sachs DPL, Glover ED, Offord KP, Johnston JA, Dale LC, Khayrallah MA, Schroeder DR, Glover PN, Sullivan CK, Croghan IT, Sullivan PMI. A comparison of sustained-release bupropion and placebo for smoking cessation. *N Engl J Med*. 1997;337:1195-1202. [Ovid Full Text](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)

16. Schatzberg AF, Nemeroff CB. *American Psychiatric Press Textbook of Pharmacology*. Washington, DC: American Psychiatric Press; 1995:147. [\[Context Link\]](#)

17. Zelman DC, Brandon TH, Jorenby DE, Baker TB. Measures of affect and nicotine dependence predict differential response to smoking cessation treatments. *J Consult Clin Psychol*. 1992;60:943-952. [\[Context Link\]](#)

18. Hall SM, Munoz R, Reus VI, Sees K, Duncan CD, Humfleet G, Hartz DT. Mood management and nicotine gum in smoking treatment: a therapeutic contact and placebo-controlled study. *J Consult Clin Psychol*. 1996;64:1003-1009. [\[Context Link\]](#)

19. Robins L, Helzer J, Croughan J, Ratcliff K. National Institute of Mental Health Diagnostic Interview Schedule: its history, characteristics, and validity. *Arch Gen Psychiatry*. 1981;38:381-389. [SFX](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)

20. McNair D, Lorr M, Droppleman L. *Manual for the Profile of Mood States*. San Diego, Calif: Edits; 1971. [\[Context Link\]](#)

21. Shiffman SM, Jarvik ME. Smoking withdrawal symptoms in two weeks of abstinence. *Psychopharmacology*. 1976;50:35-39. [SFX](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)

22. Beck AT, Ward CH, Mendelsohn M, Mock J, Erbaugh J. An inventory for measuring depression. *Arch Gen Psychiatry*. 1961;4:561-571. [SFX](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)

23. Blouin AG, Perez EL, Blouin JH. Computerized administration of the Diagnostic Interview Schedule. *Psychiatry Res*. 1988;23:335-344. [SFX](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)

24. Spielberger DC, Gorsuch RL, Lustene RE. *State-Trait Anxiety Inventory*. Palo Alto, Calif: Consulting Psychologists Press; 1970. [\[Context Link\]](#)

25. Fagerstrom K. Measuring degree of physical dependence to tobacco smoking in reference to

individualization of treatment. *Addict Behav.* 1978;3:235-241. [\[Context Link\]](#)

26. Fagerstrom KO. Measuring nicotine dependence: a review of the Fagerstrom Tolerance Questionnaire. *J Behav Med.* 1989;12:159-182. [SFX](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)

27. McCullagh P, Nelder JA. *Generalized Linear Models*. 2nd ed. New York, NY: Whapman & Hall; 1989. [\[Context Link\]](#)

28. Gibbons R, Hedeker D, Waternaux C. Random regression models: a comprehensive approach to the analysis of longitudinal psychiatric data. *Psychopharmacology.* 1988;24:438-443. [\[Context Link\]](#)

29. Gibbons R, Hedeker D, Elkin I, Waternaux C, Kraemer KH, Greehouse J, Shea T, Imber S, Sotsky S, Watkins J. Some conceptual and statistical issues in the analysis of longitudinal psychiatric data. *Arch Gen Psychiatry.* 1993;50:739-750. [Ovid Full Text](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)

30. Littell RC, Milliken GA, Stroup WW, Wolfinger RD. *SAS System for Mixed Models*. Cary, NC: SAS Institute Inc; 1996. [\[Context Link\]](#)

31. Linnoila M, George L, Guthrie S, Leventhal B. Effects of alcohol consumption and cigarette smoking in antidepressant levels of depressed patients. *Am J Psychiatry.* 1981;138:841-842. [SFX](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)

32. Glassman AH, Covey LS, Dalack GW, Stetner F, Rivelli S, Fleiss J, Cooper TB. Smoking cessation, clonidine, and vulnerability to nicotine among dependent smokers. *Clin Pharmacol Ther.* 1993;54:670-679. [\[Context Link\]](#)


33. Breslau N, Kilbey MM, Andreski P. Nicotine withdrawal symptoms and psychiatric disorders: findings from an epidemiologic study of young adults. *Am J Psychiatry.* 1992;149:464-469. [SFX](#) | [The Knowledge Network](#) | [Bibliographic Links](#) | [\[Context Link\]](#)

Behavior Therapy; Cognitive Therapy; Depressive Disorder; Nortriptyline; Smoking Cessation


## IMAGE GALLERY

Select All

 Export Selected to PowerPoint



☐ Table 1



☐ Figure 1



	MSD	MSD
	Money-History-Passive	History-Passive
Variables	Participants N = 134	Participants N = 80
<i>Demographics</i>		
Age <sup>a</sup>	40.3 (9.3)	41.1 (9.8)
Sex (female)	65 (49%)	49 (61%)
Years of education	13.6 (1.9)	13.6 (1.6)
Married or with a partner (yes)	56 (42%)	35 (44%)
<i>Personality</i>		
Use of cognitive workload strategy <sup>b</sup>	73.6 (15.4)	71.6 (15.4)
Aggression (between 1 and 5) <sup>c</sup>	3.5 (2.1)	3.3 (2.3)
State anxiety (before)	22.6 (9.2)	22.2 (11.4)
State anxiety (after test, after 10 min of working)	34.2 (10.4)	34.6 (12.2)
Flow state (before)	31.1 (10.7)	33.1 (12.1)
<i>Workload</i>		
Back discomfort (working period)	3.9 (2.8)	3.9 (2.8)
Rating of hand fatigue (after 10 min of working)	21.4 (12.3)	15.9 (10.8)
Handgrip strength (before test)	34.0 (7.3)	36.1 (7.7)
Handgrip force (after 10 min)	28.7 (15.4)	29.2 (15.4)
Distraction (on a 100-point scale)	46.4 (16.2)	46.4 (15.4)

<sup>a</sup> Data are given as the mean (SD).

<sup>b</sup> Data are given as the number percentage of subjects.

<sup>c</sup> Data were an aggression difference score for MSD history-passive personality score and score in an aggression test (a 5-point MSQ aggression scale) (see Table 1).

Table 2




Figure 2

Characteristics	Years			
	12	24	36	48
ACVD incidence (positive score)				
death				
Health-related interpretation				
Increasing severity	36 (3%)	30 (2%)	33 (3%)	27 (3%)
$p < 1\%$	24	27	24	26
Increasing arterial disease	74 (5%)	40 (3%)	48 (4%)	44 (3%)
$p < 0.01$	27	28	28	17
Engelbrekt's placement				
Intermediate group				
Increasing stroke	30 (2%)	24 (2%)	20 (2%)	24 (3%)
$p < 0.01$	24	24	22	24
Increasing total group	66 (4%)	67 (5%)	39 (3%)	41 (3%)
$p < 0.01$	29	28	22	23
ACVD incidence (positive score)				
death				
Health-related interpretation				
Increasing severity	21 (2%)	15 (1%)	20 (2%)	15 (1%)
$p < 1\%$	14	13	15	13
Increasing arterial disease	47 (3%)	28 (2%)	46 (4%)	39 (3%)
$p < 1\%$	15	15	17	15
Engelbrekt's placement				
Intermediate group				
Increasing stroke	54 (4%)	21 (2%)	27 (2%)	26 (3%)
$p < 1\%$	14	14	12	14
Increasing total group	62 (4%)	50 (3%)	30 (3%)	29 (3%)
$p < 1\%$	13	13	13	14

Table 3




Figure 3

Major Depressive Disorder History	Baseline Participants	Training Participants
<i>N</i> = 24		
Week 10	10 (41.7%)	10 (41.7%)
Week 12	9 (37.5%)	10 (41.7%)
Week 14	10 (41.7%)	10 (41.7%)
Week 16	9 (37.5%)	10 (41.7%)
Week 18	10 (41.7%)	10 (41.7%)
Week 20	10 (41.7%)	10 (41.7%)
Week 22	10 (41.7%)	10 (41.7%)
Week 24	10 (41.7%)	10 (41.7%)
Week 26	10 (41.7%)	10 (41.7%)
Week 28	10 (41.7%)	10 (41.7%)
Week 30	10 (41.7%)	10 (41.7%)
Week 32	10 (41.7%)	10 (41.7%)
Week 34	10 (41.7%)	10 (41.7%)
Week 36	10 (41.7%)	10 (41.7%)
Week 38	10 (41.7%)	10 (41.7%)
Week 40	10 (41.7%)	10 (41.7%)
Week 42	10 (41.7%)	10 (41.7%)
Week 44	10 (41.7%)	10 (41.7%)
Week 46	10 (41.7%)	10 (41.7%)
Week 48	10 (41.7%)	10 (41.7%)
Week 50	10 (41.7%)	10 (41.7%)
Week 52	10 (41.7%)	10 (41.7%)
Week 54	10 (41.7%)	10 (41.7%)
Week 56	10 (41.7%)	10 (41.7%)
Week 58	10 (41.7%)	10 (41.7%)
Week 60	10 (41.7%)	10 (41.7%)
Week 62	10 (41.7%)	10 (41.7%)
Week 64	10 (41.7%)	10 (41.7%)
Week 66	10 (41.7%)	10 (41.7%)
Week 68	10 (41.7%)	10 (41.7%)
Week 70	10 (41.7%)	10 (41.7%)
Week 72	10 (41.7%)	10 (41.7%)
Week 74	10 (41.7%)	10 (41.7%)
Week 76	10 (41.7%)	10 (41.7%)
Week 78	10 (41.7%)	10 (41.7%)
Week 80	10 (41.7%)	10 (41.7%)
Week 82	10 (41.7%)	10 (41.7%)
Week 84	10 (41.7%)	10 (41.7%)
Week 86	10 (41.7%)	10 (41.7%)
Week 88	10 (41.7%)	10 (41.7%)
Week 90	10 (41.7%)	10 (41.7%)
Week 92	10 (41.7%)	10 (41.7%)
Week 94	10 (41.7%)	10 (41.7%)
Week 96	10 (41.7%)	10 (41.7%)
Week 98	10 (41.7%)	10 (41.7%)
Week 100	10 (41.7%)	10 (41.7%)

Table 5



© 2015 [Ovid Technologies, Inc.](#) All rights reserved.

[About Us](#) | [Contact Us](#) | [Terms of Use](#)

OvidSP\_UI03.18.00.124, SourceID 69505