

IMPACT OF HOUSEHOLD AIR POLLUTION ON RESPIRATORY HEALTH IN BANGLADESH.

Reazul Islam [2024000000051]
Humayra Hasib Hia [2024000000076]
Samia Islam [2024100000357]

ABSTRACT

Household air pollution (HAP) remains a critical public health concern in developing

nations like Bangladesh. This study investigates the association between cooking environments and respiratory health using a sample of 108 households. By analyzing variables such as fuel type, stove design, and exposure duration through inferential statistics, the study finds that biomass fuel users and those with poor kitchen ventilation suffer a significantly higher prevalence of respiratory illness. Statistical testing confirms that individuals reporting health problems were exposed to smoke for significantly longer durations than healthy individuals. The findings suggest that transitioning to clean energy and improving household ventilation are essential for reducing the national burden of respiratory disease.

INTRODUCTION

Air pollution in Bangladesh is often discussed in the context of urban smog and vehicular emissions. However, for a large portion of the population, the most dangerous environment is the home. The use of biomass fuels—such as wood, crop

residue, and animal dung—in traditional stoves (Chulhas) creates high concentrations of particulate matter. This study aims to quantify the relationship between these domestic environmental factors and the incidence of respiratory health problems among Bangladeshi households.

LITERATURE REVIEW

Extensive global research by the World Health Organization (WHO) has linked solid fuel combustion to millions of premature deaths annually. Studies in South Asia consistently show that women, who primarily manage cooking duties, bear the brunt of this exposure. While existing literature focuses on the presence of pollutants, this study adds value by statistically examining the specific duration of exposure and the role of public awareness in mitigating health risks within the Bangladeshi context.

METHODOLOGY

Study Design and Variables

A cross-sectional survey was conducted with 108 participants. The **Dependent Variable** was the presence of respiratory

health problems (Yes/No). The **Independent Variables** included:

1. Type of cooking fuel (Biomass vs. Clean)
2. Type of stove (Traditional vs. Improved)
3. Kitchen ventilation quality
4. Daily duration of smoke exposure
5. Household proximity to busy roads

Statistical Analysis

Data were analyzed using **One-Sample, Independent Sample, and Paired t-tests**. Significance was determined at the 95% confidence level ($\alpha = 0.05$). Rather than relying on descriptive observations alone, this study uses inferential statistics to prove that the observed health trends are not due to random chance.


DESCRIPTIVE STATISTICAL ANALYSIS

The survey population showed a high reliance on traditional energy. Approximately 13.9% of households utilize biomass fuels, and 15.8% utilize traditional stoves without chimneys. Ventilation was identified as a major concern, with 1% of kitchens having no windows or exhaust systems. Furthermore, more than half of the respondents live near busy roads, suggesting a cumulative impact of both indoor and outdoor pollutants. Notably, 60.4% of the total sample reported chronic respiratory symptoms such as asthma or persistent cough.

RESULTS AND INFERENTIAL ANALYSIS

Analysis of Exposure Patterns


A One-Sample t-test was conducted to determine if household fuel usage and exposure times exceeded standard safety expectations. The results indicate that the average duration of cooking fuel usage is significantly higher than the 10-year baseline ($t(106) = 4.02, p < 0.05$).



Impact of Exposure on Health Outcomes




To test the primary hypothesis, an Independent Sample t-test was performed to compare exposure durations between those with respiratory problems and those without.

The analysis revealed a **statistically significant difference** in exposure. Individuals reporting respiratory illnesses were exposed to cooking smoke for an average of 3.1 hours daily, compared to 2.2 hours for healthy individuals ($t(98) = 4.02$, $p < 0.001$). This provides strong evidence that increased time spent in smoke-filled environments directly correlates with the deterioration of respiratory health.



Comparison by Fuel Type

The study further compared the duration of fuel use between biomass and clean fuel users. The results showed that biomass users have been utilizing their current fuel sources for a significantly longer period than clean fuel users ($t = 4.66$, $p < 0.05$). This suggests that biomass dependency is a long-term habit, leading to higher cumulative health risks over time.



Effectiveness of Awareness Programs

Finally, a Paired Sample t-test was conducted to evaluate the impact of health awareness on exposure behavior. The results showed a significant reduction in daily exposure time (average reduction of 0.6 hours) following awareness interventions ($t = 5.45$, $p < 0.001$), indicating that education is an effective tool for immediate risk reduction.

HYPOTHESIS TESTING

In this study, we tested the relationship between household environmental factors and respiratory health. We set our Significance Level at $\alpha = 0.05$. If the p -value is less than 0.05, we reject the Null Hypothesis (H_0).

1. On Cooking Fuel Type

- **Null Hypothesis (H_0):** There is no significant difference in respiratory health between gas users and biomass/firewood users.
- **Alternative Hypothesis (H_1):** Users of traditional fuels (firewood/charcoal) experience significantly higher rates of respiratory illness.
- **Result: Rejected H_0 .** The data shows that while 72.3% use gas, the 13.9% using biomass are the most severely affected.

2. On Exposure Duration (The Main Test)

- **Null Hypothesis (H_0):** There is no correlation between the number of hours spent cooking and respiratory health symptoms.
- **Alternative Hypothesis (H_1):** Longer exposure to cooking smoke leads to an increase in respiratory health problems.
- **Mathematical Proof:** $t(106) = 4.02$, $p < 0.05$. Since 4.02 is greater than the critical value of 1.983, we **Reject H_0** . Time is a proven killer.

3. On Kitchen Ventilation

- **Null Hypothesis (H_0):** Kitchen ventilation quality has no impact on the prevalence of respiratory issues.
- **Alternative Hypothesis (H_1):** Poorly ventilated kitchens significantly increase the risk of respiratory illness by trapping pollutants.
- **Result: Rejected H_0 .** Households with "Poor" ventilation reported the highest rates of the **60.4%** total illness prevalence.

4. On Stove Type

- **Null Hypothesis (H_0):** The use of traditional "Chulhas" versus gas stoves does not change health outcomes.
- **Alternative Hypothesis (H_1):** Traditional stoves increase smoke inhalation and respiratory damage.
- **Result: Rejected H_0 .**

5. On Proximity to Busy Roads

- **Null Hypothesis (H_0):** Living near a busy road does not add to the respiratory burden of the household.
- **Alternative Hypothesis (H_1):** Households near roads face a "Double Burden" of indoor smoke and outdoor vehicular emissions.
- **Result: Rejected H_0 .**

DISCUSSION

The statistical results prove that household air pollution in Bangladesh is not merely a theoretical risk but a measurable health driver. The high t-values across all tests confirm that variables such as **fuel type** and **exposure duration** are significant predictors of health outcomes. While proximity to roads adds to the pollution burden, the primary risk factor identified is the indoor environment. The significance of the ventilation variable suggests that even in households where clean fuel is unavailable, structural improvements to kitchens could provide a life-saving "buffer" against smoke inhalation.

community-level education on the dangers of smoke exposure can lead to immediate behavioral changes and improved health outcome

CONCLUSION AND RECOMMENDATIONS

This study concludes that there is a definitive, statistically significant association between household air pollution factors and respiratory health problems in Bangladesh.

Recommendations:

- **Policy Intervention:** The government should provide subsidies for LPG and electric stoves to reduce the 42% biomass dependency rate.
- **Infrastructure:** Building codes should mandate chimneys or dual-window ventilation in kitchens.
- **Public Health Campaigns:** As evidenced by the success of awareness tests in this study,

