

Problem Set 3

Michael Fryer

2025-11-20

Collaborators: ...

1 Entropy & Divergence

Suppose biased coin lands heads 60% of the time.

a) What is the entropy of this coin?

```
# 60% heads, 40% tails
p <- c(0.6, 0.4)
cat("Entropy: ", -sum(p*log(p)))
```

```
## Entropy: 0.6730117
```

b) Suppose another biased coin lands tails 30% of the time. Which coin has higher entropy? Justify and explain your answer.

```
# 70% heads, 30% tails
q <- c(0.7, 0.3)
cat("Entropy: ", -sum(q*log(q)))
```

```
## Entropy: 0.6108643
```

The coin in part a) has a higher entropy than the coin in b). The variance in the probabilities in b) is larger than in a) i.e. the more unique events there are with similar probabilities of occurring, the larger the entropy will be.

c) Suppose the first coin is the true distribution and the second coin's distribution is your approximation of the first coin. What is the KL divergence between these two distributions? Explain and justify your answer.

```
cat("KL divergence: ", sum(p*log(p/q)))
```

```
## KL divergence: 0.02258242
```

We observe a KL divergence of 0.022. We expect this value to be small as p and q are relatively close together and the KL divergence is the average difference in log probabilities between the two models. This value would be much larger if instead q = 0.9, 0.1.

2 Collider Bias and Information Criteria

Return to the textbook example in §6.3.1, which explores the relationship between age, marriage and happiness. This example makes use of the simulation model `sim_happiness()`, which is part of the `rethinking` package:

```
d <- sim_happiness( seed=1515 , N_years=1000)
d2 <- d[ d$age>17 , ] # only adults
```

```
d2$A <- (d2$age - 18) / (65 - 18)
d2$mid <- d2$married + 1
```

Compare the two models, `m6.9` and `m6.10`, using both PSIS and WAIC scores using `compare()`.

a) Which model is expected to make better predictions according to these two information criteria? Explain your answer.

b) On the basis of the causal model, how should you interpret the parameter estimates from the model preferred by PSIS and WAIC?

3 Tulips and Interactions

Tulips need both water and light to bloom. But do water and light work independently, or do they interact? The `tulips` dataset in the `rethinking` package contains experimental data on tulip blooms under different conditions.

```
data(tulips)
d <- tulips
str(d)
```

```
## 'data.frame': 27 obs. of 4 variables:
## $ bed : Factor w/ 3 levels "a","b","c": 1 1 1 1 1 1 1 1 1 2 ...
## $ water : int 1 1 1 2 2 2 3 3 3 1 ...
## $ shade : int 1 2 3 1 2 3 1 2 3 1 ...
## $ blooms: num 0 0 111 183.5 59.2 ...
```

The dataset contains:

- **bed**: Bed identifier (grouping variable)
- **water**: Amount of water (1 = low, 2 = medium, 3 = high)
- **shade**: Amount of shade (1 = low, 2 = medium, 3 = high)
- **blooms**: Number of blooms produced

Before fitting any models, standardize the predictor variables and the outcome:

a) Fit a model (`m3.1`) that predicts blooms using water and shade as independent predictors (no interaction). That is, assume the effect of water on blooms is the same regardless of shade level, and vice versa.

Use prior predictive simulation to justify your priors. Write 2-3 sentences explaining your choice of priors.

Hint: With standardized variables, you should use relatively weak priors. A slope of 2.0 would be implausibly large for most ecological relationships.

b) Now fit a model (`m3.2`) that includes an interaction between water and shade. This allows the effect of water on blooms to depend on the level of shade (and vice versa).

Hint: In the model formula, you can specify the interaction as $W * S$, which is shorthand for $W + S + W * S$. Or you can write out all three terms explicitly.

Plot the posterior predictions for both models (`m3.1` and `m3.2`) side by side. For each model, create a plot showing how predicted blooms change with water at different levels of shade (e.g., plot three lines for shade = -1, 0, 1).

c) Use both PSIS and WAIC to compare your two models. Which model is expected to make better predictions? Is the difference substantial?

Create the comparison plot using `plot(compare(m3.1, m3.2))` and interpret what you see. Write 3-4 sentences explaining:

- Which model performs better
- How large the difference is relative to the standard error
- What this tells you about whether water and shade interact in affecting tulip blooms

d) Examine the pointwise PSIS diagnostics for your preferred model. Are there any problematic observations (high Pareto k values)? If so, which observations are influential and why might they be outliers?

Hint: Use `PSIS(model, pointwise=TRUE)` and examine the k values.

e) Based on your analysis, write 2-3 sentences answering: Do water and light work independently to affect tulip blooms, or do they interact? What does this mean biologically?

4 Reflection

Look back at Problems 2 and 3. Both involve interactions, but of different kinds:

- In Problem 2, you examined collider bias, where conditioning on a variable (marriage status) creates a spurious association between age and happiness
- In Problem 3, you examined multiplicative interactions, where the effect of one variable (water) depends on the level of another (shade)

Write 3-4 sentences explaining: What is the difference between these two types of “interactions”? How are they similar and how are they different from a causal modeling perspective?

5 AI Declaration

Please declare your collaborators in the class and how you used AI (if at all) to complete this assignment. If you used AI, include the prompts you used and explain what you learned from its responses that you didn't understand initially.

6 Appendix