

Chapter 1

Introduction to Graph

Graph theory on August 8, 2023

Contents

Graph definitions

- Terminology
- Special Graphs
- Bipartite graph

Representing Graphs and Graph Isomorphism

- Representing Graphs
- Graph Isomorphism

Exercise

- Graph
- Bipartite graph
- Isomorphism

Vo Dang Khoa, Huynh Tuong Nguyen
Faculty of Information Technology
Industrial University of Ho Chi Minh City
{khoavo;htnguyen}@iuh.edu.vn

① Graph definitions

Terminology

Special Graphs

Bipartite graph

② Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

③ Exercise

Graph

Bipartite graph

Isomorphism

Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism

Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism

Course learning outcomes

CLO.1 Understanding of the basic concepts of graphs
Special types of graph,
computer based graph representation, isomorphism,
planar graph, connectivity in graph, graph traversal.

CLO.2 Describe definition of path and circuit
Identify the existence of Euler path & circuit
Identify the existence of Hamilton path & circuit

CLO.3 Compute minimum spanning tree in a (weighted) graph
Use algorithms: Prim, Kruskal

CLO.4 Determine shortest path in a weighted graph
Use algorithms: Dijkstra, Bellman-Ford, Floyd-Warshall

CLO.5 Solve maximum flow problem
Use Ford-Fulkerson's algorithm

The need of the graph

- Representation/Storing
- Searching/sorting
- Optimization

Its applications

- Electric circuit/board
- Chemical structure
- Networking
- Map, geometry, ...


- Graph theory is useful for analysing “things that are connected to other things”.
- Some difficult problems become easy when represented using a graph.

Definition


A graph (đồ thị) G is a pair of (V, E) , which are:

- V – nonempty set of **vertices** (nodes) (đỉnh)
- E – set of **edges** (cạnh)

A graph captures abstract relationships between vertices.



Undirected graph




Directed graph

Definition (Simple graph (đơn đồ thị))

- Each edge connects two different vertices, and
- No two edges connect the same pair of vertices


An edge between two vertices u and v is denoted as $\{u, v\}$



Definition (Multigraph (đa đồ thị))

Graphs that may have multiple edges connecting the same vertices.


An unordered pair of vertices $\{u, v\}$ are called **multiplicity m** (*bội m*) if it has m different edges between.



Definition (Pseudograph (giả đồ thị))

Are multigraphs that have

- **loops** (*khuyên*)– edges that connect a vertex to itself




Definition (Directed Graph (đồ thị có hướng))

A directed graph G is a pair of (V, E) , in which:

- V – nonempty set of vertices
- E – set of directed edges (*cạnh có hướng, arcs*)

A directed edge **start** at u and **end** at v is denoted as (u, v) .



Neighborhood

In an undirected graph $G = (V, E)$,

- two vertices u and $v \in V$ are called **adjacent** (*liền kề*) if they are **end-points** (*điểm đầu mút*) of edge $e \in E$, and
- e is **incident with** (*cạnh liên thuộc*) u and v
- e is said to **connect** (*cạnh nối*) u and v ;

The degree of a vertex

The **degree of a vertex** (*bậc của một đỉnh*), denoted by $\deg(v)$ is the number of edges incident with it, except that a loop contributes twice to the degree of that vertex.

- **isolated** vertex (*đỉnh cô lập*): vertex of degree **0**
- **pendant** vertex (*đỉnh treo*): vertex of degree **1**

Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph



Bipartite graph

Isomorphism

Example

Example

What are the degrees and neighborhoods of the vertices in these graphs?



Solution

In G , $\deg(a) = 2$, $\deg(b) = \deg(c) = \deg(f) = 4$, $\deg(d) = 1, \dots$

Neighborhoods of these vertices are

$$N(a) = \{b, f\}, N(b) = \{a, c, e, f\}, \dots$$

In H , $\deg(a) = 4$, $\deg(b) = \deg(e) = 6$, $\deg(c) = 1, \dots$

Neighborhoods of these vertices are

$$N(a) = \{b, d, e\}, N(b) = \{a, b, c, d, e\}, \dots$$

Theorem (The Handshaking Theorem)

Let $G = (V, E)$ be an undirected graph with m edges. Then

$$2m = \sum_{v \in V} \deg(v)$$

(Note that this applies even if multiple edges and loops are present.)

Theorem

An undirected graph has an even number of odd-degree vertices.

Prove that ...

...

If the number of vertices in an undirected graph is an odd number, then there exists an even-degree vertex.

...

If the number of vertices in an undirected graph is an odd number, then the number of vertices with even degree is odd.

...

If the number of vertices in an undirected graph is an even number, then the number of vertices with even degree is even.

Neighborhood

In an directed graph $G = (V, E)$,

- u is said to be **adjacent to** (*nối tới*) v and v is said to be **adjacent from** (*được nối từ*) u if (u, v) is an arc of G , and
- u is called **initial vertex** (*đỉnh đầu*) of (u, v)
- v is called **terminal** (*đỉnh cuối*) or **end vertex** of (u, v)
- the initial vertex and terminal vertex of a loop are the same.

The degree of a vertex

In a graph G with directed edges:

- **in-degree** (*bậc vào*) of a vertex v , denoted by $\deg^-(v)$, is the number of arcs with v as their terminal vertex.
- **out-degree** (*bậc ra*) of a vertex v , denoted by $\deg^+(v)$, is the number of arcs with v as their initial vertex.

Note: a loop at a vertex contributes **1** to both the in-degree and the out-degree of this vertex.

Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism



Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism


Theorem

Let $G = (V, E)$ be a graph with directed edges. Then


$$\sum_{v \in V} \deg^-(v) = \sum_{v \in V} \deg^+(v) = |E|.$$

Complete Graphs

A complete graph (*đồ thị đầy đủ*) on n vertices, K_n , is a simple graph that contains **exactly one edge** between each pair of distinct vertices.





K_5



K_4

A cycle (đồ thị vòng) C_n , $n \geq 3$, consists of n vertices v_1, v_2, \dots, v_n and edges $\{v_1, v_2\}, \{v_2, v_3\}, \dots, \{v_{n-1}, v_n\}$, and $\{v_n, v_1\}$.

 C_5  C_4

Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism



Exercise

Graph

Bipartite graph

Isomorphism

We obtain a wheel (*đồ thị hình bánh xe*) W_n when we add an additional vertex to a cycle C_n , for $n \geq 3$, and connect this new vertex to each of the n vertices in C_n .

 W_5  W_4

Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism



Exercise

Graph

Bipartite graph

Isomorphism

An n -dimensional hypercube (*khối n chiều*), Q_n , is a graph that has vertices representing the 2^n bit strings of length n . Two vertices are adjacent iff the bit strings that they represent differ in exactly one bit position.

 Q_1  Q_2  Q_3

What's about Q_4 ?

Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism

Applications of Special Graphs

- Local networks topologies
 - Star, ring, hybrid
- Parallel processing
 - Linear array
 - Mesh network

Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism

Exercise (5)

Give the number of edges in function of number of vertices in a complete graph K_n .

Exercise (6)

Give an undirected simple graph $G = (V, E)$ with $|V| = n$, show that

- a $\forall v \in V, \deg(v) < n$,
- b there does not exist simultaneously both a vertex of degree 0 and a vertex of degree $(n - 1)$ with $n \geq 2$,
- c deduce that there are at least two vertices of the same degree.

Exercise (7)

Is it possible that each person has exactly 3 friends in the same group of 9 people ?

Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism




Bipartite Graphs

Definition

A simple graph G is called bipartite (*đồ thị phân đôi*) if its vertex set V can be partitioned into two disjoint sets V_1 and V_2 such that every edge in the graph connects a vertex in V_1 and a vertex in V_2 (so that no edge in G connects either two vertices in V_1 or two vertices in V_2)

Example


C_6 and C_3 are bipartite ?



Definition

A complete bipartite $K_{m,n}$ is a graph that

- has its vertex set partitioned into **two subsets** of m and n vertices, respectively,
- with an edge between two vertices iff one vertex is in the first subset and the other is in the second one



$$K_{3,3}$$

Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise


Graph

Bipartite graph

Isomorphism

Example (Bipartite graphs?)

- C_6
- C_n
- K_3
- K_n
- Q_n
- the following graph



Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism






Exercise

Graph

Bipartite graph

Isomorphism

Bipartite graph



Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism




New Graph From Old

Definition

A **subgraph** (đồ thị con) of a graph $G = (V, E)$ is a graph $H = (W, F)$ where $W \subseteq V$ and $F \subseteq E$.

Definition

The **union** (hợp) of two simple graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ is a simple graph with vertex set $V_1 \cup V_2$ and edge set $E_1 \cup E_2$. The union of G_1 and G_2 is denoted by $G_1 \cup G_2$.

 G_1  G_2  $G_1 \cup G_2$

Contents

Graph definitions

[Terminology](#)[Special Graphs](#)[Bipartite graph](#)


Representing Graphs and Graph Isomorphism

[Representing Graphs](#)[Graph Isomorphism](#)

Exercise

[Graph](#)[Bipartite graph](#)[Isomorphism](#)

Planar Graphs



Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise



Graph

Bipartite graph

Isomorphism

Definition

- A graph is called **planar** (*phẳng*) if it can be drawn in the plane **without any edges crossing**.
- Such a drawing is called **planar representation** (*biểu diễn phẳng*) of the graph.

 K_4  K_4 with no crossing

Contents

Graph definitions

[Terminology](#)[Special Graphs](#)[Bipartite graph](#)

Representing Graphs and Graph Isomorphism


[Representing Graphs](#)[Graph Isomorphism](#)

Exercise


[Graph](#)[Bipartite graph](#)[Isomorphism](#)

Corollary

- If G is a **connected planar simple graph** with e edges and v vertices where $v \geq 3$, then $e \leq 3v - 6$.
- If G is a **connected planar simple graph** with e edges and v vertices where $v \geq 3$, and no circuits of length 3, then $e \leq 2v - 4$.



$K_{3,3}$
Non-planar



K_5
Non-planar

Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise


Graph

Bipartite graph

Isomorphism

Definition

- Given a planar graph G , an **elementary subdivision** (*phân chia sơ cấp*) is removing an edge $\{u, v\}$ and adding a new vertex w together with edges $\{u, w\}$ and $\{w, v\}$.
- Graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ are called **homeomorphic** (*đồng phôi*) if they can be obtained from the same graph by a sequence of elementary subdivisions.



Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism

Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise




Graph

Bipartite graph

Isomorphism


Theorem

A graph is nonplanar iff it contains a *subgraph homeomorphic to $K_{3,3}$ or K_5* .




Exercise

- Is K_4 planar?
- Is Q_3 planar?



K_4



Q_3

Adjacency Lists (Danh sách kề)

Contents

Graph definitions

- Terminology
- Special Graphs
- Bipartite graph


Representing Graphs and Graph Isomorphism

- Representing Graphs
- Graph Isomorphism


Exercise

- Graph
- Bipartite graph
- Isomorphism

Vertex	Adjacent vertices
a	b, c, e
b	a
c	a, d, e
d	c, e
e	a, c, d




Initial vertex	Terminal vertices
a	b, c, d, e
b	b, d
c	a, c, e
d	c, e
e	b, c, d



Example

Determine adjacency list of vertices A & B from the following graph (Write the elements in the alphabetical order)



Contents

Graph definitions

- Terminology
- Special Graphs
- Bipartite graph

Representing Graphs and Graph Isomorphism


- Representing Graphs
- Graph Isomorphism

Exercise

- Graph
- Bipartite graph
- Isomorphism

Example

Determine adjacency list of vertices D & F from the following graph (Write the elements in the alphabetical order)



Contents

Graph definitions

- Terminology
- Special Graphs
- Bipartite graph

Representing Graphs and Graph Isomorphism

- Representing Graphs
- Graph Isomorphism


Exercise

- Graph
- Bipartite graph
- Isomorphism

Definition

Adjacency matrix (*Ma trận kề*) A_G of $G = (V, E)$

- Dimension $|V| \times |V|$
- Matrix elements
$$a_{ij} = \begin{cases} 1 & \text{if } (v_i, v_j) \in E \\ 0 & \text{otherwise} \end{cases}$$



$$\begin{matrix} a \\ b \\ c \\ d \end{matrix} \begin{bmatrix} & a & b & c & d \\ a & 0 & 1 & 1 & 1 \\ b & 1 & 0 & 1 & 0 \\ c & 1 & 1 & 0 & 0 \\ d & 1 & 0 & 0 & 0 \end{bmatrix}$$

Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise


Graph

Bipartite graph

Isomorphism


Example

Give adjacency matrix according to the following graph



Example

Give adjacency matrix according to the following graph



Example

Give the graph defined by the following adjacency matrix

$$\begin{array}{c} A \\ B \\ C \\ D \\ E \end{array} \left[\begin{array}{ccccc} A & B & C & D & E \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{array} \right]$$

Contents

Graph definitions

- Terminology
- Special Graphs
- Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

- Graph Isomorphism

Exercise

- Graph
- Bipartite graph
- Isomorphism

Example

Give the directed graph defined by the following adjacency matrix

$$\begin{array}{c} A \\ B \\ C \\ D \\ E \end{array} \left[\begin{array}{ccccc} A & B & C & D & E \\ 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 \end{array} \right]$$

Contents

Graph definitions

- Terminology
- Special Graphs
- Bipartite graph

Representing Graphs and Graph Isomorphism

- Representing Graphs
- Graph Isomorphism


Exercise

- Graph
- Bipartite graph
- Isomorphism

Definition

Incidence matrix (*ma trận liên thuộc*) M_G of $G = (V, E)$

- Dimension $|V| \times |E|$
- Matrix elements
$$m_{ij} = \begin{cases} 1 & \text{if } e_j \text{ is incident with } v_i \\ 0 & \text{otherwise} \end{cases}$$



$$\begin{matrix} a \\ b \\ c \\ d \end{matrix} \begin{bmatrix} e_1 & e_2 & e_3 & e_4 \\ 1 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

Contents

Graph definitions

[Terminology](#)[Special Graphs](#)[Bipartite graph](#)

Representing Graphs and Graph Isomorphism


[Representing Graphs](#)[Graph Isomorphism](#)

Exercise

[Graph](#)[Bipartite graph](#)[Isomorphism](#)


Example

Give incidence matrix according to the following graph



Example

Give incidence matrix according to the following graph



Contents

Graph definitions

- Terminology
- Special Graphs
- Bipartite graph

Representing Graphs and Graph Isomorphism


- Representing Graphs
- Graph Isomorphism

Exercise

- Graph
- Bipartite graph
- Isomorphism

Example

Give incidence matrix according to the following graph



Contents

Graph definitions

- Terminology
- Special Graphs
- Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

- Graph Isomorphism


Exercise

- Graph
- Bipartite graph
- Isomorphism

Definition

$G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ are **isomorphic** (đẳng cấu) if there is a **one-to-one function** f from V_1 to V_2 with the property that a and b are adjacent in G_1 iff $f(a)$ and $f(b)$ are adjacent in G_2 , for all a and b in V_1 . Such a function f is called an **isomorphism** (một đẳng cấu).

(i.e. there is a one-to-one correspondence between vertices of the two graphs that preserves the adjacency relationship.)



Isomorphism function $f : U \rightarrow$

V with

$$f(u_1) = v_1 \quad f(u_2) = v_4$$

$$f(u_3) = v_3 \quad f(u_4) = v_2$$

Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism






Exercise

Graph

Bipartite graph

Isomorphism

Bipartite graph



Contents

Graph definitions

Terminology
Special Graphs
Bipartite graph




Representing Graphs and Graph Isomorphism

Representing Graphs
Graph Isomorphism

Exercise

Graph
Bipartite graph
Isomorphism

Isomorphism ?



Contents

Graph definitions

- Terminology
- Special Graphs
- Bipartite graph

Representing Graphs and Graph Isomorphism

- Representing Graphs
- Graph Isomorphism

Exercise

- Graph
- Bipartite graph
- Isomorphism

Isomorphism?

Contents

Graph definitions


- Terminology
- Special Graphs
- Bipartite graph

Representing Graphs and Graph Isomorphism



- Representing Graphs
- Graph Isomorphism

Exercise

- Graph
- Bipartite graph
- Isomorphism



Isomorphism?



Contents

Graph definitions

- Terminology
- Special Graphs
- Bipartite graph

Representing Graphs and Graph Isomorphism

- Representing Graphs
- Graph Isomorphism

Exercise

- Graph
- Bipartite graph
- Isomorphism

Are the simple graphs with the following adjacency matrices isomorphic ?

$$① \quad \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}$$

$$② \quad \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix}$$

$$③ \quad \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix}$$

Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism

Contents

Graph definitions

Terminology

Special Graphs

Bipartite graph

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism

Determine whether the graphs (without loops) with the incidence matrices are isomorphic.

- $\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix}$
- $\begin{pmatrix} 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 \end{pmatrix}$
- Extend the definition of isomorphism of simple graphs to undirected graphs containing loops and multiple edges.
- Define isomorphism of directed graphs