

Experiments with microwaves

Nele Wurster and Hannes Winkler

Abstract

Executed: 10th June 2026

Submitted: 24rd June 2026

1 Introduction

The spectrum of electromagnetic radiation covers a wide range of wavelengths, ranging from high-energy γ -rays to low-energy radio waves, all of which play an important role in today's world. Examples include X-rays used in medical diagnostics, the radio waves employed by mobile phones to transmit information, and microwaves used to heat your lunch.

In this experiment, we take a closer look at the latter type of radiation: microwaves. Their wavelengths range from one meter to one millimeter, giving them a wide variety of applications. Furthermore, microwaves play a significant role in our understanding of the universe through the study of the cosmic microwave background, which is believed to be a remnant of the Big Bang, the event that marked the beginning of the universe. Microwaves exhibit many interesting properties. In the following, several of these properties will be examined and discussed, namely the angular distribution and polarization properties of microwaves, their wavelengths, and the formation of evanescent waves. Using these phenomena, we will also determine the lattice constant of a macroscopic cube.

2 Angular distribution of microwaves

To examine the angular distribution of microwaves, the transmitter was placed on a goniometertable, such that the horn is located approximately at the pivot point. The angular distribution was measured to both sides $\pm 90^\circ$ in equidistant steps. The measurements are done with a photo diode that is connected to an Agilent 34402A. The measured intensity was normalized to 1 for $\theta = 0^\circ$ and is shown in fig. 1. The density of measured points around $\theta = 0^\circ$ is quite low, which is not ideal. One should see the symmetry around the 0-point, but with our




Figure 1: Measured voltage within the relevant angular range normalized to 1 in dependence of the angle θ . The left and right full width at half maximum are marked.

measurements, one can only guess it.

Nevertheless was the angle stretch of the radiation quantified by calculating the halfangle θ_H where the full width at half maximum lies.

$$\theta_H = (28.4 \pm 2.5)^\circ \quad (1)$$

As expected due to the transmitter being a horn antenna, this confirms that the microwave radiation is pointed towards the 0° direction.

3 Focal length of paraffin wax lens

To further investigate microwaves, it is necessary to produce collimated rays. For that a paraffin wax lens was used which focal point has to be determined. This was achieved by placing a wax lens between transmitter and receiver. The distance between the transmitter and the lens is $g = (80 \pm 1) \text{ cm}$. The re-




Figure 2: Intensity profile of the microwave radiation in relation to it's distance from the lens.

ceiver is placed with maximal possible distance to the lense and then progressively moved towards it. The radiation was measured in every step. To determine the peak radiation level, a smooth curve was fitted to the experimental data and it's maximum was determined to $b' = (151.8 \pm 6.0) \text{ cm}$. The plot is shown in fig. 2.

The focal length was then calculated using the thin-lense equation:

$$f = \frac{g \cdot b}{g + b} \quad \text{with} \quad b = b' - g \quad (2)$$

which results in a focal lenght of ($f = 37.84 \pm 0.48 \text{ cm}$). In the following parts of the experiment, the focal lenght was roughly placed at 40 cm, such that the rays would be collimated.

4 Determination of the lattice constant of a cubic crystal

In solid state physics, the θ -rocking-curve method is often used to examine the characteristics of an atomic lattice, namely the lattice constant. In this part of the experiment, this method was tested with microwaves on a macroscopic styrofoam cube doted with aluminum wrapped balls, such that a cubic lattice emerges. The bragg's law states that monochromatic radiation of wavelenght λ interferes constructive under the angle θ if

$$n\lambda = 2d\sin(\theta) \quad (3)$$

is realised. Here n is the number of the maximum or diffraction order, λ the wavelenght measured in a previous part, d the spacing between adjacent crystal planes, and θ is the angle at which the maximum occurs. The setup used in the experiment is shown in




Figure 3: Experimental setup to measure the lattice constant of an macroscopic cube

fig. 3. Starting the measurement at an angle $\theta = 0^\circ$, the crystal was rotated in steps of 2° while the detector was rotated to be at 2θ to ensure the symmetric condition required for Bragg reflection.

While Bragg's law is independent of the orientation of the crystal planes, it acutally plays a role in the measurement. That is why we need to introduce Miller indices h, k, l to describe which plane we are currently measuring. For a cubic crystal, the spacing between lattice plane can be expressed as:

$$d_{hkl} = \frac{d_0}{\sqrt{h^2 + k^2 + l^2}} \quad (4)$$

Using equations 3 and 4 one can calculate the lattice constant d_0 with:

$$d_0 = d_{hkl} \sqrt{h^2 + k^2 + l^2} = \frac{n\lambda}{2\sin(\theta)} \sqrt{h^2 + k^2 + l^2} \quad (5)$$

The sides with the Miller indicies (100) and (110) were measured and the intensity in terms of the rotation angle is shown in fig. 4 and 5 respectively.




Figure 4: Measured Bragg diffraction from the (100) plane for angles $\theta = [0, 60]$. The first maximum was determined and marked.

The first maximum for the (100) plane was measured at an angle $\theta = (25.89 \pm 0.5)^\circ$.




Figure 5: Measured Bragg diffraction from the (110) plane for angles $\theta = [0, 50]$. The first maximum was determined and marked.

The first maximum for the (110) plane was measured at an angle $\theta = (34.95 \pm 0.5)^\circ$.

Using eq. 5 and the wavelength $\lambda = \text{hehrher}$ the lattice constant can be calculated:

$$\begin{aligned} (100) : d_0 &= (3.81 \pm 0.59) \text{ cm} \\ (110) : d_0 &= (4.11 \pm 0.63) \text{ cm} \end{aligned} \quad (6)$$

Hier noch die diskussion, wenn dann in der rechnung die tatsächliche wellenlänge eingetragen ist

References

- [1] https://www.physik.uni-wuerzburg.de/fileadmin/11000000/03_Studium/02_Bachelor/Grundpraktikum/_imported/fileadmin/11016800/Modulbeschreibungen/C-Modul/Versuch_42_ueberarbeitet.pdf last visited on 22.06.2026

- [2] https://www.physik.uni-wuerzburg.de/fileadmin/11000000/03_Studium/02_Bachelor/Grundpraktikum/_imported/fileadmin/11016800/Geraete/Agilent-34405a.pdf last visited on 22.06.2026