

Stability of the Transonic Plasma Flow in the Magnetic Nozzle

Hunt Feng
Supervisor: Andrei Smolyakov

Faculty of Physics and Engineering Physics
University of Saskatchewan



Goals and Significance of the Research

Goals:

- Investigate the stability of transonic plasma flow in magnetic nozzle.
- Understand the plasma behavior near the nozzle throat.
- Dealing with singularity at nozzle throat both analytically and numerically.

Significance:


- Understand better the transonic plasma flow in magnetic nozzle.
- Addresses the complexity introduced by the singularity in the governing equations.
- Offer insights into plasma flow in magnetic mirror configuration.

Linear Instability of Plasma Flow


- The instability of plasma flow refers to the tendency of a plasma system to deviate from a stable, equilibrium state and exhibit perturbations or fluctuations in its behavior. [4]
- To investigate linear instability, we assume oscillating perturbed quantities, $\tilde{n}, \tilde{v} \sim \exp(-i\omega t)$.
 - ① If $\text{Im}(\omega) > 0$, then it is unstable flow since the perturbations grow exponential in time, $\exp(\text{Im}(\omega)t)$.
 - ② If $\text{Im}(\omega) \leq 0$, then it is stable flow since the perturbations decay/unchanged in time.
- In this research we will focus on the linear instability only.

Magnetic Nozzle

- A magnetic nozzle is a device that uses a magnetic field to shape and control the flow of charged particles in a plasma propulsion system.
- Instabilities may affect magnetic nozzle operation and the resulting thrust. [7]



(a) Simplified representation of magnetic nozzle.



(b) A simplified magnetic field of magnetic nozzle.

Figure 1: Simplified representation of magnetic nozzle. Length is normalized.

Governing Equations

The nondimensionalized governing equations for the plasma flow in magnetic nozzle are [14]

$$\text{Cons. of Den.} \quad \frac{\partial n}{\partial t} + n \frac{\partial v}{\partial z} + v \frac{\partial n}{\partial z} - nv \frac{\partial_z B}{B} = 0 \quad (1)$$

$$\text{Cons. of Mom.} \quad n \frac{\partial v}{\partial t} + nv \frac{\partial v}{\partial z} = - \frac{\partial n}{\partial z} \quad (2)$$

where n, v are density and velocity, respectively.

The equilibrium quantities n_0, v_0 must satisfy the condition,

$$\frac{\partial}{\partial z} \left(\frac{n_0 v_0}{B} \right) = 0 \quad (3)$$

$$v_0 \frac{\partial v_0}{\partial z} = - \frac{1}{n_0} \frac{\partial n_0}{\partial z} \quad (4)$$

Equilibrium Velocity Profiles




Figure 2: Velocity if normalized to sound speed. There are 4 different cases for velocity profile: subsonic, supersonic, accelerating and decelerating case. In this research we are focusing on the accelerating velocity profile.

Polynomial Eigenvalue Problem

By linearizing the governing equations, and assume oscillating perturbed quantities, $\tilde{n}, \tilde{v} \sim \exp(-i\omega t)$. We can derive the following polynomial eigenvalue problem,

$$\begin{aligned} & \omega^2 \tilde{v} \\ & + 2i\omega \left(v_0 \frac{\partial}{\partial z} + \frac{\partial v_0}{\partial z} \right) \tilde{v} \\ & + \left[(1 - v_0^2) \frac{\partial^2}{\partial z^2} - \left(3v_0 + \frac{1}{v_0} \right) \frac{\partial v_0}{\partial z} \frac{\partial}{\partial z} \right. \\ & \quad \left. - \left(1 - \frac{1}{v_0^2} \right) \left(\frac{\partial v_0}{\partial z} \right)^2 - \left(v_0 + \frac{1}{v_0} \right) \frac{\partial^2 v_0}{\partial z^2} \right] \tilde{v} = 0 \end{aligned} \tag{5}$$

Notice that the highest derivative term (labeled in red) vanishes at the throat of the nozzle, $z = 0$. Spectral method fails to resolve eigenmodes because of the existence of this singularity.

Existence of Singularity




Figure 3: Spectral method failed to resolve meaningful eigenfunctions (and eigenvalues) due to the existence of the singularity at $z = 0$.

Shooting Method

- Expand \tilde{v} near the singularity using Frobenius method.
- Pick up regular solutions and shoot them to the left.
- Eigenvalues are found by matching the Dirichlet BC.




Figure 4: The solutions cross the singular point smoothly. All modes are stable.

References I

- [1] T. Aikawa.
The stability of spherically symmetric accretion flows.
Astrophys Space Sci, 66(2):277–285, Dec. 1979.
- [2] H. Bondi.
On Spherically Symmetrical Accretion.
Monthly Notices of the Royal Astronomical Society, 112(2):195–204, Apr. 1952.
- [3] R. L. Burden, J. D. Faires, and A. M. Burden.
Numerical Analysis.
2016.
- [4] F. F. Chen.
Introduction to Plasma Physics and Controlled Fusion.
Springer International Publishing, Cham, 2016.
- [5] H. Furuhashi, Y. Nambu, and H. Saida.
Simulation of an acoustic black hole in a Laval nozzle.
Class. Quantum Grav., 23(17):5417–5438, Sept. 2006.
- [6] K. Jockers.
On the stability of the solar wind.
Sol Phys, 3(4):603–610, Apr. 1968.
- [7] I. D. Kaganovich, A. Smolyakov, Y. Raitses, E. Ahedo, I. G. Mikellides, B. Jorns, F. Taccogna, R. Gueroult, S. Tsikata, A. Bourdon, J.-P. Boeuf, M. Keidar, A. T. Powis, M. Merino, M. Cappelli, K. Hara, J. A. Carlsson, N. J. Fisch, P. Chabert, I. Schweigert, T. Lafleur, K. Matyash, A. V. Khrabrov, R. W. Boswell, and A. Fruchtman.
Physics of $E \times B$ discharges relevant to plasma propulsion and similar technologies.
Physics of Plasmas, 27(12):120601, 12 2020.

References II

- [8] E. Keto.
Stability and solution of the time-dependent Bondi–Parker flow.
Monthly Notices of the Royal Astronomical Society, 493(2):2834–2840, Apr. 2020.
- [9] O. Koshkarov, A. I. Smolyakov, I. D. Kaganovich, and V. I. Ilgisonis.
Ion sound instability driven by the ion flows.
Phys. Plasmas, 22(5):052113, May 2015.
- [10] A. A. Litvak and N. J. Fisch.
Rayleigh instability in Hall thrusters.
Physics of Plasmas, 11(4):1379–1383, Apr. 2004.
- [11] S. Mazouffre.
Electric propulsion for satellites and spacecraft: established technologies and novel approaches.
Plasma Sources Sci. Technol., 25(3):033002, June 2016.
- [12] I. Romadanov, A. Smolyakov, Y. Raitses, I. Kaganovich, T. Tian, and S. Ryzhkov.
Structure of nonlocal gradient-drift instabilities in Hall $E \times B$ discharges.
Physics of Plasmas, 23(12):122111, Dec. 2016.
- [13] D. D. Ryutov, P. N. Yushmanov, D. C. Barnes, and S. V. Putvinski.
Divertor for a linear fusion device.
page 060003, Newport Beach, CA, USA, 2016.
- [14] A. I. Smolyakov, A. Sabo, P. Yushmanov, and S. Putvinskii.
On quasineutral plasma flow in the magnetic nozzle.
Physics of Plasmas, 28(6):060701, June 2021.
- [15] R. F. Stellingwerf and J. Buff.
Stability of astrophysical gas flow. I - Isothermal accretion.
ApJ, 221:661, Apr. 1978.

References III

- [16] L. N. L. N. Trefethen.
Spectral methods in MATLAB.
Software, environments, tools. Society for Industrial and Applied Mathematics, Philadelphia, PA, 2000.
- [17] C. H. Williams.
Fusion Propulsion Through a Magnetic Nozzle and Open Divertor.
In *AIP Conference Proceedings*, volume 654, pages 510–515, Albuquerque, New Mexico (USA), 2003. AIP.
ISSN: 0094243X.