

Focus Fusion: p-B¹¹ Fusion with DPF

Hunt Feng¹

¹Department of Physics and Engineering Physics
University of Saskatchewan

October 23, 2023

Outline of Presentation

- 1 Introduction - p-B¹¹ Fusion
- 2 Theoretical Advances by LPPFusion Researchers

Outline of Presentation

1 Introduction - $p\text{-B}^{11}$ Fusion

2 Theoretical Advances by LPPFusion Researchers

p-B¹¹ as Fusion Fuel

Advantages:

- No neutrons are produced in this reaction, $p + B^{11} \rightarrow 3He^4$.
- Released energy is carried only by charged particles.
- No need to use heat-exchanger, charged particles create electricity directly.
- Only tiny amount of neutrons is produced in secondary reaction, $He^4 + B^{11} \rightarrow N^{14} + n$.
- No radioactive waste.

Challenges:

- Requires average ion energies above 100keV. (DT fusion requires 40keV).
- Plasma density-confinement product $n\tau$ requirement is 15 times higher than DT fusion.
- Boron ions lead far greater amounts of X-ray energy than DT fusion.

Dense Plasma Focus (DPF)

Advantages:

- Extremely compact.
- Simple in construction.
- No need for external magnets nor lasers.
- Utilize the instabilities of plasma rather than fighting them.

Challenges:

- Very bad fusion yield, efficiency 1.25×10^{-5} .

Fusion Yield




Figure 1: Up to peak currents of 1MA, DPF fusion yield rises sharply with increasing current, but plateaus above 1MA. At lower currents, FF-1's performance exceeds those of other DPFs, but was comparable to other best results at 1MA. [5]

E. J. Lerner, S. M. Hassan, I. Karamitsos-Zivkovic, and R. Fritsch. Focus Fusion: Overview of Progress Towards p-B11 Fusion with the Dense Plasma Focus.

Outline of Presentation

1 Introduction - p-B¹¹ Fusion

2 Theoretical Advances by LPPFusion Researchers

Quantitative Model of DPF Functioning

$$r_c = 1.32 \times 10^{-3} (\mu \cdot z)^{-2/3} r \quad (1)$$

$$B_c = 4z \left(\frac{\mu M}{m} \right) B \quad (2)$$

$$n_c = 3.7 \times 10^{10} \frac{\mu^2 z I^2}{r^2} \quad (3)$$

$$Y \sim \mu^{2.75} I^4 f(T_i) \quad (4)$$

where B is peak field at cathode (G), B_c is the field in the core of the plasmoid, r is cathode radius (cm), r_c is the plasmoid core radius, n_c is plasmoid ion density, I is peak current (A), μ is average ionic mass, z is ionic charge, Y is fusion reaction number and $f(T_i)$ is the reaction rate as a function of ion temperature T_i .

Conclusions:

- I^4 scaling is correctly predicted by the model.
- Yield increases with increasing atomic mass of the fill gas.

Quantum Magnetic Field(QMF) Effect

- Problem of high X-ray emission with p-B¹¹ can be mitigated through the use of quantum magnetic field (QMF) effect.
- In strong magnetic field, electrons can have only discrete energy levels, termed Landau levels.

$$E_b = \left(n + \frac{1}{2} \right) \frac{e\hbar B}{mc}$$

- Very little energy can be transferred from ions to electrons by collisions.

Control of Angular Momentum and Efficiency of Energy Transfer to the Plasmoid

- For high efficiency, control of angular momentum is required.
- The plasmoid formation requires certain amount of angular momentum, so that kink instability can occur.
- Angular momentum can be imparted to the plasma sheath during the rundown by the interaction of the inward flowing electron flows and any small initial axial magnetic field. $\mathbf{J} \times \mathbf{B}$ accelerates the electron in the azimuthal direction.

Viscous Heating Mechanism

- Ions in the plasmoid might be heated by viscous heating.
- As plasmoid contracts, ions moving inward at different velocities start to mix. The ordered velocity of motion is converted into random velocity of heat.

$$T_i = 6.2 \times 10^{-4} z_{\text{eff}}^{1.6} n_i^{0.2} (\ln \Lambda L_{\text{max}} B)^{0.4}$$

where $z_{\text{eff}} = (\sum f z^2)^{1/2}$, f is the number fraction of a given ion, z is the ionic charge and the summation is over all ions. z_{eff} is thus a dimensionless number, the effective number of charges per ion. L_{max} is the distance around the plasmoid, which in our model is $9.7(\mu z)^{1/3} L_p$, where L_p is the observed length of the plasmoid core along its axis. T_i is the ion temperature in eV.

- Higher T_i can be expected with higher z fill gas.

Induced Current Heating Mechanism

- Electron beam induces currents in the plasmoid electrons.
- Plasmoid has a great density, this current is distributed over these electrons.
- The slow electrons undergo collisions and converted their kinetic energy to heat.

$$T_e = 0.19(z_{eff}/r_b)^{0.8}(\ln \Lambda LI/z(\gamma - 1))^{0.4}$$

where r_b is the beam radius and γ is the relativistic factor for the beam electrons.

Lower Hybrid Heating Mechanism

- For the densest plasmoids, lower hybrid instability will lead to wave heating of ions by the electron beam.
- For current more than $I > 5.0/\mu^{1/6}z^{5/6}\text{MA}$, the majority of electron beam energy will go to ion heating, rather than electron heating.

Role of Impurities of Disrupting Filaments and Source of Impurities

- High z impurities increase radiation, limiting fusion yield.
- High z impurities decrease the conductivity of the plasma in the current-carrying sheath, leading to disruption of the filaments through over heating.
- High z impurities reduces density (and hence yield) due to asymmetric compression.
- The increasing presence of impurities is hypothesized to be the main reason for the plateauing of fusion yield.



Dense plasma focus — Plasma-Universe.com — plasma-universe.com.
<https://www.plasma-universe.com/dense-plasma-focus/>.
[Accessed 16-10-2023].



DPF Device - LPP Fusion — lppfusion.com.
<https://www.lppfusion.com/technology/focus-fusion-energy/dpf-device/>.
[Accessed 16-10-2023].



R. Behbahani.
Enhancement of Charged Particle Emission from a Plasma Focus Device.
Phd thesis, University of Saskatchewan, 2017.



M. Krishnan.
The dense plasma focus: A versatile dense pinch for diverse applications.
IEEE transactions on plasma science, 40(12):3189–3221, 2012.



E. J. Lerner, S. M. Hassan, I. Karamitsos-Zivkovic, and R. Fritsch.
Focus Fusion: Overview of Progress Towards p-B11 Fusion with the
Dense Plasma Focus.
Journal of Fusion Energy, 42(1), mar 9 2023.