Hunter Lybbert
Student ID: 2426454
11-22-24
AMATH 561

PROBLEM SET 7

Note: Exercises 1-4 are from Matt Lorig's notes (link on course website).

1. Exercise 4.1.
A six-sided die is rolled repeatedly. Which of the following a Markov chains? For those that are, find the one-step transition matrix.

(a) X_n is the largest number rolled up to the nth roll.

Solution:
TODO:

(b) $X - n$ is the number of sixes rolled in the first n rolls.

Solution:
TODO:

(c) At time n, X_n is the time since the last six was rolled.

Solution:
TODO:

(d) At time n, X_n is the time until the next six is rolled.

Solution:
TODO:

2. Exercise 4.2.
Let $Y_n = X_{2n}$. Compute the transition matrix for Y when

(a) X is a simple random walk (i.e., X increases by one with probability p and decreases by 1 with probability q.

Solution:
TODO:

(b) X is a branching process where G is the generating function of the number of offspring from each individual.

Solution:
TODO:

3. Exercise 4.3.
Let X be a Markov chain with state space S and absorbing state k (i.e., $p(k, j) = 0$ for all $j \in S$). Suppose $j \to k$ for all $j \in S$. Show that all states other than k are transient.

Solution:
TODO:

4. Exercise 4.4.
Suppose two distinct states i, j satisfy

$$P(\tau_j < \tau_i | X_0 = i) = P(\tau_i < \tau_j | X_0 = j)$$

where $\tau := \inf\{n \geq 1 : X_n = j\}$. Show that, if $X_0 = i$, the expected number of visits to j prior to re-visiting i is one.

Solution:
TODO:

5. Stationary distribution of Ehrenfest chain. (a) Let X_n be the number of balls in the left urn at time n (total number of balls in both urns is r). At each time step, one of the r balls is picked at random and moved to the other urn.

(a) Let $G_n(s)$ be the generating function of X_n. Derive a formula for G_{n+1} as a function of G_n.

Solution:
TODO:

(b) Let $G(s) = \lim_{n\to\infty} G_n(s)$. Use the relation in part a) to derive an equation for G. Solve it and find G.

Solution:
TODO:

(c) Find the stationary distribution π of Ehrenfest chain. What is the connection between G and π?

Solution:
TODO: