

Hunter Lybbert
Student ID: 2426454
10-28-24
AMATH 567

HOMEWORK 5

Collaborators*:

*Listed in no particular order. And anyone I discussed at least part of one problem with is considered a collaborator.

1: From A&F: 2.6.5

Consider two entire functions with no zeroes and having a ratio equal to unity at infinity. Use Liouville's Theorem to show that they are in fact the same function.

Solution:

(**1 part**, except maybe if there are multiple things to prove here)

2: From A&F: 2.6.10

... deduce

$$f(z) = \frac{1}{2\pi} \int_0^{2\pi} \frac{f(\xi)\xi}{\xi - z} d\theta$$

... explain why we have

$$0 = \frac{1}{2\pi} \int_0^{2\pi} \frac{f(\xi)\xi}{\xi - 1/\bar{z}} d\theta$$

... use something to show

$$f(z) = \frac{1}{2\pi} \int_0^{2\pi} f(\xi) \left(\frac{\xi}{\xi - z} \pm \frac{\bar{z}}{\xi - \bar{z}} \right) d\theta$$

then ...

Solution:

This is a beast of a problem there are **approximately 9** things to show...

3: Suppose Ω is an open simply connected region and $z_0 \in \Omega$. Assume that $f(z)$ is analytic in $\Omega \setminus \{z_0\}$ and satisfies

$$|f(z)| \leq M|z - z_0|^{-\gamma}, \quad \gamma < 1.$$

Show that if the a specific choice for $f(z_0)$ is made then f extends to an analytic function on Ω .

(**1 part**, except maybe if there are multiple things to prove here)

Solution:

4: Establish the following lemma:

Lemma 1

Suppose Ω is an open region and $f(z)$ is continuous on $\overline{\Omega}$. Let Γ be a contour in $\overline{\Omega}$. Suppose a sequence of contours $\Gamma_n \subset \overline{\Omega}$ converge to Γ in the sense that there exists parameterizations $z(t)$ of Γ and $z_n(t)$ of Γ_n defined on $[a, b]$ satisfying

$$\begin{aligned} z_n(t) &\xrightarrow{n \rightarrow \infty} z(t), & \text{uniformly on } [a, b], \\ z'_n(t) &\xrightarrow{n \rightarrow \infty} z'(t), & \text{uniformly on } [a, b]. \end{aligned}$$

Then

$$\int_{\Gamma_n} f(z) dz \xrightarrow{n \rightarrow \infty} \int_{\Gamma} f(z) dz.$$

Hint: Use that f is uniformly continuous on $\overline{\Omega}$.

(1 part, except maybe if there are multiple things to prove here)

Solution:

5: for any $r, R > 0$, let $C = \partial\Sigma$, $\Sigma = \{z \in \mathbb{C} : |\operatorname{Re} z| \leq r \text{ and } 0 \leq -\operatorname{Im} z \leq R, R > 0\}$. In this problem \sqrt{z} denotes the principal branch with $\arg z \in [-\pi, \pi)$.

- Show that if $f(z)$ is analytic in a region that contains Σ ,

$$\oint_C f(z) \sqrt{z-1} \sqrt{z+1} dz = 0.$$

(1 part)

Solution:

- Show that if $f(z)$ is analytic in a region that contains Σ

$$\oint_C \frac{f(z) dz}{\sqrt{z-1} \sqrt{z+1}} = 0.$$

(1 part)

Solution:

6: From A&F: 3.1.1 b,d

In the following we are given sequences. Discuss their limits and whether the convergence is uniform, in the region $\alpha \leq |z| \leq \beta$, for finite $\alpha, \beta > 0$.

b)

$$\left\{ \frac{1}{z^n} \right\}_{n=1}^{\infty}$$

(2 parts)

Solution:

d)

$$\left\{ \frac{1}{1 + (nz)^2} \right\}_{n=1}^{\infty}$$

(2 parts)

Solution:

7: From A&F: 3.1.2 b,d

For each sequence in problem 1, what can be said if

(a) $\alpha = 0$

(b) $\alpha > 0, \quad \beta = \infty$

(4 parts 2x2) *Solution:*

8: From A&F: 3.1.3 Compute the integrals

$$\lim_{n \rightarrow \infty} \int_0^1 n z^{n-1} dz \quad \text{and} \quad \int_0^1 \lim_{n \rightarrow \infty} (n z^{n-1}) dz$$

and show that they are not equal. Explain why this is not a counter example to Theorem 3.1.1. (A &F pg. 111)

(3 parts) *Solution:*

There are approximately 25 things to do