

Notes on Introductory Series

John Hunter Kohler

March 2020

Though the formal definition of sequences and series limits will not be given. The notation used will be standard and informal. No proofs will be given similar to the entirety of this document. Assume all sequences and series are real.

Foundation of Sequences

This section focuses on the basics of sequences and methods of computation on the infinite terms of some sequence. A sequence $\{a_n\}$ is said to be convergent if

$$\lim_{n \rightarrow \infty} a_n \in \mathbb{R} \quad (1)$$

If the above property is not held, the sequence is said to be divergent. When computing limits of two convergent sequences $\{a_n\}, \{b_n\}$, it is possible to separately evaluate the sequences.

$$\lim_{n \rightarrow \infty} f(a_n, b_n) = f(\lim_{n \rightarrow \infty} a_n, \lim_{n \rightarrow \infty} b_n) \quad (2)$$

A simple tool for the evaluation of limits upon convergent series is the squeeze theorem for series.

$$a_n \leq c_n \leq b_n, \text{ for } n > N \quad (N \in \mathbb{N}) \quad (3)$$

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = L \implies \lim_{n \rightarrow \infty} c_n = L \quad (4)$$

Sequences, like functions, are said to be monotonic if they are strictly increasing or strictly decreasing. Like functions, sequences may be bounded above or below by some constant. If a sequence is bounded and monotonic, then it must be convergent¹.

¹Obviously, the same can be said about those sequences which are simply strictly increasing and bounded above as well as strictly decreasing and bounded below.

Convergence of Series

Here, let us take a look at simple rules for figuring if a series converges or diverges. Obviously a convergent series is the sum of every term of a series such that the value is finite. If some series does not exist or is infinite, it is said to be divergent. A simple proposition is that

$$\text{If } \sum a_n \text{ converges, then } \lim_{n \rightarrow \infty} a_n = 0. \quad (5)$$

Absolute and Conditional Convergence

Now recognize two more properties of series. If some series $\sum a_n$ is convergent yet $\sum |a_n|$ is not, then the series $\sum a_n$ is said to be conditionally convergent. If, along with $\sum |a_n|$, $\sum a_n$ is convergent, we say that the series is absolutely convergent. Otherwise, the series is divergent. From this it is important to recognize one thing: if a series is conditionally convergent, then when algebraically rearranging² its explicit terms will allow you to, without logical error, compute the sum of the series as any real number.

Geometric Sums and P-Series

Firstly define a partial sum -this usually uses confusing notation- as

$$S_n := \sum_{i=0}^n a_i \quad (6)$$

The lower bound of the sum may change based on context, but this is the general notation with the exception of the differing use of capitalization of s . For a geometric sum, based on a sequence where the ratio between each term is a constant, the partial sum formula is slightly unintuitive, though important. It is vital in the proof of the infinite sum, though that will not be exemplified.

$$\text{for } a, r \text{ constant } S_n = \sum_{i=0}^n ar^i = \frac{a(1 - r^{n+1})}{1 - r} \quad (7)$$

$$\lim_{n \rightarrow \infty} S_n = \frac{a}{1 - r} \text{ when } |r| < 1 \quad (8)$$

As you will see later, this is a form of a power series, with radius of convergence³ one. The other series to look at of the form for which we know

²For the purposes of computation.

³Put simply, as expressed above, the infinite geometric sum only converges for $|r| < 1$.

convergence is the p-series ⁴.

$$\sum_{n=1}^{\infty} \frac{1}{n^p} \text{ converges for } p > 1 \quad (9)$$

Integral Test

If there is some continuous, positive, strictly decreasing function $f(x)$ on,

$$x \in [k, \infty) \text{ and } f(n) = a_n, n \in \mathbb{N}, \text{ then} \quad (10)$$

$$\int_k^{\infty} f(x)dx \text{ and } \sum_{n=k}^{\infty} a_n \text{ converge or diverge together.} \quad (11)$$

Comparison Test

This test is similar to, and essentially a lemma⁵ of, the last theorem:

$$\text{If there is some } \sum a_n, \sum b_n \text{ for } a_n \geq b_n \geq 0, a_n \leq b_n, n \in \mathbb{N}, \quad (12)$$

$$\text{then } \sum b_n \text{ converges } \implies \sum a_n \text{ converges,} \quad (13)$$

$$\text{and } \sum a_n \text{ diverges } \implies \sum b_n \text{ diverges.} \quad (14)$$

Limit Comparison Test

By taking the associated sequences of two series and compare their infinite terms, a similar association of convergence as before can be formed⁶.

$$\text{Let } \sum a_n, \sum b_n \text{ such that } a_n \geq 0, b_n > 0, \quad (15)$$

$$n \in \mathbb{N}, \text{ and let } C := \lim_{n \rightarrow \infty} \frac{a_n}{b_n} \quad (16)$$

$$\text{If } C \in \mathbb{R}^+ \text{ then } \sum a_n \text{ and } \sum b_n \quad (17)$$

$$\text{converge or diverge together.} \quad (18)$$

⁴The special case is $p = 1$ which is the harmonic series. Not discussed here.

⁵Due to the fact an integral would simply be the application of this theorem through its definition of a Riemann Sum.

⁶These associations will let us extend knowledge of convergence from what we have proved to unknowns more easily. Also this would give the option of working the more simple of the two series.

Alternating Series Test

To deal with series of the form below, this test is created. Once used, if it is possible to do so, another test may be required to determine the properties of the subseries. An alternating series is of the form:

$$a_n = (-1)^n b_n \text{ or } a_n = (-1)^{n+1} b_n, \text{ for } (b_n \geq 0 \text{ or } b_n \leq 0)^7, n \in \mathbb{N} \quad (19)$$

Furthermore, the test states:

$$\text{If } \lim_{n \rightarrow \infty} b_n = 0 \text{ and } \{b_n\} \text{ is monotonic}^7 \sum_{n=1}^{\infty} a_n \text{ is convergent.} \quad (20)$$

Ratio Test

The ratio test will be the best tool presented here to deal with sequences involving factorials as well as rational equations of polynomials. By reviewing the test below you will see that it takes a form in which, for the described cases, we may cancel out confounding terms.

$$\text{Let } L := \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| \quad (21)$$

$$\text{If } L < 1, \sum a_n \text{ is absolutely convergent.} \quad (22)$$

$$\text{If } L > 1, \sum a_n \text{ is divergent.} \quad (23)$$

$$\text{If } L = 1, \text{ the test is inconclusive.} \quad (24)$$

Root Test

This test is similar to the ratio test. Almost⁸ all sequences produce the same results in the ratio and root test. This test is best utilized for series similar to $a_n = (b_n)^n$. The test functions thus:

$$\text{Let } L := \lim_{n \rightarrow \infty} |a_n|^{\frac{1}{n}} \quad (25)$$

$$\text{If } L < 1, \sum a_n \text{ is absolutely convergent.} \quad (26)$$

$$\text{If } L > 1, \sum a_n \text{ is divergent.} \quad (27)$$

$$\text{If } L = 1, \text{ the test is inconclusive.} \quad (28)$$

⁷For $(b_n \geq 0 \text{ and strictly decreasing})$, or $(b_n \leq 0 \text{ and strictly increasing})$. Note that $\sum b_n$ does not have to converge.

⁸Most cases that do not abide by this principle should not be presented in an introductory course. It may be worthwhile checking both either way as either one may have been applied incorrectly

Series Estimation

Bounding by Integration

Utilizing function integrals, we can find bounds for the value of an infinite series. Even if the representation of a series can not be integrated, we know enough integral approximation techniques to approximate the improper integrals. Using notation of the partial sum S_n , we may manually compute the partial sum of a series up to some point. Next, define the E_n as below as what we call the remaining sum -for obvious reasons-.

$$E_n := S - S_n = \sum_{i=0}^{\infty} a_i - \sum_{i=0}^n a_i = \sum_{i=n+1}^{\infty} a_i \quad (29)$$

Now it is possible to bound E_n by way of integration for some function $f(x)$ that satisfies all the requirements of the Integral Test. To complete the series bound, simply recognize the following:

$$\int_{n+1}^{\infty} f(x) dx \leq E_n = S - S_n \leq \int_n^{\infty} f(x) dx \quad (30)$$

$$S_n + \int_{n+1}^{\infty} f(x) dx \leq S \leq S_n + \int_n^{\infty} f(x) dx \quad (31)$$

One more extension can be made to this method; namely that if we utilize a series b_n that functions, in relation to a_n , similarly to our convergence test by comparison. This can make the computation of the approximation easier by trading in accuracy.

Approximating Alternating Series

The equation given below is derived from the proof of the Alternating Series Test. It should be intuitive however due to, again mirroring the conditions of the test, b_n must be strictly decreasing and positive, or the inverse. Understand that b_n is our subseries of the alternating series. The sum and partial sums below are of the series of interest a_n .

$$|E_n| = |S - S_n| \leq b_{n+1} \quad (32)$$

It is important to realize that E_n is simply the error of our partial sum. From above we know that we can bound the error and therefore simply bound the total sum of the sequence:

$$S_n - |E_n| \leq S \leq S_n + |E_n| \quad (33)$$

Bounding by Ratio

As goes all approximations in this section, we must have some a_n that follows the requirements of the corresponding test -here, the ratio test-. The formulas below are notably useful for bounding E_n for sequences which are best analyzed with the Ratio Test. A last obvious precondition is that a_n converges.

$$\text{Define a new geometric sequence by: } r_n := \frac{a_{n+1}}{a_n} \quad (34)$$

$$\text{Case 1: Sequence } r_n \text{ is decreasing.} \quad (35)$$

$$|E_n| \leq \frac{a_{n+1}}{1 - r_{n+1}} \quad (36)$$

$$\text{Case 2: Sequence } r_n \text{ is increasing.} \quad (37)$$

$$|E_n| \leq \frac{a_{n+1}}{1 - L} \quad (38)$$

These are the only two cases that happen due to $\{r_n\}$ being convergent if its ratio⁹ $r_n < 1$. In case two, though the geometric series $\{r_n\}$ has a ratio greater than 1, we know that as n goes to infinity, that this become less than one, so we use L as the general ratio⁹.

Power Series

The general¹⁰ power series is thus for a constant and sequence $\{c_n\}$:

$$p(x) = \sum_{n=0}^{\infty} c_n (x - a)^n \quad (39)$$

Let us call $\{c_n\}$ the sequence of coefficients of $p(x)$. Briefly mentioned in the section on geometric series, the radius of convergence R is the bound¹¹ on x for which $p(x)$ converges. Furthermore, we may call the range of x for which $p(x)$ converges, its interval of convergence. When attempting to find the radius of convergence for some series, the end goal tends to be to find some inequality involving x in solitude. To do this one would use one

⁹When speaking about the ratio of $\{r_n\}$, it is meant that for some n , namely that for which we choose E_n , has such a ratio

¹⁰This is the general power series along \mathbb{R} . Things such as radius of convergence make more sense when extended to the complex generalization; for example, this radius truly refers to the disk in \mathbb{C} for which some series representation of $p(z)$ converges

¹¹Meaning that $|x| \leq R$ or $|x| < R$.

or multiple of the tests described in the section of series convergence. The ratio test is a particularly effective tool in this case. It is not always the case that $|x| = R$ is not in the interval of convergence, thus one must find both R and check the endpoints of said interval. Note that R is in $\overline{\mathbb{R}}$, though x only varies along real numbers. This implies two important points. The first of which is x will have an interval of all reals for $R = \infty$. Next, even in the case of $R = 0$, the interval of x will always be non null due to $p(a) = 0$.

Expressing Analytic Functions

We will not describe the definition of an analytic function beyond saying that they are functions that can be expressed locally as a power series. Below¹² are some frequent analytic functions¹³ described by their power series.

$$(1+x)^k = \sum_{n=0}^{\infty} \binom{k}{n} x^n = 1 + kx + \frac{k(k-1)}{2!} + \dots \quad |x| < 1 \quad (40)$$

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \frac{x^6}{6!} \dots \quad x \in \mathbb{R} \quad (41)$$

$$\sin(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots \quad x \in \mathbb{R} \quad (42)$$

$$\cos(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \dots \quad x \in \mathbb{R} \quad (43)$$

$$\sinh(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!} = x + \frac{x^3}{3!} + \frac{x^5}{5!} + \frac{x^7}{7!} + \dots \quad x \in \mathbb{R} \quad (44)$$

$$\cosh(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \frac{x^6}{6!} + \frac{x^8}{8!} + \dots \quad x \in \mathbb{R} \quad (45)$$

The next step is to bring calculus into our representations. Obviously the integral or derivative of a series is simply that of each individual term. Our first theorem is thus:

$$\text{For some differentiable, continuous } f(x) = \sum_{n=0}^{\infty} c_n(x-a)^n \quad (46)$$

$$\frac{df}{dx} \text{ and } \int f(x) dx \text{ have the equivalent } R \text{ to } f(x) \quad (47)$$

¹²All will be easier to prove and recognize after learning the Taylor series. Note that the series below are Maclauren series.

¹³In non general form for the purposes of simplicity.

Recognize in many situations it may be beneficial to take the n th derivative of a function so to find a easily recognizable power series, then integrate n times -while solving for the integration constants- to find the power series of the original function. A simple example is the power series of $\ln(1-x)$. In the next section we will learn how to find the power series of this easily, but we may also find it through simply differentiating it so that it is represented as the geometric series.

Taylor Series

Put simply, the Taylor series generalizes a power series to any infinitely differentiable function. These series still have radii of convergence and are said to be centered around some a ¹⁴. Assume $f(x)$ is an infinitely differentiable function that is continuous and differentiable at a .

$$f(x) = \sum_{i=0}^{\infty} \frac{f^{(i)}(a)}{i!} (x-a)^i = f(a) + f'(a)(x-a) + \dots \quad (48)$$

Note that the Taylor series is simply the semi-general power series described with $\{c_n\} = \{f^{(n)}(a)\}$.¹⁵ As described before, this equation only holds true for $x \in R_a$, where we say R_a is the radius of convergence around a . If we have some series, we can approximate the value of $f(x)$, $x \in R_a$ to any level of specificity. This is extremely useful in the cases of transcendental functions that cannot be calculated. I will again mention that a Taylor series around $a = 0$ is referred to as a Maclaurin series. To finish, one piece of frequent notation is that of the n th degree of a Taylor series denoted by T_n .

$$T_n = \sum_{i=0}^n \frac{f^{(i)}(a)}{i!} (x-a)^i = f(a) + f'(a)(x-a) + \dots + \frac{f^{(n)}(a)}{n!} (x-a)^n \quad (49)$$

¹⁴This central value -this again makes more sense over a complex field - will also be notated frequently as x_0 . See the below that a is simply some value on the domain of f for which we will find $f(a)$.

¹⁵To generalize this series we obviously must find a general form for $f^{(n)}$, which is not always the easiest task