AS · Cambridge (CIE) · Physics

25 mins 25 questions

Multiple Choice Questions

Stress & Strain

Extension & Compression / Hooke's Law / The Young Modulus

Easy (9 questions)	/9
Medium (10 questions)	/10
Hard (6 questions)	/6
Total Marks	**/25**

Scan here to return to the course
or visit savemyexams.com

© 2026 Save My Exams, Ltd.

Easy Questions

1 A student wanted to measure the Young modulus of a wire, to do this they needed to take a number of measurements.

In which row could the measurement **not** be made **directly** with the listed apparatus?

	Measurement	Apparatus
A	extension of wire	vernier scale
B	area of cross-section of wire	micrometer screw gauge
C	original length of wire	metre rule
D	mass of load applied to a wire	electronic balance

(1 mark)

2 What is the unit of the Young modulus?

A. $N\ m^{-2}$

B. $N\ m$

C. $N\ m^{-1}$

D. $N\ m^{2}$

(1 mark)

© 2026 Save My Exams, Ltd.

3 A material is extended elastically within the limit of proportionality. What is equal to the Young modulus of the material?

A. gradient of the stress-strain graph

B. area under the force-extension graph

C. area under the stress-strain graph

D. gradient of the force-extension graph

(1 mark)

4 An experiment was carried out to determine the Young modulus of steel using a steel wire.

The uncertainties in the measurements are given below.

Measurement	Uncertainty
load on wire	±2%
length of wire	±0.2%
diameter of wire	±1.5%
extension	±1%

What is the percentage uncertainty in the Young modulus?

A. 1.3%

B. 1.8%


C. 4.7%

D. 6.2%

(1 mark)

© 2026 Save My Exams, Ltd.

5 A glass rod was put under stress and strain until it broke. The graph of this is shown below.



Which of the following statements is correct about the glass rod?

A. the glass shows plastic deformation

B. when the cross-sectional area of the rod is doubled, the ultimate tensile stress of the rod is halved

C. Hooke's law is obeyed for all values of stress up to the breaking point


D. the glass is ductile

(1 mark)

6 Two wires S and T, with the same initial dimensions, were put under stress and strain, the

© 2026 Save My Exams, Ltd.

graph shows the results up to their breaking points.



Which statement is **not** correct?

A. material S has a larger ultimate tensile stress

B. material S extends elastically

C. material S extends more than material T when loaded with the same force


D. material S is brittle

(1 mark)

7 The graphs below show the stress-strain graphs of three materials. The graphs do not have

© 2026 Save My Exams, Ltd. | Get more and ace your exams at savemyexams.com

the same scales.



The three materials are copper, rubber and glass.


Which materials are represented by the graphs?

	1	2	3
A	rubber	copper	glass
B	rubber	glass	copper
C	glass	copper	rubber
D	copper	glass	rubber

(1 mark)

© 2026 Save My Exams, Ltd.

8 The stress-strain graph for bone is shown below.



What is the Young modulus of bone?

A. 2×10^6 N m^{-2}

B. 2×10^8 N m^{-2}

C. 1×10^6 N m^{-2}

D. 1×10^8 N m^{-2}

(1 mark)

9 A student wanted to determine the spring constant of a spring. They did this by using different loads to extend the spring by varying amounts.

To find the spring constant, which quantities are required?

A. the elastic limit, extensions and the length of the spring

B. the loads and the extensions of the spring

C. the elastic limit and the loads


D. the loads and the length of the spring

(1 mark)

© 2026 Save My Exams, Ltd. | Get more and ace your exams at savemyexams.com

Medium Questions

1 A metal tube of diameter d with a thin wall thickness W is shown in the diagram. The thickness of the wall is small when compared to the diameter of the tube.



A force of magnitude T is applied parallel to the axis of the tube, putting the tube under tension. It is proposed that making the wall thicker would reduce the stress on the tube.

If the tube diameter and the tension remain the same, which wall thickness would half the stress?

A. $\frac{W}{2}$

B. $\sqrt{2}W$

C. $2W$

D. $4W$

(1 mark)

2 A steel metal wire has the following properties:

© 2026 Save My Exams, Ltd. | Get more and ace your exams at savemyexams.com

diameter = 5.0×10^{-4} m

Young modulus = 2.0×10^{11} Pa

tension = 20 N

The string snaps and contracts elastically.

By what percentage does a length *l* of a piece of the string contract?

A. $1.3 \times 10^{-4}\%$

B. $5.1 \times 10^{-4}\%$

C. $1.3 \times 10^{-2}\%$

D. $5.1 \times 10^{-2}\%$

(1 mark)

3 The Young modulus of a metal wire is dependent on which property?

A. spring constant

B. ductility

C. ultimate tensile stress

D. elastic limit

(1 mark)

4 The Young modulus *E* and the force per unit extension *k* describes the behaviour of a wire under tensile stress.

For a wire of length *L* and cross-sectional area *A*, what is the relationship between *E* and *k* ?

A. $E = \frac{A}{kL}$

B. $E = \frac{kA}{L}$


C. $E = \frac{kL}{A}$

© 2026 Save My Exams, Ltd.

D. $E = \frac{L}{kA}$

(1 mark)

5 A steel bar, shown in the diagram, has a circular cross-section that is under tension *T*.



The diameter of the thinner portion is half the wider portion.

What is the value of $\frac{stress\ on\ the\ wide\ portion}{stress\ on\ the\ narrow\ portion}$?

A. 4.0

B. 2.0

C. 0.50

D. 0.25

(1 mark)

6 A student wanted to find the young modulus of a wire. The equation for the young modulus *E* is:

$$E = \frac{4Fl}{\pi d^2 x}$$

The student extended the wire with a known force and made a series of measurements.


Which measurement has the largest effect on the uncertainty in the value of the calculated Young modulus?

© 2026 Save My Exams, Ltd. | Get more and ace your exams at savemyexams.com

	Measurement	Symbol	Value
A	length of wire before force applied	l	2.043 ± 0.002 m
B	force applied	F	19.62 ± 0.01 N
C	extension of wire with force applied	x	5.2 ± 0.2 mm
D	diameter of wire	d	0.54 ± 0.02 mm

(1 mark)

7 A spring was stretched with increasing load. The graph of the results is shown below.



What is the spring constant?

A. 0.080 N m^{-1}

B. 0.13 N m^{-1}


© 2026 Save My Exams, Ltd.

C. 2.7 N m^{-1}

D. 8.0 N m^{-1}

(1 mark)

8 The graph below shows the stress-strain for a metal.



When the strain on the rod is 0.010, what is the strain energy per unit volume of a rod made from this metal?

A. 10 kJ m^{-3}

B. 10 MJ m^{-3}

C. 100 kJ m^{-3}

D. 1.0 MJ m^{-3}

(1 mark)

9 A steel spring was stretched to a length of 55 cm when a 25 N weight was hung from it. The spring has a spring constant of 150 N m^{-1}.

What was the original length of the spring?

A. 0.61 m

B. 0.72 m

C. 0.38 m

D. 0.49 m

© 2026 Save My Exams, Ltd.

(1 mark)

10 A steel cable used in a construction crane has a Young's modulus of $2.1 \times 10^9\,\text{Pa}$. The cable has a cross-sectional area of $0.05\,\text{mm}^2$ and is subjected to a tensile force of $210\,\text{N}$. The cable obeys Hooke's law.

What is the stress in the steel cable?

A. $4.2 \times 10^3\,\text{Pa}$

B. $4.2 \times 10^6\,\text{Pa}$

C. $4.2 \times 10^9\,\text{Pa}$


D. $4.2 \times 10^{12}\,\text{Pa}$

(1 mark)

© 2026 Save My Exams, Ltd.

Hard Questions

1 A glass-reinforced plastic rod and a nylon rod attached, as shown, was used to make a composite rod.



Each rod has a length of 1.00 m and the same cross-sectional area. The Young modulus E_n of the nylon is 2.0 GPa, and the Young modulus E_p of the plastic is 40 GPa

The composite rod will break when its total extension reaches 3.0 mm.

What is the greatest tensile stress that can be applied to the composite rod before it breaks?

A. 5.7×10^{6} Pa

B. 5.7×10^{9} Pa

C. 7.1×10^{-14} Pa


D. 7.1×10^{-2} Pa

(1 mark)

2 A child with a mass of 40 kg was standing on a diving board. The diving board has a length of 5.0 m and is hinged at one end then supported by a spring 2.0 m from the hinged end. The

© 2026 Save My Exams, Ltd.

spring has a spring constant of 10 kN m^{-1}



What is the extra compression of the spring caused by the child standing on the end of the board?

A. 16 cm

B. 9.8 cm

C. 1.6 cm


D. 1.0 cm

(1 mark)

3 The diagram shows a metal cube is placed between two board and compressed elastically

© 2026 Save My Exams, Ltd. | Get more and ace your exams at savemyexams.com

by two opposing forces *F*. The cube has a length of each side of *l*



How will Δ*l*, the amount of compression, relate to *l* ?

A. $\Delta l \propto l$

B. $\Delta l \propto l^2$

C. $\Delta l \propto \frac{1}{l^2}$

D. $\Delta l \propto \frac{1}{l}$

(1 mark)

4 A metal wire that is supported vertically from a fixed point has a load of 84 N applied to the lower end. The wire has a cross-sectional area of 0.02 mm^2 and obeys Hooke's law. The length of the wire increases by 0.30 %.

What is the Young modulus of the metal wire?

A. 1.4×10^5 Pa

B. 1.4×10^8 Pa


C. 1.4×10^9 Pa

D. 1.4×10^{12} Pa

© 2026 Save My Exams, Ltd. | Get more and ace your exams at savemyexams.com

(1 mark)

5 Two wires S and T is joined, S has a length of 3.0 m and T a length of 1.0 m, both wires have a uniform cross-sectional area.



When a load was hung from the wires S stretched by 1.5 mm and T by 1.0 mm.

The same load is then hung from a second wire of the same cross-section, consisting of 1.0 m of metal S and 3.0 m of metal T.

What is the total extension of this second wire?

A. 3.5 mm

B. 2.5 mm

C. 5.0 mm

D. 4.8 mm

(1 mark)

6 Two steel wires X and Y that both obey Hooke's law were used in an experiment. Wire X has a length l and a cross-sectional area A. Wire Y has a length of $2l$ and cross-sectional area of

© 2026 Save My Exams, Ltd. | Get more and ace your exams at savemyexams.com

$\frac{A}{2}$.

What is the ratio $\frac{tension\ in\ X}{tension\ in\ Y}$ when both wires are stretched to the same extension?

A. $\frac{1}{4}$

B. $\frac{4}{1}$

C. $\frac{2}{1}$

D. $\frac{1}{2}$

(1 mark)

© 2026 Save My Exams, Ltd.