Cambridge Assessment International Education

Cambridge International AS & A Level

BIOLOGY

9700/08

Paper 1 Multiple Choice – Transport of Oxygen &

MARK SCHEME

Maximum Mark: 30

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the 2026 series for most Cambridge IGCSE™, Cambridge International A and AS Level components, and some Cambridge O Level components.

This document consists of 4 printed pages.

© UCLES 2026

Question	Answer	Marks
1	**A** ; *Option A correctly fulfills the Cambridge 9700 learning criteria for Transport of Oxygen &. In accordance with syllabus standards, the distractors represent common misconceptions or inverted relationships.*	1
2	**B** ; *Option B correctly fulfills the Cambridge 9700 learning criteria for Transport of Oxygen &. In accordance with syllabus standards, the distractors represent common misconceptions or inverted relationships.*	1
3	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
4	**D** ; *Option D correctly fulfills the Cambridge 9700 learning criteria for Transport of Oxygen &. In accordance with syllabus standards, the distractors represent common misconceptions or inverted relationships.*	1
5	**A** ; *Option A correctly fulfills the Cambridge 9700 learning criteria for Transport of Oxygen &. In accordance with syllabus standards, the distractors represent common misconceptions or inverted relationships.*	1
6	**B** ; *Option B correctly fulfills the Cambridge 9700 learning criteria for Transport of Oxygen &. In accordance with syllabus standards, the distractors represent common misconceptions or inverted relationships.*	1
7	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
8	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
9	**A** ; *Option A correctly fulfills the Cambridge 9700 learning criteria for Transport of Oxygen &. In accordance with syllabus standards, the distractors represent common misconceptions or inverted relationships.*	1
10	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
1	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
2	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1

© UCLES 2026

Question	Answer	Marks
3	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
4	**B** ; *The membrane is described as 'fluid' because phospholipids and proteins move laterally within the monolayer, and 'mosaic' because protein molecules are scattered within the phospholipid bilayer.*	1
5	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
6	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
7	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
8	**D** ; *Option D correctly fulfills the Cambridge 9700 learning criteria for Transport of Oxygen &. In accordance with syllabus standards, the distractors represent common misconceptions or inverted relationships.*	1
9	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
10	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
1	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
2	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
3	**B** ; *Option B correctly fulfills the Cambridge 9700 learning criteria for Transport of Oxygen &. In accordance with syllabus standards, the distractors represent common misconceptions or inverted relationships.*	1
4	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1

© UCLES 2026

Question	Answer	Marks
5	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
6	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
7	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
8	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
9	**C** ; *High partial pressure of CO2 (and lower pH) shifts the oxygen dissociation curve to the right (the Bohr effect), decreasing haemoglobin's affinity for O2 and promoting oxygen unloading at respiring tissues.*	1
10	**C** ; *Option C correctly fulfills the Cambridge 9700 learning criteria for Transport of Oxygen &. In accordance with syllabus standards, the distractors represent common misconceptions or inverted relationships.*	1

© UCLES 2026