Transport in plants – AS 9700 Biology June 2022

1. June/2022/Paper_11/No.26

Which row is correct for the movement of water in a root?

	pathway	molecule present in Casparian strip
A	apoplast pathway through intercellular spaces	suberin
B	apoplast pathway through plasmodesmata	lignin
C	symplast pathway through plasmodesmata	lignin
D	symplast pathway through intercellular spaces	suberin

apoplast pathway is through intercelular spaces

Symplast takes the pathways through the plasmodesmata

2. June/2022/Paper_11/No.27

The table contains some information about uptake and movement of water and of mineral ions in plants.

	water	mineral ions
mechanism for uptake	osmosis	diffusion, active transport
site of uptake	root hair cells	root hair cells
mechanism of release	evaporation, diffusion	diffusion, active transport
site of release	stomata	plant cells

Using the information provided, which factors will affect the uptake and movement of water or of mineral ions in plants?

1 humidity – Low humidity faster transpiration

2 surface area of root hair cell – increase water uptake in plants

3 oxygen concentration – Increased respiration produce energy for active transport

4 temperature – transpiration (water loss at leaves) is faster temperature increase evaporate rate. This reduce rate of photosynthesis

A 1, 2, 3 and 4

B 1, 3 and 4 only

C 1 and 3 only

D 4 only

PapaCambridge

3. June/2022/Paper_11/No.28

Which changes to the water potential and the volume of solution in the phloem sieve tube occur when sucrose is moved from a photosynthesising leaf into the phloem sieve tube?

	water potential in the phloem sieve tube becomes	volume of solution in the phloem sieve tube
A	higher	decreases
B	higher	increases
C	lower	decreases
D	lower	increases

Moving sucrose decrease water potential in the sieve tubes of phloem. Therefore water moves into phloem sieve tube

4. June/2022/Paper_11/No.29

A student wrote the following statements about a possible mechanism for loading sucrose from a source.

1 When energy is released from ATP, the released energy is used to move sucrose through a co-transporter protein in the companion cell membrane. - false - Co-transporter protein transport is passive

2 As sucrose is moved into a companion cell the pH in the cell wall of the companion cell decreases. Moving H+ ions affect pH; H+ comes along with sucrose decreased H+ increase acidity

3 Proton pumps in the cell membrane of a companion cell move sucrose into the phloem sieve tube element. - false movement of sucrose by co-transporter protein does not use energy (passive)

Which statements are correct?

A 1, 2 and 3 B 1 and 2 only C 1 and 3 only D 2 only

energy is used to pump H+

PapaCambridge

5. June/2022/Paper_12/No.24

Some of the features present in transport tissues are listed.

1 ✓ lignified walls – Lignin reinforce walls of xylem tissues

2 cytoplasm – xylem cells are hollow and have no nucleus

3 mitochondria – no cell organelles in xylem – they are made of dead tissue

4 ✓ pits – allows for water to move across the xylem tissues.

5 plasmodesmata

Which features are present in xylem vessel elements?

A 1, 2, 3 and 4

B 1, 4 and 5

C ✓ 1 and 4 only

D 2, 3 and 5

6. June/2022/Paper_12/No.25

5 The diagrams show transverse sections through parts of plants.



phloem transports sugars from source (leaf) to sink (storage organ)

Which row is correct?

	contains lignin	transports organic solutes
A	1, 4, 5	2, 3, 6
B	1, 4, 6	2, 3, 5
C	2, 3, 5	1, 4, 6
D	2, 4, 6	1, 3, 5

xylem is reinforced by lignin

7. June/2022/Paper_12/No.26

Which molecules form hydrogen bonds with water during transpiration?

1 ✓ cellulose in the xylem wall

2 suberin in the xylem wall – suberin present in endodermis

3 ✓ other water molecules in the xylem

A 1, 2 and 3 B 1 and 2 only C 1 and 3 only D 2 and 3 only

PapaCambridge

8. June/2022/Paper_12/No.27

Some plant species can take up heavy metal contaminants that are dissolved in soil water and then transport them within the plant. Within plant cells, the heavy metals accumulate mainly in the vacuole.

tonoplast is membrane of vacuole

Which suggestions about the transport and accumulation of heavy metals are valid?

1 After initial entry into the root, some of the heavy metals can pass through the tonoplast to be stored in the vacuole of cells in the cortex.

2 The heavy metals take an apoplastic pathway in the xylem but at the endodermis must take a symplastic pathway.

3 The rate of accumulation of the heavy metals in leaf cells will be faster at night, when photosynthesis is not occurring, than during the day.

photosynthesis increases rate of transpiration.

4 The presence of heavy metals causes the transpiration stream to slow down and reduce the rate of transpiration.

A 1 and 2 B 1 and 4 C 2 and 3 D 3 and 4

9. June/2022/Paper_12/No.28

What is the correct route for the movement of water from cell to cell in the apoplast pathway?

apoplastic path exist between two walls - intracellular

A through adjacent cell surface membranes

B through intercellular spaces

C through plasmodesmata - intracellular

D through the Casparian strip

10. June/2022/Paper_12/No.29

Which row shows the correct sequence for the movement of sucrose into phloem sieve tubes?

Pump H^+ into companion cell passively

	first	second	third
A	diffusion of sucrose into the companion cell cytoplasm	active transport of protons into the companion cell cytoplasm	cotransport of protons and sucrose into the sieve tubes
B	cotransport of protons and sucrose into the companion cell cytoplasm	active transport of protons out of the companion cell cytoplasm	diffusion of sucrose into the sieve tubes
C	active transport of protons into the companion cell cytoplasm	diffusion of sucrose into the companion cell cytoplasm	cotransport of protons and sucrose into the sieve tubes
D	active transport of protons out of the companion cell cytoplasm	cotransport of protons and sucrose into the companion cell cytoplasm	diffusion of sucrose into the sieve tubes

PapaCambridge

11. June/2022/Paper_12/No.39

A person's blood group is determined by antigens present on the red blood cells. The table shows the antigens and antibodies in the blood of people with different blood groups.

blood group	presence of A or B antigens on red blood cells	presence of antibodies to A or B in plasma
A	A only	anti-B only
B	B only	anti-A only
AB	A and B	neither
O	neither	anti-A and anti-B

During a blood transfusion, it is essential that the person who receives the blood does **not** have antibodies to the donor's blood.

AB has no antibodies (universal recipient) O can recieve from O only but is universal donor

Which blood groups can be given to a person with blood group B?

A A and B **B** AB and B **C** AB and O **D** B and O


Recipient must not have antibodies against donors blood

PapaCambridge

12. June/2022/Paper_13/No.26


The diagram shows a plant organ.

Which letter correctly labels the xylem?



13. June/2022/Paper_13/No.27

The photomicrograph shows a vascular bundle found in a plant organ.



Companion cells beside sieve tube cells have lots of mitochondrion for energy production.

Which statements about this vascular bundle are correct?

1 The vascular bundle is part of the structure of a root. - from stem

2 Some of the cells in region X have very large numbers of mitochondria.

3 Region Y is made up of a number of different cell types. sclerenchyma

A 1 and 2 B 1 only C 2 and 3 D 3 only

14. June/2022/Paper_13/No.28

Which changes to the water potential and the volume of solution in the phloem sieve tube element occur when sucrose is moved from the phloem sieve tube element to an actively dividing shoot tip?

unloading of sucrose

	water potential in phloem sieve tube element becomes	volume of solution in phloem sieve tube element
A	higher	decreases
B	higher	increases
C	lower	decreases
D	lower	increases

sucrose moves down to sink cell by hydrostatic pressure, diffusion also moved by some water

water potential in sieve tube phloem, becomes higher

PapaCambridge

15. June/2022/Paper_13/No.29

Which processes occur during the loading of sucrose into phloem sieve tubes?

Loading of sucrose

1 Protons are pumped out of the cytoplasm of the companion cell into its cell wall.

2 There is a higher concentration of protons in the symplastic pathway outside the companion cell. apoplastic movement of H2O between Cell wall intercelular spaces through X

3 Protons are unable to move back into the companion cell.

4 A co-transporter molecule acts as a carrier for protons and sucrose. H+ and sucrose move back to cell through Co-transporter

A 1 and 2 B 1 and 4 C 2 and 3 D 3 and 4

PapaCambridge

16. June/2022/Paper_22/No.6

The transport tissues of plants are phloem and xylem.

The role of xylem is the transport of water and mineral ions from the soil solution to the different parts of the plant body.

The role of phloem is the translocation of assimilates and other substances from sources to sinks.

(a) The source of mineral ions for the plant is the soil solution. These mineral ions are transported from the roots in the xylem. Mineral ions are also found in the phloem sap within phloem sieve tubes.

Suggest why mineral ions are found within phloem sieve tubes **and** state how they are transported within phloem sieve tubes.

Mineral ions enter the phloem from xylem through plasmodesmata of the companion cells. The mineral ions are dissolved in water. So they flow with water entering the phloem. Materials move within the phloem by massflo

[3]

(b) Phloem tissue consists of different cell types.

Complete the passage using the most appropriate terms to summarise some of the features of phloem tissue.

The end walls of sieve tube elements are modified to allow efficient flow of phloem sap by the formation of Sieve plates . These structures also prevent the cells from bursting under pressure. The cytoplasm of sieve tube elements is very much reduced and is found at the edge of the cells. Most of the organelles in the cell are absent. Adjacent to sieve tube elements are Companion cells that carry out the metabolic processes of the missing organelles, allowing the sieve tube elements to function. [3]

[Total: 6]

PapaCambridge

17. June/2022/Paper_23/No.3

Hakea spp. are xerophytic plants native to Australia. The leaves of *Hakea* have adaptations for a xerophytic mode of life.

(a) Fig. 3.1 is a photomicrograph of a transverse section of part of a leaf of *H. laurina*.




Fig. 3.1

Describe the xerophytic features of the waxy cuticle and the stoma shown in Fig. 3.1 **and** explain how these features adapt the plant to a xerophytic mode of life.

The stoma is sunken to reduce rate of water through evaporation. It also lowers rate of transpiration.
The thick waxy cuticle is water proof. This reduces rate of water loss.

[3]

(b) When the availability of phosphate ions and other soil nutrients is limited, a number of changes occur in the roots of *Hakea* spp.:

- Regions of meristematic tissue are active for a few days.
- Root clusters are formed. A root cluster is a dense arrangement of tiny side roots known as rootlets.
- Most of the epidermal cells of the rootlets are root hair cells.
- Rootlets release compounds into the soil that make phosphates and other mineral ions more soluble for uptake.
- Uptake of phosphate ions and the absorption of water from the soil increases.

PapaCambridge

(i) Meristem cells have a similar role to stem cells in animals.

Suggest how meristematic tissue activity leads to the formation of root clusters.

Meristem tissue cells divide by mitosis. This produce many cells that are genetically identical. Increase in cells at the meristem leads to formation of rootlets. Many rootlets will amplify the rate of uptake of phosphate ions. [3]

(ii) Organic anions (negatively charged organic compounds) are released into the soil by rootlets. The concentration of these organic anions can become higher in the soil solution than in the rootlet cells.

Suggest **and** explain how the concentration of organic anions in the soil solution can become higher than in the rootlet cells.

The anions are pumped out of cells to the soil against their concentration gradient. They move out through the chanel proteins. Pumping of anions require energy. The enzyme ATPase stimulate breakdown of ATP to provide energy. The anions are moved by active transport [3]

(iii) Explain how the formation of root clusters can lead to an increase in the uptake of phosphate ions and absorption of water from the soil solution.

Root clusters increase surface area because they have more root hair cells. This increase carrier proteins for the uptake [2]

[Total: 11]

PapaCambridge