

Please check the examination details below before entering your candidate information

Candidate

surname:

USMAN

Centre Number

Candidate Number

Other names:

Grade A* Scholar

--	--	--	--	--	--

--	--	--	--	--	--

Pearson Edexcel International Advanced Level

Chemistry

International Advanced Level

UNIT 4: Rates, Equilibria and Further Organic Chemistry

Paper Reference: WCH14/01

Time Allowed: 2 hours 30 minutes

Topic Focus: 11A.1 Rate Measurement & 11A.2 Rate Equations

Total Marks: 150 Marks

INSTRUCTIONS

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Answer the questions in the spaces provided — there may be more space than you need.
- Show all the steps in any calculations and state the units of numerical values.
- Calculators may be used.

INFORMATION

- The total mark for this paper is 150.
- The marks for each question are shown in brackets — use this as a guide as to how much time to spend on each question.
- In questions marked with an asterisk (*), marks will be awarded for your ability to structure your answer logically, showing how the points that you make are related or follow on from each other.
- A Periodic Table is printed on the back page of this question paper.

ADVICE

- Read each question carefully before you start to answer it.
- Keep an eye on the time.
- Try to answer every question.
- Check your answers if you have time at the end.

MENTORA ACADEMY • WEEK 1 CURRICULUM MASTERY & A* SCHOLAR EXAM

Prepared exclusively for candidate USMAN · Strict timed simulation conditions

SECTION A

Answer ALL questions in this section.

You should spend no more than 40 minutes on this section.

Use the spaces provided in this question paper.

For each question, select one answer from A to D and put a cross in the box [X].

If you change your mind, put a line through the box and then mark your new answer with a cross.

1 A reaction has the rate equation: $\text{rate} = k[\text{A}]^{0.5} [\text{B}]^2$. What are the correct SI units for the rate constant k ?

- ☐ A $\text{mol}^{-1.5} \text{dm}^4.5 \text{s}^{-1}$
- ☐ B $\text{mol}^{-0.5} \text{dm}^{1.5} \text{s}^{-1}$
- ☐ C $\text{mol}^{-2} \text{dm}^6 \text{s}^{-1}$
- ☐ D $\text{dm}^3 \text{mol}^{-1} \text{s}^{-1}$

(Total for Question 1 = 1 mark)

2 For the gas-phase reaction $2\text{NO}(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{NO}_2(\text{g})$, the rate equation is $\text{rate} = k[\text{NO}]^2 [\text{O}_2]$. The volume of the closed reaction vessel is doubled at constant temperature. By what factor does the initial rate change?

- ☐ A Halves (0.5x)
- ☐ B Decreases to one-quarter (0.25x)
- ☐ C Decreases to one-eighth (0.125x)
- ☐ D Remains unchanged

(Total for Question 2 = 1 mark)

3 In the acid-catalysed reaction between propanone and iodine, the rate equation is $\text{rate} = k[\text{propanone}][\text{H}^+]$. Why does the concentration of iodine NOT appear in the rate equation?

- ☐ A Iodine acts as a heterogeneous catalyst.
- ☐ B Iodine is involved in a fast step after the rate-determining step.
- ☐ C Iodine is in zero-order large excess in all experiments.
- ☐ D The reaction between iodine and propanone is reversible.

(Total for Question 3 = 1 mark)

4 A student uses a colorimeter to follow the oxidation of ethanedioic acid by acidified potassium manganate(VII). Manganate(VII) ions are intensely purple, absorbing light strongly between 520 nm and 550 nm. Which colour filter should be chosen for maximum sensitivity?

- ☐ A Purple filter
- ☐ B Red filter
- ☐ C Green filter
- ☐ D Blue filter

(Total for Question 4 = 1 mark)

5 In a sampling and titrimetric study of the alkaline hydrolysis of ethyl ethanoate, $\text{CH}_3\text{COOCH}_2\text{CH}_3 + \text{OH}^- \rightarrow \text{CH}_3\text{COO}^- + \text{CH}_3\text{CH}_2\text{OH}$, aliquots are withdrawn at regular intervals. What is the most reliable method to quench the reaction immediately?

- ☐ A Add the aliquot to an excess of ice-cold standard hydrochloric acid.
- ☐ B Cool the aliquot rapidly in an ice-water bath without adding chemicals.
- ☐ C Add an excess of ethanol to shift the equilibrium position.
- ☐ D Filter the aliquot immediately to remove unreacted ester.

(Total for Question 5 = 1 mark)

6 A reaction produces 120 cm^3 of gas at r.t.p. On an electronic balance reading to $\pm 0.01 \text{ g}$, why is the 'loss-in-mass' method completely unsuited for reactions evolving hydrogen gas (H_2) compared to carbon dioxide (CO_2)?

- ☐ A Hydrogen gas dissolves completely in aqueous acid.
- ☐ B The mass of 120 cm^3 of H_2 is 0.010 g , giving an unacceptable apparatus percentage uncertainty ($\sim 100\%$).
- ☐ C Hydrogen gas reacts with atmospheric oxygen on escaping the flask.
- ☐ D Hydrogen gas is too dense to leave the cotton wool plug.

(Total for Question 6 = 1 mark)

7 The reaction $\text{CH}_3\text{COOCH}_2\text{CH}_3(\text{aq}) + \text{NaOH}(\text{aq}) \rightarrow \text{CH}_3\text{COONa}(\text{aq}) + \text{CH}_3\text{CH}_2\text{OH}(\text{aq})$ is monitored using a conductivity meter. Why does electrical conductivity decrease steadily as the reaction proceeds?

- ☐ A Covalent ethyl ethanoate is converted into ions.
- ☐ B Highly mobile hydroxide ions (OH^-) are replaced by less mobile ethanoate ions (CH_3COO^-).
- ☐ C Sodium ions are precipitated as insoluble sodium ethanoate.
- ☐ D The total concentration of charged ions drops to zero at the end.

(Total for Question 7 = 1 mark)

8 When the concentration of reactant X is increased by a factor of 4, the initial rate of reaction increases by a factor of 8. What is the order of reaction with respect to X?

- ☐ A 1
- ☐ B 1.5
- ☐ C 2
- ☐ D 3

(Total for Question 8 = 1 mark)

9 A reaction $\text{A} + 2\text{B} \rightarrow \text{C}$ follows the rate law $\text{rate} = k[\text{A}][\text{B}]^2$, with $k = 4.5 \times 10^{-3} \text{ dm}^3 \text{ mol}^{-2} \text{ s}^{-1}$. When $[\text{B}]$ is held in vast excess at 2.0 mol dm^{-3} , the reaction displays pseudo-first-order kinetics with $\text{rate} = k_{\text{obs}}[\text{A}]$. What is the numerical value of k_{obs} ?

- ☐ A $9.0 \times 10^{-3} \text{ s}^{-1}$
- ☐ B $1.8 \times 10^{-2} \text{ s}^{-1}$
- ☐ C $4.5 \times 10^{-3} \text{ s}^{-1}$
- ☐ D $3.6 \times 10^{-2} \text{ s}^{-1}$

(Total for Question 9 = 1 mark)

10 In the peroxydisulfate-iodide clock reaction, which assumption is essential for the relationship (initial rate proportional to $1/t$) to remain valid?

- ☐ A The reaction goes to 100% completion before the colour change occurs.
- ☐ B The amount of thiosulfate consumes a very small fraction (<10%) of the total reactants.
- ☐ C The starch indicator acts as an active catalyst.
- ☐ D The reaction is zero order with respect to all reactants.

(Total for Question 10 = 1 mark)

11 A tangent drawn to a concentration-time curve at $t = 0$ passes through (0 s, 0.40 mol dm^{-3}) and (80 s, 0.00 mol dm^{-3}). What is the initial rate of reaction?

- ☐ A $5.0 \times 10^{-3} \text{ mol dm}^{-3} \text{ s}^{-1}$
- ☐ B $2.0 \times 10^{-3} \text{ mol dm}^{-3} \text{ s}^{-1}$
- ☐ C $3.2 \times 10^{-2} \text{ mol dm}^{-3} \text{ s}^{-1}$
- ☐ D $0.40 \text{ mol dm}^{-3} \text{ s}^{-1}$

(Total for Question 11 = 1 mark)

12 In the iodine clock reaction $\text{S}_2\text{O}_8^{2-} + 2\text{I}^- \rightarrow 2\text{SO}_4^{2-} + \text{I}_2$, what is the exact chemical role of the added sodium thiosulfate ($\text{Na}_2\text{S}_2\text{O}_3$)?

- ☐ A It oxidises sulfate ions to regenerate persulfate.
- ☐ B It reacts instantaneously with I_2 to regenerate I^- until thiosulfate is completely consumed.
- ☐ C It complexes with starch to create the blue-black species.
- ☐ D It buffers the pH of the aqueous solution at pH 7.0.

(Total for Question 12 = 1 mark)

13 Which of the following experimental techniques allows continuous monitoring of reaction rate WITHOUT the need to withdraw and quench reaction samples?

- ☐ A Sampling followed by acid-base titration
- ☐ B Sampling followed by iodine titration
- ☐ C Continuous measurement of electrical conductivity
- ☐ D Precipitation and gravimetric weighing

(Total for Question 13 = 1 mark)

14 Dilatometry is an experimental technique that measures the rate of reaction by monitoring:

- ☐ A The small volume change of a liquid reaction mixture inside a precision capillary.
- ☐ B The change in refractive index of the solution.
- ☐ C The rate of heat dissipation to the surroundings.
- ☐ D The pressure of an evolved gas in a manometer.

(Total for Question 14 = 1 mark)

15 Consider the initial rate data: Exp 1: $[A]=0.1$, $[B]=0.1$, $\text{Rate}=2 \times 10^{-4}$; Exp 2: $[A]=0.2$, $[B]=0.1$, $\text{Rate}=4 \times 10^{-4}$; Exp 3: $[A]=0.3$, $[B]=0.2$, $\text{Rate}=2.4 \times 10^{-3}$. What is the overall order of the reaction?

- ☐ A 1
- ☐ B 2
- ☐ C 3
- ☐ D 4

(Total for Question 15 = 1 mark)

16 When the temperature of an aqueous reaction mixture is increased by 10 K at constant reactant concentrations, what happens to the rate constant k and the reaction orders?

- ☐ A Both k and the reaction orders double.
- ☐ B k increases substantially (approx. doubles); reaction orders remain unchanged.
- ☐ C k remains unchanged; reaction orders increase.
- ☐ D k increases; reaction orders decrease to zero.

(Total for Question 16 = 1 mark)

17 A plot of $\log_{10}(\text{initial rate})$ against $\log_{10}[A]$ yields a straight line with gradient = +2.0 and y-intercept = -1.30. What is the value of the rate constant k ?

- ☐ A $0.050 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$
- ☐ B $0.20 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$
- ☐ C $1.30 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$
- ☐ D $2.00 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$

(Total for Question 17 = 1 mark)

18 When calcium carbonate chips react with dilute hydrochloric acid in a conical flask connected to a gas syringe, why is there often a noticeable delay (lag phase) before gas volume begins to register?

- ☐ A Friction in the syringe plunger prevents immediate movement.
- ☐ B Carbon dioxide is moderately soluble in water and the solution must become saturated before gas is evolved.
- ☐ C The activation energy of the reaction increases initially.
- ☐ D The reaction is initially endothermic and cools the flask.

(Total for Question 18 = 1 mark)

19 The acid-catalysed hydrolysis of sucrose to glucose and fructose is called the 'inversion of sucrose'. Which optical instrument can be used to follow the reaction rate continuously?

- ☐ A Colorimeter
- ☐ B Spectrophotometer
- ☐ C Polarimeter
- ☐ D Refractometer

(Total for Question 19 = 1 mark)

20 In a titrimetric study of the decomposition of hydrogen peroxide, $2\text{H}_2\text{O}_2(\text{aq}) \rightarrow 2\text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g})$, aliquots are titrated with standard KMnO_4 . Why is quenching in ice-cold dilute H_2SO_4 highly effective?

- ☐ A Sulfuric acid acts as an inhibitor and destroys the MnO_2 catalyst, while low temperature slows the reaction.
- ☐ B Sulfuric acid precipitates the hydrogen peroxide as an insoluble sulfate.
- ☐ C Sulfuric acid oxidises water to stop the backward reaction.
- ☐ D Sulfuric acid acts as an indicator in the titration.

(Total for Question 20 = 1 mark)

21 Marble chips (CaCO_3) react with excess 1.0 mol dm^{-3} HCl . What is the order of reaction with respect to solid calcium carbonate?

- ☐ A Zero order
- ☐ B First order
- ☐ C Second order
- ☐ D Half order

(Total for Question 21 = 1 mark)

22 For a reaction rate = $k[P]^2 [Q]$, where rate is measured in $\text{mol dm}^{-3} \text{ min}^{-1}$ and concentrations in mol dm^{-3} , what are the units of k ?

- ☐ A $\text{dm}^6 \text{ mol}^{-2} \text{ min}^{-1}$
- ☐ B $\text{dm}^3 \text{ mol}^{-1} \text{ min}^{-1}$
- ☐ C $\text{mol}^{-2} \text{ dm}^6 \text{ s}^{-1}$
- ☐ D min^{-1}

(Total for Question 22 = 1 mark)

23 For a zero-order reaction $A \rightarrow \text{products}$, which of the following plots gives a straight line with a negative gradient?

- ☐ A $[A]$ against time t
- ☐ B $\ln[A]$ against time t
- ☐ C $1/[A]$ against time t
- ☐ D rate against $[A]$

(Total for Question 23 = 1 mark)

24 For a second-order reaction with respect to reactant A, which plot yields a straight line with a POSITIVE gradient equal to k ?

- ☐ A $[A]$ against time t
- ☐ B $\ln[A]$ against time t
- ☐ C $1/[A]$ against time t
- ☐ D $[A]^2$ against time t

(Total for Question 24 = 1 mark)

25 Why must a colorimeter be calibrated with a cuvette containing only pure solvent (a 'blank') before recording absorbance data?

- ☐ A To set the absorbance to 1.00 before the reaction begins.
- ☐ B To compensate for light absorption, reflection, and scattering by the solvent and cuvette walls.
- ☐ C To heat the cuvette to the reaction temperature.
- ☐ D To ensure the detector operates at maximum wavelength.

(Total for Question 25 = 1 mark)

26 In a clock reaction investigation using varying volumes of potassium iodide, why MUST deionised water be added to keep the total volume constant across all runs?

- ☐ A To maintain constant ionic strength and ensure volume of reactant is directly proportional to concentration.
- ☐ B To prevent the solution from overheating during the reaction.
- ☐ C To dilute the starch indicator so it does not coagulate.
- ☐ D To keep the pH strictly buffered at pH 4.0.

(Total for Question 26 = 1 mark)

27 For the stoichiometric reaction $2A + B \rightarrow 3C$, the rate of consumption of A is $-d[A]/dt = 0.048 \text{ mol dm}^{-3} \text{ s}^{-1}$. What is the rate of formation of C ($d[C]/dt$)?

- ☐ A $0.024 \text{ mol dm}^{-3} \text{ s}^{-1}$
- ☐ B $0.032 \text{ mol dm}^{-3} \text{ s}^{-1}$
- ☐ C $0.072 \text{ mol dm}^{-3} \text{ s}^{-1}$
- ☐ D $0.048 \text{ mol dm}^{-3} \text{ s}^{-1}$

(Total for Question 27 = 1 mark)

28 The nucleophilic substitution $\text{CH}_3\text{Br} + \text{OH}^- \rightarrow \text{CH}_3\text{OH} + \text{Br}^-$ has the rate law $\text{rate} = k[\text{CH}_3\text{Br}][\text{OH}^-]$. If $[\text{CH}_3\text{Br}]$ is tripled and $[\text{OH}^-]$ is divided by 3, what happens to the rate?

- ☐ A Increases by a factor of 9
- ☐ B Decreases by a factor of 9
- ☐ C Remains exactly the same
- ☐ D Increases by a factor of 3

(Total for Question 28 = 1 mark)

29 Why is the half-life of a first-order reaction completely independent of the starting concentration?

- ☐ A Because rate is inversely proportional to concentration.
- ☐ B Because rate is directly proportional to concentration, so as concentration doubles, the reaction proceeds twice as fast.
- ☐ C Because the rate constant k decreases as concentration falls.
- ☐ D Because the activation energy decreases as the reaction proceeds.

(Total for Question 29 = 1 mark)

30 A student carries out a continuous colorimetric rate experiment with iodine in a colorimeter without temperature control. Over 15 minutes, the cell temperature rises by 4.2°C due to lamp heating. What systematic error does this introduce?


- ☐ A The calculated rate constant k will be progressively overestimated as the run proceeds.
- ☐ B The order with respect to iodine will change from zero to second order.
- ☐ C The absorbance reading will become negative.
- ☐ D The rate of reaction will decrease towards the end of the run.

(Total for Question 30 = 1 mark)

SECTION B

Answer ALL questions. Write your answers in the spaces provided.

31 Propanone reacts with iodine in aqueous acid according to the equation:



A student investigates the kinetics of this reaction at 298 K using continuous colorimetry.

(a) State why colorimetry is an ideal experimental technique for continuously monitoring the rate of this reaction. Identify the species responsible for the colour change and suggest a suitable colour for the filter used in the colorimeter.

(3)

Species responsible for colour:

Colour of filter:

(b) The student performs four initial rate experiments at 298 K. The results are shown in the table below:

Experiment	$[\text{CH}_3\text{COCH}_3] / \text{mol dm}^{-3}$	$[\text{I}_2] / \text{mol dm}^{-3}$	$[\text{H}^+] / \text{mol dm}^{-3}$	Initial Rate / $\text{mol dm}^{-3} \text{ s}^{-1}$
1	0.400	0.00200	0.400	1.40×10^{-6}
2	0.800	0.00200	0.400	2.80×10^{-6}
3	0.400	0.00400	0.400	1.40×10^{-6}
4	0.400	0.00200	0.800	2.80×10^{-6}

Deduce the order of reaction with respect to propanone, iodine, and hydrogen ions. Fully justify your deduction for each reactant by comparing appropriate experiments.

(6)

Order with respect to propanone =
Order with respect to iodine =
Order with respect to H^+ =

(c) (i) Write the overall rate equation for the reaction. (1)

Rate equation =

(ii) Calculate the rate constant, k , at 298 K using data from Experiment 1, and state its units. (2)

k = units

(d) (i) On the axes below, sketch the graph of $[I_2]$ against time when propanone and H^+ are present in large excess. (2)




(ii) Explain how the feature of your sketched graph confirms the order with respect to iodine. (1)


(e) Explain why $[H^+]$ appears in the rate equation even though hydrogen ions are regenerated in the stoichiometric equation and do not appear as a net reactant in the overall reaction. (3)

(Total for Question 31 = 18 marks)

32 Persulfate ions oxidise iodide ions in aqueous solution according to Reaction 1:



In a clock experiment, a small fixed amount of sodium thiosulfate ($\text{Na}_2\text{S}_2\text{O}_3$) and starch indicator are added:



(a) Explain the mechanism of the colour change in this clock reaction. Why does the mixture remain completely colourless initially, and what causes the sudden appearance of a dark blue-black colour at time t ?

(4)

(b) A series of experiments is conducted at 20 °C. In each run, exactly 5.0 cm³ of 0.010 mol dm⁻³ $\text{Na}_2\text{S}_2\text{O}_3$ and starch are added, and deionised water is used to keep the total volume at exactly 50.0 cm³.

Run	Vol 0.050 M $\text{S}_2\text{O}_8^{2-} / \text{cm}^3$	Vol 0.100 M I^- / cm^3	Vol Water / cm ³	Time t / s
1	10.0	10.0	25.0	88.0
2	20.0	10.0	15.0	44.0
3	10.0	20.0	15.0	44.0
4	15.0	20.0	10.0	29.3

(i) Deduce the order of reaction with respect to $\text{S}_2\text{O}_8^{2-}$ and I^- . Show your reasoning clearly.

(2)

Order with respect to $\text{S}_2\text{O}_8^{2-}$ =

Order with respect to I^- =

(ii) Calculate the concentration of $\text{S}_2\text{O}_8^{2-}$ and I^- present in the reaction vessel in Run 1 before the reaction begins.

(2)

$[\text{S}_2\text{O}_8^{2-}] = \dots\dots\dots \text{mol dm}^{-3}$ $[\text{I}^-] = \dots\dots\dots \text{mol dm}^{-3}$

(iii) In Run 1, 5.0×10^{-5} mol of $\text{S}_2\text{O}_3^{2-}$ was consumed before the colour change. Calculate the initial rate of reaction in Run 1 in $\text{mol dm}^{-3} \text{s}^{-1}$ with respect to $\text{S}_2\text{O}_8^{2-}$.

(2)

Initial rate = $\dots\dots\dots \text{mol dm}^{-3} \text{s}^{-1}$

(iv) Calculate the rate constant, k , at 20 °C using your answers to (b)(ii) and (b)(iii). Include units in your answer.

(2)

$k = \dots\dots\dots \text{units} \dots\dots\dots$

(c) Explain why the amount of sodium thiosulfate added must be significantly less than the amounts of persulfate and iodide present in the mixture.

(3)

(d) State two distinct reasons why deionised water was added to maintain a constant total volume of 50.0 cm^3 across all runs.

(3)

(Total for Question 32 = 18 marks)

33 Ethyl ethanoate is hydrolysed by sodium hydroxide according to the second-order reaction:



Rate = k [ester] [OH⁻]. A student studies this reaction using both conductimetry and titrimetry.

- (a) Explain in terms of ionic mobilities why the electrical conductivity of the reaction mixture decreases as this reaction progresses, and state one advantage of conductimetry over titrimetry. **(3)**

- (b) In a sampling study, a student withdraws 10.0 cm³ aliquots every 2 minutes. Describe the precise procedure to quench each aliquot, and explain why pouring into ice-cold water alone is inadequate compared to quenching in excess standard acid followed by back-titration. **(4)**

- (c) (i) In Experiment 1, the student uses [ester]₀ = 0.0150 mol dm⁻³ and [OH⁻]₀ = 0.600 mol dm⁻³. Explain why the reaction obeys pseudo-first-order kinetics under these conditions. **(2)**

(ii) Under the conditions of Experiment 1, a plot of $\ln[\text{ester}]$ against time t yields a straight line with gradient $= -0.0660 \text{ s}^{-1}$. Calculate the apparent pseudo-rate constant k' and the true second-order rate constant k , stating the units of k .

(5)

$k' = \dots\dots\dots \text{s}^{-1}$

$k = \dots\dots\dots \text{units} \dots\dots\dots$


(d) Calculate the half-life ($t_{1/2}$) of ethyl ethanoate in Experiment 1. State and explain what happens to the half-life if the initial concentration of sodium hydroxide is doubled to 1.20 mol dm^{-3} .

(4)

$t_{1/2} = \dots\dots\dots \text{s}$

(Total for Question 33 = 18 marks)

34 Aqueous hydrogen peroxide decomposes catalytically according to the equation:



*(a) A teacher presents two laboratory methods to follow the rate of this reaction:

Method 1: Collecting oxygen gas in a 100 cm³ gas syringe.

Method 2: Monitoring the loss in mass on an electronic balance reading to 0.01 g.

Compare and evaluate both methods. In your response, discuss: gas solubility, temperature and friction errors, apparatus percentage uncertainties, and aerosol spray loss.

(6)

(b) A student collects oxygen in a gas syringe using 25.0 cm³ of H₂O₂ at 298 K. The total volume at completion (V_{inf}) was 68.0 cm³.

Time t / s	0	30	60	90	120	150	180	240	300	inf
$V(\text{O}_2) / \text{cm}^3$	0.0	18.0	31.5	41.5	49.0	54.5	58.5	63.5	66.0	68.0
$(V_{\text{inf}} - V) / \text{cm}^3$	68.0	50.0	36.5	26.5	19.0	13.5	9.5	4.5	2.0	0.0

Show that the reaction is first-order with respect to H₂O₂ by determining two successive half-lives from the data above. Clearly state the values of both half-lives.

(6)

First half-life, $t_{1/2}(1) = \dots\dots\dots$ s

Second half-life, $t_{1/2}(2) = \dots\dots\dots$ s

(c) A tangent drawn to the $V(\text{O}_2)$ vs time curve at $t = 0$ passes through (0 s, 0 cm³) and (50 s, 32.0 cm³). Calculate the initial rate in cm³ s⁻¹, and convert this into mol dm⁻³ s⁻¹ of H₂O₂ consumed in the 25.0 cm³ solution ($V_m = 24,000 \text{ cm}^3 \text{ mol}^{-1}$).

(3)

Initial rate = $\dots\dots\dots$ mol dm⁻³ s⁻¹

(d) Calculate the rate constant, k , for this decomposition from your average half-life using $t_{1/2} = \ln 2 / k$. Include units.

(3)

$k = \dots\dots\dots$ units $\dots\dots\dots$

(Total for Question 34 = 18 marks)

35 Nitrogen monoxide reacts with hydrogen at 1000 K according to the equation:



(a) Explain how monitoring the total pressure of the reaction vessel at constant volume allows the reaction rate to be determined, and state why the reaction temperature must be kept strictly above 100 °C.

(3)

(b) The following initial rate data were obtained at 1000 K:

Experiment	$[\text{NO}] / 10^{-3} \text{ mol dm}^{-3}$	$[\text{H}_2] / 10^{-3} \text{ mol dm}^{-3}$	Initial Rate / $10^{-5} \text{ mol dm}^{-3} \text{ s}^{-1}$
1	1.50	2.00	3.60
2	3.00	2.00	14.40
3	1.50	6.00	10.80
4	4.50	4.00	R_4

(i) Deduce the orders with respect to NO and H_2 with full reasoning.

(4)

Order with respect to NO =

Order with respect to H_2 =

(ii) Write the rate equation and calculate the rate constant, k , at 1000 K with units.

(3)

Rate equation =

k = units

(iii) Calculate the predicted initial rate, R_4 , in Experiment 4.

(2)

R_4 = $\text{mol dm}^{-3} \text{s}^{-1}$

(c) The reaction mixture is placed in a cylinder with a movable piston. The volume of the cylinder is compressed to one-third of its original volume at constant temperature.

(i) State what happens to the concentration of each gas.

(ii) Calculate the factor by which the initial rate will increase.

(iii) State and explain whether the rate constant k changes.

(6)

Factor increase in rate =

(Total for Question 35 = 18 marks)

SECTION C

Answer ALL questions. Write your answers in the spaces provided.

36 Crystal violet (CV^+) is an intensely violet triphenylmethane dye. It reacts with hydroxide ions to form a colourless carbinol base (CV-OH):



Rate = $k [\text{CV}^+]^m [\text{OH}^-]^n$. A student investigates the kinetics spectrophotometrically.

(a) Describe how a student prepares a calibration curve of absorbance vs $[\text{CV}^+]$ using a stock solution of $1.00 \times 10^{-4} \text{ mol dm}^{-3}$ crystal violet, a 590 nm filter, and serial dilution with standard volumetric glassware.

(4)

(b) In the kinetic run, $[\text{CV}^+]_0 = 2.00 \times 10^{-5} \text{ mol dm}^{-3}$ and $[\text{OH}^-]_0 = 0.100 \text{ mol dm}^{-3}$. Explain why this 5000-fold excess of OH^- allows the isolation method to be used.

(3)

(c) Prove mathematically that for a first-order reaction obeying Beer-Lambert's Law ($A = \epsilon c l$), a plot of $\ln(\text{Absorbance})$ against time gives a straight line of gradient $= -k'$. If the experimental gradient is -0.0185 s^{-1} , state the order m and calculate k' .

(5)

Order with respect to crystal violet, $m = \dots\dots\dots$

$k' = \dots\dots\dots \text{s}^{-1}$

(d) When $[\text{OH}^-]$ is increased from $0.100 \text{ mol dm}^{-3}$ to $0.200 \text{ mol dm}^{-3}$, the gradient becomes -0.0370 s^{-1} . Deduce the order n with respect to OH^- , write the overall rate equation, and calculate the true rate constant k .

(3)


Order with respect to OH^- , $n = \dots\dots\dots$

Rate equation = $\dots\dots\dots$

$k = \dots\dots\dots \text{units} \dots\dots\dots$

(Total for Question 36 = 15 marks)

37 Sucrose hydrolyses in aqueous acid to produce glucose and fructose:



Sucrose is dextrorotatory (+66.5°), but the equimolar product mixture is laevorotatory (–19.7°).

(a) Explain how a polarimeter works to continuously follow this reaction, and why the process is referred to historically as the 'inversion of sucrose'.

(4)

(b) The angle of rotation α_t was recorded over time at 298 K. The concentration of unreacted sucrose is directly proportional to $(\alpha_t - \alpha_{\text{inf}})$:

Time t / min	0	20	40	60	80	120
	32.0	22.6	16.0	11.3	8.0	4.0

Prove that the reaction is first-order with respect to sucrose by determining two successive half-lives from this data. Calculate the rate constant k in min^{-1} .

(4)

First half-life = min
 Second half-life = min
 k = min^{-1}

(c) Explain why a dilatometer can follow this reaction even though no gas is evolved and no colour changes occur.

(3)

(d) Explain why measuring electrical conductivity is completely unsuitable for following the rate of this reaction.

(2)

(e) Compare the initial rate of inversion when $1.0 \text{ mol dm}^{-3} \text{ HCl}$ is used as catalyst compared to $1.0 \text{ mol dm}^{-3} \text{ CH}_3\text{COOH}$. Explain your answer with reference to $[\text{H}^+]$.

(2)

(Total for Question 37 = 15 marks)

TOTAL FOR SECTION C = 30 MARKS

TOTAL FOR PAPER = 150 MARKS

**END OF QUESTION
PAPER**

DATA SHEET – THE PERIODIC TABLE OF THE ELEMENTS

Group 1	Group 2	Group 3	Group 4	Group 5	Group 6	Group 7	Group 0
1 H 1.0	-	-	-	-	-	-	2 He 4.0
3 Li 6.9	4 Be 9.0	5 B 10.8	6 C 12.0	7 N 14.0	8 O 16.0	9 F 19.0	10 Ne 20.2
11 Na 23.0	12 Mg 24.3	13 Al 27.0	14 Si 28.1	15 P 31.0	16 S 32.1	17 Cl 35.5	18 Ar 39.9
19 K 39.1	20 Ca 40.1	31 Ga 69.7	32 Ge 72.6	33 As 74.9	34 Se 79.0	35 Br 79.9	36 Kr 83.8
37 Rb 85.5	38 Sr 87.6	49 In 114.8	50 Sn 118.7	51 Sb 121.8	52 Te 127.6	53 I 126.9	54 Xe 131.3
55 Cs 132.9	56 Ba 137.3	81 Tl 204.4	82 Pb 207.2	83 Bi 209.0	84 Po [209]	85 At [210]	86 Rn [222]

Physical Constant / Formula	Symbol / Expression	Value & Units
Gas constant	R	8.314 J mol ⁻¹ K ⁻¹
Molar volume of gas at r.t.p.	V _m	24.0 dm ³ mol ⁻¹ (24,000 cm ³ mol ⁻¹)
First-order half-life relationship	t _{1/2}	t _{1/2} = ln 2 / k = 0.693 / k
Beer-Lambert Law	A	A = l c

PEARSON EDEXCEL INTERNATIONAL ADVANCED LEVEL

CHEMISTRY UNIT 4 (WCH14/01) – ADVANCED MASTERY QUIZ

CONFIDENTIAL TEACHER'S MARK SCHEME & EXAMINER GUIDANCE (150 MARKS)

SECTION A ANSWER KEY & EXPLANATIONS (30 MARKS)

Q1	Key: A	Overall order = $0.5 + 2 = 2.5$. Units of $k = (\text{mol dm}^{-3} \text{ s}^{-1}) / (\text{mol dm}^{-3})^{2.5} = \text{mol}^{-1.5} \text{ dm}^{4.5} \text{ s}^{-1}$. (1)
Q2	Key: C	Doubling volume halves all concentrations: rate = $k(0.5[\text{NO}])^2(0.5[\text{O}_2]) = (0.5)^3 = 0.125x$ (decreases by factor of 8). (1)
Q3	Key: B	Iodine is involved in a fast step AFTER the rate-determining step, so its concentration does not affect the rate. (1)
Q4	Key: C	Manganate(VII) absorbs strongly in the green region (520–550 nm); complementary green filter provides maximum absorbance sensitivity. (1)
Q5	Key: A	Adding excess standard ice-cold acid removes OH^- instantaneously (quenching) and cools mixture; unreacted acid is back-titrated. (1)
Q6	Key: B	Mass of 120 cm^3 of H_2 is only 0.010 g. On a $\pm 0.01 \text{ g}$ balance, % uncertainty is 100%, destroying reliability. (1)
Q7	Key: B	Highly mobile OH^- ions (ionic conductivity $198 \text{ S cm}^2 \text{ mol}^{-1}$) are replaced by bulky, slower CH_3COO^- ions ($41 \text{ S cm}^2 \text{ mol}^{-1}$). (1)
Q8	Key: B	$4^n = 8 \Rightarrow (2^2)^n = 2^3 \Rightarrow 2n = 3 \Rightarrow n = 1.5$. (1)
Q9	Key: B	$k_{\text{obs}} = k[\text{B}]^2 = (4.5 \times 10^{-3})(2.0)^2 = 1.8 \times 10^{-2} \text{ s}^{-1}$. (1)
Q10	Key: B	Initial rate approx $1/t$ requires $\Delta[\text{reactant}]$ to be negligible ($<10\%$) so concentrations remain constant during time t . (1)
Q11	Key: A	Gradient magnitude = $0.40 / 80 = 5.0 \times 10^{-3} \text{ mol dm}^{-3} \text{ s}^{-1}$. (1)
Q12	Key: B	Thiosulfate rapidly reduces I_2 back to I^- as fast as it forms until all $\text{S}_2\text{O}_3^{2-}$ is exhausted, then free I_2 forms blue-black starch complex. (1)
Q13	Key: C	Electrical conductivity probe stays in the reaction vessel and measures continuously without removing or quenching any liquid. (1)
Q14	Key: A	Dilatometry measures small liquid volume changes in a narrow capillary due to bond formation/breakage and hydration changes. (1)
Q15	Key: C	From Exp 1 & 2: doubling $[\text{A}]$ doubles rate $\Rightarrow$ order in A = 1. Exp 1 & 3: $[\text{A}]$ triples ($3x$) and rate increases $12x \Rightarrow (3)^1 \times (2)^n = 12 \Rightarrow 2^n = 4 \Rightarrow n = 2$. Overall order = $1 + 2 = 3$. (1)
Q16	Key: B	k increases exponentially with temperature due to more molecules having $E > E_a$; reaction orders depend strictly on mechanism and remain unchanged. (1)
Q17	Key: A	$\log_{10}(\text{rate}) = \log_{10}(k) + 2 \log_{10}[\text{A}]$. Intercept = $\log_{10}(k) = -1.30 \Rightarrow k = 10^{-1.30} = 0.050 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$. (1)
Q18	Key: B	CO_2 is moderately soluble in water; bubbles only evolve once the liquid becomes saturated with dissolved CO_2 . (1)
Q19	Key: C	Polarimeter measures rotation of plane-polarised light as sucrose ($+66.5^\circ$) hydrolyses to invert sugar (-19.7°). (1)
Q20	Key: A	Cold acid stops the catalytic activity of MnO_2 and lowers temperature drastically so decomposition freezes before titration. (1)
Q21	Key: A	Pure solid reactants have constant chemical activity/concentration; reaction rate depends only on exposed surface area, which is zero order. (1)
Q22	Key: A	Units of $k = (\text{mol dm}^{-3} \text{ min}^{-1}) / (\text{mol dm}^{-3})^3 = \text{dm}^6 \text{ mol}^{-2} \text{ min}^{-1}$. (1)

Q23	Key: A	For zero order, rate = $-d[A]/dt = k \Rightarrow [A]_t = -k t + [A]_0$ (linear negative slope). (1)
Q24	Key: C	Integrated second order rate law: $1/[A]_t = kt + 1/[A]_0$ (linear positive slope). (1)
Q25	Key: B	Blank calibration eliminates absorbance, reflection, and scatter from cuvette walls and solvent. (1)
Q26	Key: A	Total volume must be constant so that volume of reactant solution is directly proportional to its concentration. (1)
Q27	Key: C	Rate of formation of C = $(3/2) \times$ rate of consumption of A = $1.5 \times 0.048 = 0.072 \text{ mol dm}^{-3} \text{ s}^{-1}$. (1)
Q28	Key: C	New rate proportional to $(3) \times (1/3) = 1$. Rate remains completely unchanged. (1)
Q29	Key: B	Rate is proportional to $[A]$; doubling $[A]$ doubles rate, so the time taken to halve any concentration is constant ($t_{1/2} = \ln 2 / k$). (1)
Q30	Key: A	Reaction accelerates as temperature rises, so measured rates and apparent k will increase over time. (1)

QUESTION 31: Acid-Catalysed Iodination of Propanone (18 Marks)

(a) Suitability of Colorimetry	Brown/yellow iodine is consumed so absorbance decreases continuously without opening reaction vessel (1). Species responsible: Iodine (I_2) or triiodide (I_3^-) (1). Filter: 400–470 nm / blue filter (complementary colour to yellow/brown) (1).	[3]
(b) Order Deductions	Compare Exp 1 & 2: $[I_2]$ and $[H^+]$ constant; [propanone] doubles ($0.400 \rightarrow 0.800$), rate doubles ($1.40 \times 10^{-6} \rightarrow 2.80 \times 10^{-6}$) $\Rightarrow$ Order with respect to propanone = 1 (2). Compare Exp 1 & 3: [propanone] and $[H^+]$ constant; $[I_2]$ doubles ($0.00200 \rightarrow 0.00400$), rate unchanged (1.40×10^{-6}) $\Rightarrow$ Order with respect to $I_2 = 0$ (2). Compare Exp 1 & 4: [propanone] and $[I_2]$ constant; $[H^+]$ doubles ($0.400 \rightarrow 0.800$), rate doubles ($1.40 \times 10^{-6} \rightarrow 2.80 \times 10^{-6}$) $\Rightarrow$ Order with respect to $H^+ = 1$ (2).	[6]
(c) Rate Equation & k	(i) Rate = $k[CH_3COCH_3][H^+]$ (1). (ii) $k = \text{Rate} / ([CH_3COCH_3][H^+]) = 1.40 \times 10^{-6} / (0.400 \times 0.400) = 8.75 \times 10^{-6}$ (1). Units: $\text{mol dm}^{-3} \text{ s}^{-1} / (\text{mol dm}^{-3})^2 = \text{dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ (1).	[3]
(d) $[I_2]$ vs Time Graph	(i) Straight line with constant negative gradient starting from $[I_2]_0$ down to zero on the time axis (2). (ii) Constant gradient ($-d[I_2]/dt = \text{constant}$) demonstrates that the rate is independent of $[I_2]$, confirming zero order (1).	[3]
(e) Role of H^+ Catalyst	H^+ acts as a homogeneous catalyst (1). Participates in the slow rate-determining step (protonation of carbonyl oxygen / enolisation) (1). Regenerated in a subsequent rapid step when the enol intermediate attacks iodine, so it does not appear as a net reactant in the overall equation (1).	[3]

QUESTION 32: Peroxodisulfate–Iodide Clock Reaction (18 Marks)

(a) Mechanism of Clock	Reaction 1 produces I_2 slowly. Reaction 2 consumes I_2 instantaneously by reacting with $S_2O_3^{2-}$ to form I^- and $S_4O_6^{2-}$ (1). As long as $S_2O_3^{2-}$ remains, no free I_2 can accumulate, so solution remains completely colourless (1). The instant all $S_2O_3^{2-}$ is completely exhausted, the very next trace of I_2 formed reacts with starch indicator (1) to form a dark blue-black starch-triiodide complex at time t (1).	[4]
-------------------------------	--	------------

(b) Data Analysis & k	(i) Compare Runs 1 & 2: $[I^-]$ constant; $[S_2O_8^{2-}]$ doubles, time t halves ($88 \rightarrow 44$ s) so rate doubles $\Rightarrow$ Order in $S_2O_8^{2-} = 1$ (1). Compare Runs 1 & 3: $[S_2O_8^{2-}]$ constant; $[I^-]$ doubles, time halves $\Rightarrow$ Order in $I^- = 1$ (1). (ii) In Run 1: $[S_2O_8^{2-}] = (10.0 \times 0.050) / 50.0 = 0.010 \text{ mol dm}^{-3}$ (1); $[I^-] = (10.0 \times 0.100) / 50.0 = 0.020 \text{ mol dm}^{-3}$ (1). (iii) Moles $S_2O_8^{2-}$ reacted = $0.5 \times$ moles $S_2O_3^{2-} = 2.5 \times 10^{-5} \text{ mol}$. $\Delta[S_2O_8^{2-}] = 2.5 \times 10^{-5} / 0.050 \text{ dm}^3 = 5.0 \times 10^{-4} \text{ mol dm}^{-3}$. Initial rate = $5.0 \times 10^{-4} / 88.0 \text{ s} = 5.68 \times 10^{-6} \text{ mol dm}^{-3} \text{ s}^{-1}$ (2). (iv) $k = \text{Rate} / ([S_2O_8^{2-}][I^-]) = 5.68 \times 10^{-6} / (0.010 \times 0.020) = 0.0284$ (or $2.84 \times 10^{-2} \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$) (2).	[8]
(c) Thiosulfate Amount	Thiosulfate must be in small limiting amount so only a tiny fraction (<10%) of persulfate and iodide react before the colour change (1). This ensures concentrations of $S_2O_8^{2-}$ and I^- remain practically constant, so average rate measured equals the true initial rate (2).	[3]
(d) Constant Total Volume	(1) Ensures the volume of each reactant solution is directly proportional to its concentration (1). (2) Keeps total ionic strength and depth of solution constant across all runs (2).	[3]

QUESTION 33: Alkaline Hydrolysis of an Ester (18 Marks)

(a) Conductimetry	OH^- ions have very high molar conductivity (mobility), whereas CH_3COO^- ions formed have much lower mobility (1). As OH^- is consumed and replaced by CH_3COO^-, total conductance decreases linearly with conversion (1). Conductivity probe is non-invasive and provides continuous real-time data without removing samples (1).	[3]
(b) Quenching Procedure	Pipette 10.0 cm^3 aliquot into a known excess of ice-cold standard dilute HCl (1). Acid neutralises OH^- instantaneously, stopping the forward reaction immediately (1). Ice-water alone only slows the reaction by lowering temperature, but hydrolysis continues during titration, introducing systematic error (1). Unreacted HCl is back-titrated with standard NaOH using phenolphthalein indicator (1).	[4]
(c) Pseudo-First-Order & k	(i) $[OH^-]$ is in 40-fold excess (0.600 vs $0.0150 \text{ mol dm}^{-3}$); change in $[OH^-]$ is negligible (<2.5%), so $[OH^-]$ is virtually constant (2). Rate = $k[\text{ester}][OH^-] = k'[\text{ester}]$ where $k' = k[OH^-]$ (1). (ii) Gradient of $\ln[\text{ester}]$ vs $t = -k' \Rightarrow k' = 0.0660 \text{ s}^{-1}$ (2). $k = k' / [OH^-] = 0.0660 / 0.600 = 0.110 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ (2).	[7]
(d) Half-Life Evaluation	$t_{1/2} = \ln 2 / k' = 0.693 / 0.0660 = 10.5 \text{ s}$ (2). If $[OH^-]$ doubles to 1.20 mol dm^{-3}, k' doubles ($k' = k[OH^-]$) (1). Since $t_{1/2} = \ln 2 / k'$, the half-life will halve to 5.25 s (1).	[4]

QUESTION 34: Decomposition of H_2O_2 & Practical Evaluation (18 Marks)

*(a) Level of Response Evaluation	Level 3 (5–6 marks): Comprehensive comparative evaluation addressing both methods across all 4 key descriptors: (1) Gas syringe: Direct, non-invasive volume reading; errors include friction in plunger causing sticking/pressure build-up, and gas volume sensitivity to temperature fluctuations ($PV=nRT$). (2) Loss-in-mass: Low sensitivity because O_2 has low molar mass (32 g/mol); for 100 cm³ O_2, mass lost is only 0.133 g, which on a 0.01 g balance has ~7.5% apparatus uncertainty. (3) Aerosol/spray loss: Effervescence ejects liquid droplets causing overestimated mass loss unless a loose cotton wool plug is used. (4) Solubility: O_2 has slight aqueous solubility in both methods, causing minor lag at start. Level 2 (3–4 marks): Discusses both methods with 2–3 valid points. Level 1 (1–2 marks): Superficial comparison with single valid point.	[6]
(b) Half-Life Verification	$V_{\infty} = 68.0 \text{ cm}^3$. ($V_{\infty} - V$) represents unreacted $[H_2O_2]$ (1). At $t = 0$: ($V_{\infty} - V$) = 68.0 cm³. First half-life: value drops to 34.0 cm³. Occurs at $t = 68 \text{ s}$ ($t_{1/2}(1) = 68 \text{ s}$) (2). Second half-life: value drops to 17.0 cm³. Occurs at $t = 136 \text{ s}$ ($t_{1/2}(2) = 136 - 68 = 68 \text{ s}$) (2). Successive half-lives are constant (68 s and 68 s) within experimental error, confirming first-order kinetics with respect to H_2O_2 (1).	[6]
(c) Initial Rate Calculation	Initial rate in gas volume = $32.0 \text{ cm}^3 / 50 \text{ s} = 0.640 \text{ cm}^3 \text{ s}^{-1}$ (1). Moles of O_2 per second = $0.640 / 24,000 = 2.67 \times 10^{-5} \text{ mol s}^{-1}$ (1). Moles of H_2O_2 consumed = $2 \times 2.67 \times 10^{-5} = 5.33 \times 10^{-5} \text{ mol s}^{-1}$. Rate of H_2O_2 consumption = $(5.33 \times 10^{-5}) / 0.0250 \text{ dm}^3 = 2.13 \times 10^{-3} \text{ mol dm}^{-3} \text{ s}^{-1}$ (1).	[3]
(d) Rate Constant k	$k = \ln 2 / t_{1/2} = 0.693 / 68 \text{ s} = 0.0102 \text{ s}^{-1}$ (or $1.02 \times 10^{-2} \text{ s}^{-1}$) (2). Units: s^{-1} (1).	[3]

QUESTION 35: Gas-Phase Kinetics of NO and H₂ (18 Marks)

(a) Pressure Monitoring	4 moles of gaseous reactants ($2NO + 2H_2$) produce 3 moles of gaseous products ($N_2 + 2H_2O$ at $T > 100^\circ\text{C}$) (1). Total moles of gas decrease, so total pressure drops at constant volume and temperature, directly proportional to reaction extent (1). Temperature must exceed 100°C to prevent water vapour from condensing to liquid water, which would cause an unpredictable large pressure drop (1).	[3]
(b) Rate Law, k and R₄	(i) Compare Exp 1 & 2: $[H_2]$ constant; $[NO]$ doubles, rate quadruples ($3.60 \rightarrow 14.40 \times 10^{-5}$) $\Rightarrow$ Order with respect to $NO = 2$ (2). Compare Exp 1 & 3: $[NO]$ constant; $[H_2]$ triples ($2.00 \rightarrow 6.00$), rate triples ($3.60 \rightarrow 10.80 \times 10^{-5}$) $\Rightarrow$ Order with respect to $H_2 = 1$ (2). (ii) Rate = $k[NO]^2[H_2]$ (1). $k = \text{Rate} / ([NO]^2[H_2]) = 3.60 \times 10^{-5} / ((1.50 \times 10^{-3})^2 \times (2.00 \times 10^{-3})) = 8.00 \times 10^3$ (or 8000) (2). Units: $(\text{mol dm}^{-3} \text{ s}^{-1}) / (\text{mol dm}^{-3})^3 = \text{dm}^6 \text{ mol}^{-2} \text{ s}^{-1}$ (1). (iii) $R_4 = (8000) \times (4.50 \times 10^{-3})^2 \times (4.00 \times 10^{-3}) = 6.48 \times 10^{-4} \text{ mol dm}^{-3} \text{ s}^{-1}$ (1).	[9]
(c) Compression Effects	(i) Compressing volume to $1/3$ triples the concentration of each gas ($3\times$) (1). (ii) Rate proportional to $[NO]^2[H_2] \Rightarrow$ new rate = $(3)^2 \times (3)^1 = 27$ times initial rate (3). (iii) The rate constant k is unaffected because k is strictly constant at a fixed temperature and is independent of concentration or volume changes (2).	[6]

QUESTION 36: Decolorisation of Crystal Violet (15 Marks)

(a) Calibration Protocol	Pipette precise volumes of $1.00 \times 10^{-4} \text{ mol dm}^{-3}$ stock CV+ into volumetric flasks to produce 5 standard concentrations (e.g. 2.0, 4.0, 6.0, 8.0, $10.0 \times 10^{-5} \text{ mol dm}^{-3}$) (1). Zero colorimeter using cuvette of pure deionised water (blank) (1). Measure absorbance of each standard at 590 nm (1). Plot absorbance against [CV+] to confirm a straight line passing through the origin (Beer-Lambert Law) (1).	[4]
(b) Isolation Method	[OH ⁻] is 5000 times greater than [CV+] ($0.100 \text{ vs } 2.00 \times 10^{-5} \text{ mol dm}^{-3}$) (1). Even after complete reaction, [OH ⁻] drops by at most 0.02%, so [OH ⁻] is effectively constant (1). Rate = $k[\text{CV}+]^m [\text{OH}^-]^n = k'[\text{CV}+]^m$ where $k' = k[\text{OH}^-]^n$ (pseudo-order rate law) (1).	[3]
(c) Proof & k'	$A = \epsilon \cdot l \cdot [\text{CV}+] \Rightarrow [\text{CV}+] = A / (\epsilon \cdot l)$ (1). For first-order: $\ln[\text{CV}+]_t = -k' t + \ln[\text{CV}+]_0$ (1). Substituting gives $\ln(A_t / \epsilon \cdot l) = -k' t + \ln(A_0 / \epsilon \cdot l) \Rightarrow \ln A_t - \ln(\epsilon \cdot l) = -k' t + \ln A_0 - \ln(\epsilon \cdot l) \Rightarrow \ln A_t = -k' t + \ln A_0$ (1). Linear plot confirms order $m = 1$ with respect to crystal violet (1). $k' = -\text{gradient} = 0.0185 \text{ s}^{-1}$ (1).	[5]
(d) Order in OH⁻ and k	When [OH ⁻] doubles from 0.100 to 0.200 mol dm^{-3} , gradient k' doubles from 0.0185 to 0.0370 $\text{s}^{-1} \Rightarrow$ Order $n = 1$ with respect to OH ⁻ (1). Rate = $k[\text{CV}+][\text{OH}^-]$ (1). $k = k' / [\text{OH}^-] = 0.0185 / 0.100 = 0.185 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ (1).	[3]

QUESTION 37: Acid-Catalysed Inversion of Sucrose (15 Marks)

(a) Polarimeter Principles	Monochromatic light passes through a polarising filter to produce plane-polarised light (1). Light passes through the sample cell containing optically active sugar solution; plane of polarisation is rotated by angle α (1). Analyser prism rotated until light transmission matches reference point (1). Named 'inversion' because optical rotation inverts from positive dextrorotatory (+66.5 deg) to net negative laevorotatory (-19.7 deg) because fructose rotates light more strongly to the left (-92.4 deg) than glucose rotates to the right (+52.7 deg) (1).	[4]
(b) Half-Life Proof & k	At $t = 0 \text{ min}$: $(\alpha_0 - \alpha_{\infty}) = 32.0 \text{ deg}$. First half-life: drops to 16.0 deg at $t = 40 \text{ min} \Rightarrow t_{1/2}(1) = 40 \text{ min}$ (1). Second half-life: drops to 8.0 deg at $t = 80 \text{ min} \Rightarrow t_{1/2}(2) = 80 - 40 = 40 \text{ min}$ (1). Constant half-life (40 min) confirms first order with respect to sucrose (1). $k = \ln 2 / t_{1/2} = 0.693 / 40 \text{ min} = 0.0173 \text{ min}^{-1}$ (or $2.89 \times 10^{-4} \text{ s}^{-1}$) (1).	[4]
(c) Dilatometry Justification	During hydrolysis, a molecule of water is consumed and incorporated into two separate monosaccharide molecules (1). The glucose and fructose molecules pack differently with water, forming more compact hydration shells (1). This causes a slight net volume contraction of the solution, which can be measured precisely as a meniscus fall in a narrow capillary dilatometer (1).	[3]
(d) Conductivity Unsuitability	All reactants (sucrose, water) and products (glucose, fructose) are neutral non-ionic covalent molecules (1). The catalyst H^+ and its counterion remain constant in concentration, so electrical conductivity remains completely unchanged throughout the reaction (1).	[2]

(e) HCl vs CH₃COOH	HCl is a strong acid that fully dissociates ($[H^+] = 1.0 \text{ mol dm}^{-3}$), whereas CH ₃ COOH is a weak acid that only partially dissociates ($[H^+] \text{ approx } 0.0042 \text{ mol dm}^{-3}$) (1). Since rate is directly proportional to $[H^+]$, the initial rate using HCl will be roughly 240 times faster than with ethanoic acid (1).	[2]
--------------------------------------	---	------------