

The UK Crop Microbiome Cryobank

Payton Yau

2023-08-10

The UK Crop Microbiome Cryobank

The UK Crop Microbiome Cryobank (UK-CMCB) project integrates genomic (DNA) data with a cryobank collection of samples for the soil microbiomes of the UK major crop plant systems. Microbiomes are all the microbes present in any one ecosystem. For this project the microbiomes are from the rhizosphere (the soil surrounding the crop plant roots) and from bulk soil (soil outside the rhizosphere). A beneficial microbiome results in a healthy plant and can lead to improved crop yield and improved crop quality.

The UK-CMCB will provide a facility for researchers to source data and samples, including living microbial material, as well as genomic and metagenomic sequences from different soil microbiome environments. Specifically the project will produce a microbiome resource comprising

An integrated cryopreserved collection of samples (rhizosphere and bulk soil, bacterial and fungal isolates and DNA) from crop plant systems (initially barley, oats, oil seed rape, sugar beet and wheat) An open access AgMicrobiomeBase of microbiome data and associated meta-data linked to current public resources such as MGnify.

In various programming languages, including R, features are enhanced through external packages. These packages encompass a range of functions, datasets, and documentation that can be imported into R to extend its capabilities. External R packages are available from diverse sources such as CRAN (The Comprehensive R Archive Network), Bioconductor, GitHub, and more. The process of acquiring packages varies. After installation, a package must be loaded into the R session using the `library()` function before its functions can be utilised.

Qiime2 to Phyloseq Qiime2 is a powerful tool for microbial community analysis which we used for sequencing analysis; Phyloseq is a package in R that provides a flexible and comprehensive framework for the analysis of high-throughput sequencing data. To convert Qiime2 data to Phyloseq, users can use an external package called Qiime2R for the conversion. Qiime2R facilitates the transformation of Qiime2 data structures into phyloseq data structures within the R programming environment.

Once the data is imported, users can manipulate, analyse, and visualise the data using the various functions available in Phyloseq. This conversion allows users to take advantage of the many analysis tools available in R, such as ggplot2 for data visualisation, dplyr for data manipulation, and vegan for ecological community analysis.

```
# Download qiime2R from Github
# if (!requireNamespace("devtools", quietly = TRUE)){install.packages("devtools")}
# devtools::install_github("jbisanz/qiime2R")
library("qiime2R")

# Download phyloseq from Bioconductor
# if (!require("BiocManager", quietly = TRUE))
```

```
# install.packages("BiocManager")
# BiocManager::install("phyloseq")
library("phyloseq")

# Convert qiime2 results to phyloseq format
physeq <- qza_to_phyloseq(
  features = "~/GitHub/agmicrobiomebase/16s/[Qiime2]Silva_138/428_228_220_table_silva138-with-phyla-no-
  taxonomy = "~/GitHub/agmicrobiomebase/16s/[Qiime2]Silva_138/428_228_220_taxonomy_silva138.qza",
  metadata = "~/GitHub/agmicrobiomebase/16s/meta-table.txt"
  #, tree = "rooted-tree.qza"
)

physeq ## confirm the object
```

```
## phyloseq-class experiment-level object
## otu_table() OTU Table: [ 266974 taxa and 327 samples ]
## sample_data() Sample Data: [ 327 samples by 15 sample variables ]
## tax_table() Taxonomy Table: [ 266974 taxa by 7 taxonomic ranks ]
```

Remove unwanted (failed and controls) samples before the normalisation Removing unwanted samples before normalisation is a common step in microbiome data analysis pipelines. In many cases, some samples may fail during sequencing or quality control, while others may be controls or blanks that are not of interest. These samples can introduce noise and bias in downstream analyses if not removed.

In this script, the unwanted samples are removed using the **subset_samples** function from the **phyloseq** package. This function subsets a **phyloseq** object by samples that meet a certain condition, in this case, those that have a certain type (e.g., not failed or control samples).

By removing the unwanted samples before normalization, the remaining samples can be normalised based on their true biological variation, allowing for more accurate comparisons between samples.

```
# sample_names(physeq)
# rank_names(physeq) # "Kingdom" "Phylum" "Class" "Order" "Family" "Genus" "Species"

## unwanted samples removal (including failed samples)
physeq.ori <- subset_samples(physeq, Analysis == "Include")

## remove object
rm(physeq)
```

normalisation using ConQUR Constrained Quantile Normalisation (ConQUR) <https://www.nature.com/articles/s41467-022-33071-9> is a normalization technique commonly used in the analysis of high-throughput sequencing data, particularly in microbiome studies. It is a type of **quantile normalisation** approach that preserves the relative abundances of taxa between samples while simultaneously removing systematic technical variations that can arise due to differences in sequencing depth, PCR amplification bias, or other factors. ConQUR uses kernel density estimation to model the distribution of taxon abundances across all samples, and then constrains the normalisation process to maintain the relative position of each taxon within that distribution.

```
# devtools::install_github("wdl2459/ConQuR")
library(ConQuR)
```

```
# Download phyloseq from CRAN
# install.packages("doParallel")
library(doParallel)
```

```
# Convert ASV table to a data frame and transpose
B <- as.data.frame(physeq.ori@otu_table) # taxa
B <- t(B)
B <- as.data.frame(B)
```

```
# Extract batch ID from sample data
batchid = physeq.ori@sam_data$Plate # batchid
```

```
# Extract covariates
D = physeq.ori@sam_data[, c('Type', 'Soil', 'Location')] #covar
summary(D)
```

```
##           Type      Soil      Location
## Barley      :44      CL:68      AN:35
## Beans       :39      CY:69      BE:27
## Bulksoil    :45      SC:70      B0:35
## Oats        :44      SL:97      BU:35
## OilseedRape:43              HE:35
## Sugarbeet   :45              SH:70
## Wheat       :44              Y0:67
```

```
# Correct for batch effects using ConQuR package
```

```
options(warn=-1) # required to call
taxa_correct1 = ConQuR(tax_tab = B, batchid = batchid, covariates = D, batch_ref="1") # warning message
```

```
# Transpose the corrected matrix and convert it to a data frame
```

```
taxa_correct2 <- t(taxa_correct1)
taxa_correct2 <- as.data.frame(taxa_correct2)
```

```
# Create new ASV table, taxonomy table, and sample data
```

```
ASV = otu_table(taxa_correct2, taxa_are_rows = TRUE)
TAXA = tax_table(physeq.ori)
sampledata = sample_data(physeq.ori)
```

```
# repack the objects into a level 4 phyloseq structural data
```

```
physeq.norm = phyloseq(ASV, TAXA, sampledata)
```

```
# remove
```

```
rm(B, D, batchid, taxa_correct1, taxa_correct2, ASV, TAXA, sampledata, to_skip)
```

```
# install.packages("tidyverse")
library("tidyverse")
```

```
## (1) Sub-samples for Wheat and Bulk Soil groups for the comparisons
```

```
physeq.W.B <- physeq.norm %>% subset_samples(Type %in% c("Wheat", "Bulksoil"))
```

```
## (2) Subset the samples in the phyloseq object that belong to "Wheat"
## or other crops "Barley", "Beans", "Bulksoil", "Oats", "OilseedRape", "Sugarbeet"
physeq.wheat <- subset_samples(physeq.norm, Type=="Wheat")

## (3A) Merge the replicate samples for each Group
physeq.wheat.group = merge_samples(physeq.wheat, "Group") # Sum between replicate samples

# (3B) repair factors in the sample metadata
sample_data(physeq.wheat.group)$Group <- levels(sample_data(physeq.norm)$Group)[get_variable(physeq.wheat.group, "Group")]
sample_data(physeq.wheat.group)$Soil <- levels(sample_data(physeq.norm)$Soil)[get_variable(physeq.wheat.group, "Soil")]

## (4) Further subgroup for WH.CL.BO vs WH.CL.YO
physeq.wheat.vs <- physeq.wheat %>% subset_samples(Group %in% c("WH.CL.BO", "WH.CL.YO"))
```

Subgrouping

Bata diversity Beta diversity is a measure of the differences or similarities between two or more microbial communities or samples. It quantifies the degree of variation in microbial composition among different samples. Beta diversity analysis can help identify patterns in the distribution of microbial communities and reveal relationships between different factors such as environment, host, and other ecological variables.

Non-Metric Multidimensional Scaling (NMDS) and Principal Coordinates Analysis (PCoA) are both ordination techniques used in analyzing beta diversity, which captures the variation in species composition between different samples or sites. However, they have some differences in terms of methodology and interpretation:

NMDS (Non-Metric Multidimensional Scaling) NMDS is a method that aims to represent the pairwise dissimilarities between samples in a lower-dimensional space while preserving the rank order of these dissimilarities. It is considered a non-metric method because it doesn't assume a linear relationship between the original dissimilarities and the distances in the ordination space. NMDS emphasizes preserving the relative ranking of dissimilarities, making it more suitable for cases where the distances between samples are not well-preserved in a lower-dimensional space. NMDS produces an ordination plot where the positions of samples are arranged based on their pairwise dissimilarities. The distances between points on the plot are not directly interpretable but reflect the similarities or dissimilarities between samples.

PCoA (Principal Coordinates Analysis) PCoA is also used to visualize and analyse beta diversity by representing samples in a lower-dimensional space. It is a type of metric multidimensional scaling. PCoA is based on a metric approach, which means it attempts to preserve the actual distances between samples as closely as possible in the lower-dimensional space. PCoA produces an ordination plot where the distances between points reflect the actual dissimilarities between samples. Unlike NMDS, PCoA may not perform as well with non-linear or rank-based dissimilarity measures.


```
# install.packages("ggplot2")
library("ggplot2")

# method options: NMDS / PCoA
pcoa <- ordinate(physeq = physeq.W.B, method = "NMDS", distance = "bray")

## Square root transformation
## Wisconsin double standardization
## Run 0 stress 0.2043982
## Run 1 stress 0.2275976
## Run 2 stress 0.1955895
## ... New best solution
```

```
## ... Procrustes: rmse 0.03743642  max resid 0.1725148
## Run 3 stress 0.2362841
## Run 4 stress 0.2044245
## Run 5 stress 0.2061967
## Run 6 stress 0.2061968
## Run 7 stress 0.1955995
## ... Procrustes: rmse 0.002549282  max resid 0.01724757
## Run 8 stress 0.1955996
## ... Procrustes: rmse 0.002577285  max resid 0.01762842
## Run 9 stress 0.1955842
## ... New best solution
## ... Procrustes: rmse 0.001086866  max resid 0.007247389
## ... Similar to previous best
## Run 10 stress 0.2390976
## Run 11 stress 0.2108416
## Run 12 stress 0.2183071
## Run 13 stress 0.2195094
## Run 14 stress 0.2314068
## Run 15 stress 0.2339275
## Run 16 stress 0.2404871
## Run 17 stress 0.2061711
## Run 18 stress 0.2113607
## Run 19 stress 0.2189368
## Run 20 stress 0.1955842
## ... Procrustes: rmse 2.543653e-05  max resid 0.0001130426
## ... Similar to previous best
## *** Best solution repeated 2 times
```

```
# Plot ordination
plot_ordination(
  physeq = physeq.W.B,
  ordination = pcoa,
  color = "Type",
  shape = "Soil.Location"
) +
  theme_classic() +
  geom_point(aes(color = Type), alpha = 1, size = 3) +
  theme(
    text = element_text(size = 18, colour = "black"),
    axis.ticks = element_line(colour = "black", size = 1.1),
    axis.line = element_line(colour = "black", size = 1.1),
    axis.text.x = element_text(colour = "black", angle = 0, hjust = 0.5, size = 13, face = "bold"),
    axis.text.y = element_text(colour = "black", angle = 0, hjust = 0.5, size = 13, face = "bold"),
    axis.title.y = element_text(color = "black", size = 20, face = "bold"),
    axis.title.x = element_text(color = "black", size = 20, face = "bold")
  ) +
  scale_color_brewer(palette = "Dark2") +
  scale_fill_brewer(palette = "Dark2") +
  scale_shape_manual(values = c(15, 17, 3, 4, 16, 18, 21, 22, 23)) # Set custom shapes
```



```
# Clean up by removing objects that are no longer needed
rm(pcoa, physeq.W.B)
```

Alpha diversity Alpha diversity refers to the diversity of species within a single ecosystem or habitat. It is measured by analysing the number and distribution of species within a specific area or sample. Alpha diversity indices take into account the richness (number of species) and evenness (relative abundance of species) of a community. It provides insights into the complexity and stability of ecosystems. A high level of alpha diversity indicates a more complex ecosystem with a greater number of species, which is often associated with greater ecological resilience and stability. In contrast, a low level of alpha diversity can be an indicator of ecosystem disturbance, degradation, or vulnerability.

Please noted that different types of alpha diversity metrics capture various aspects of biodiversity within a specific community. Here's a brief explanation of each:

Observed: This metric simply counts the number of unique species (or operational taxonomic units) present in a sample. It provides a basic measure of species richness.

Chao1: Chao1 estimates the total number of species by considering the number of rare or singleton species. It takes into account the number of singletons (species observed only once) and doubletons (species observed only twice).

ACE (Abundance-based Coverage Estimator): Similar to Chao1, ACE also estimates species richness by accounting for rare species, but it also considers their abundance in the community.

Shannon Diversity Index: This index takes into account both species richness and evenness in the community. It considers the number of species present as well as their relative abundances.

Simpson Diversity Index: Simpson's index gives more weight to dominant species in the community. It reflects the probability that two randomly selected individuals belong to different species.


Inverse Simpson Diversity Index (InvSimpson): This index is the reciprocal of the Simpson index and is useful for emphasizing the dominance of a few species.

Fisher's Alpha: Fisher's alpha is a measure of species richness that takes into account the distribution of individuals among species. It's particularly useful for comparing species diversity between different communities.

Here, we pick Shannon as an example for the work

```
# available measurements: "Observed" "Chao1" "ACE" "Shannon" "Simpson" "InvSimpson" "Fisher"
# Calculate alpha diversity (Shannon) and store it in alpha.object
alpha.object <- cbind(
  x = sample_data(physeq.wheat),
  y = estimate_richness(physeq.wheat, measures = 'Shannon')
)

# Create a plot for alpha diversity
ggplot(data = alpha.object, aes(x = x.Group, y = Shannon, color = x.Group, fill = x.Group)) +
  theme_classic() +
  labs(
    x = element_blank(), # No x-axis label
    y = "Alpha Diversity (Shannon)" # y-axis label
  ) +
  geom_point(size = 1.75) + # Add points
  geom_boxplot(alpha = 0.5) + # Add boxplot with transparency
  theme(
    text = element_text(size = 18, colour = "black"),
    axis.ticks = element_line(colour = "black", size = 1.1),
    axis.line = element_line(colour = "black", size = 1.1),
    axis.text.x = element_text(colour = "black", angle = 270, size = 13, face = "bold"),
    axis.text.y = element_text(angle = 0, hjust = 0, colour = "black", size = 13, face = "bold"),
    axis.title.y = element_text(color = "black", size = 15, face = "bold"),
    legend.position = "none" # Hide legend
  )
```



```
# Clean up by removing the alpha.object
rm(alpha.object)
```

Calculate the number of taxa on each level and each group The purpose of this process is to visualise the distribution of the number of matched abundance across different groups and to identify any patterns in the distribution of the processed abundance within individual group.

```
# Create an empty list to store genus-level abundance data for each taxonomic level
gentab_levels <- list()

# Set observation threshold
observationThreshold <- 1

# Define the taxonomic levels
genus_levels <- c("Kingdom", "Phylum", "Class", "Order", "Family", "Genus", "Species")

# loop through all the taxonomic levels
for (level in genus_levels) {

  # create a factor variable for each level
  genfac <- factor(tax_table(physeq.wheat.group)[, level])

  # calculate the abundance of each genus within each sample
  gentab <- apply(otu_table(physeq.wheat.group), MARGIN = 1, function(x) {
    tapply(x, INDEX = genfac, FUN = sum, na.rm = TRUE, simplify = TRUE)
  })
}
```



```

})

# calculate the number of samples in which each genus is observed above the threshold
level_counts <- apply(gentab > observationThreshold, 2, sum)

# create a data frame of level counts with genus names as row names
BB <- as.data.frame(level_counts)
BB$name <- row.names(BB)

# add the data frame to the gentab_levels list
gentab_levels[[level]] <- BB
}

# Combine all level counts data frames into one data frame
B2 <- gentab_levels %>% reduce(full_join, by = "name")

# Set row names and column names
rownames(B2) <- B2$name
B2$name <- NULL
colnames(B2)[1:7] <- genus_levels

# Print the resulting data frame
print(B2)

```

```

##           Kingdom Phylum Class Order Family Genus Species
## WH.CL.BO         3      41   119   262   388   583    302
## WH.CL.YO         3      44   129   285   409   630    319
## WH.CY.BU         2      37   103   242   349   569    290
## WH.CY.YO         2      41   119   275   401   635    300
## WH.SC.HE         2      36    97   218   327   482    233
## WH.SC.SH         3      42   124   277   401   612    280
## WH.SL.AN         3      39   118   258   382   589    308
## WH.SL.BE         2      41   124   280   405   643    301
## WH.SL.SH         3      41   119   274   405   626    297

```

```

# Clean up by removing unnecessary objects
rm(gentab_levels, BB)

```

```

# Calculate the sum of ASVs across samples and create a data frame
ASV <- sample_sums(physeq.wheat)
ASV <- as.data.frame(ASV)
ASV$Wheat <- rownames(ASV)


# Remove the suffix from the ASV names
new_names <- sub("-\\d+$", "", ASV$Wheat)

# Update the row names of the data frame
ASV$Wheat <- new_names

# Create a scatter plot with jittered points and overlaid boxplots

```

```
ggplot(ASV, aes(x = Wheat, y = ASV, colour = Wheat)) +
  geom_point(alpha = 1, position = "jitter", size = 4) +
  geom_boxplot(alpha = 0, colour = "black", size = 0.8) +
  theme_classic() +
  theme(text = element_text(size=15, colour = "black"),
        axis.ticks = element_line(colour = "black", size = 1.25),
        axis.line = element_line(colour = 'black', size = 1.25),
        axis.text.x = element_text(angle=45, hjust=1, colour = "black", size = 13),
        axis.text.y = element_text(angle=0, hjust=0.5, colour = "black", size = 13),
        axis.title.y = element_text(color="black", size=15, face="bold"),
        legend.position = "none")
```



Calculate the sum of ASV (Wheat)

```
# Subset the taxa to Genus from physeq.wheat
physeq.wheat.genus <- subset_taxa(physeq.wheat, Genus == "Azospirillum")


# Calculate the total abundance of Azospirillum for each sample
meta = physeq.wheat.genus@sam_data
otudf = as.data.frame(t(as.data.frame(physeq.wheat.genus@otu_table)))
meta$Azospirillum = rowSums(otudf)

# Plot a graph of the abundance of Azospirillum for each sample grouped by Group:
ggplot(subset(meta, Group %in% c("WH.CL.BO", "WH.CL.YO",
```

```

        "WH.CY.BU", "WH.CY.YO",
        "WH.SC.HE", "WH.SC.SH",
        "WH.SL.AN", "WH.SL.BE",
        "WH.SL.SH")),
    aes(x = Group, y = Azospirillum, colour = interaction(Group))) +
  geom_point(alpha = 1, position = "jitter", size = 4) +
  geom_boxplot(alpha = 0, colour = "black", size = 0.8) +
  theme_classic() +
  theme(text = element_text(size=15, colour = "black"),
        axis.ticks = element_line(colour = "black", size = 1.25),
        axis.line = element_line(colour = "black", size = 1.25),
        axis.text.x = element_text(angle=45, hjust=1, colour = "black", size = 13),
        axis.text.y = element_text(angle=0, hjust=0.5, colour = "black", size = 13),
        axis.title.y = element_text(color="black", size=15, face="bold"), legend.position = "none")

```



Plotting for the abundance of one specific bacteria

```

# Clean up by removing unnecessary objects
rm(physeq.wheat.genus, meta, otudf)

```

Plotting the top 30 taxa at family level

```


## Transform normalised ASVs to proportions
proportions = transform_sample_counts(physeq.wheat.group, function(x) 100 * x/sum(x))

```

```
##
Top30ASVs = names(sort(taxa_sums(proportions), TRUE)[1:71])
Taxtab30 = cbind(tax_table(proportions), Family30 = NA)
Taxtab30[Top30ASVs, "Family30"] <- as(tax_table(proportions)[Top30ASVs, "Family"], "character")
tax_table(proportions) <- tax_table(Taxtab30)

Rgsm30 = prune_taxa(Top30ASVs, proportions)

# plotting
title = "Wheat - Top 30"
plot_bar(Rgsm30, "Group", fill = "Family", title = title) +
  coord_flip() +
  ylab("Percentage of Sequences") + ylim(0, 30) + theme_classic() +
  theme(text = element_text(size=15, colour = "black"),
        axis.ticks = element_line(colour = "black", size = 1.25),
        axis.line = element_line(colour = "black", size = 1.25),
        axis.text.x = element_text(angle=45, hjust=1, colour = "black", size = 13),
        axis.text.y = element_text(angle=0, hjust=0.5, colour = "black", size = 13),
        axis.title.y = element_text(colour="black", size=15, face="bold"),
        legend.position = "right",
        legend.text = element_text(size = 9.5),
        legend.key.height= unit(0.45, 'cm'),
        legend.key.width= unit(0.45, 'cm')
  )
```



```
# Clean up by removing unnecessary objects
# rm(proportions, Top30ASVs, Taxtab30, Rgsm30, title)
```

Pairwise comparison using PERMANOVA Pairwise PERMANOVA is a statistical method used to compare multiple groups or treatments in ecological and microbial community studies. It assesses dissimilarity between samples and provides a p-value to determine the significance of observed differences. This approach is valuable for targeted group comparisons, allowing researchers to investigate the effects of specific factors on microbial communities and uncover significant variations in community composition. By considering within- and between-group variation, pairwise PERMANOVA provides robust statistical analysis and insights into microbial community dynamics and functioning.

```
# devtools::install_github("pmartinezarbizu/pairwiseAdonis/pairwiseAdonis")
library("pairwiseAdonis")
```

```
metdat = as.data.frame(as.matrix(physeq.wheat@sam_data))
dat = as.data.frame(t(as.data.frame(physeq.wheat@otu_table)))
pairwise.adonis(dat, metdat$Group, sim.function = "vegdist",
  sim.method = "bray", p.adjust.m = "bonferroni",
  reduce = NULL, perm = 10000)
```

##		pairs	Df	SumsOfSqs	F.Model	R2	p.value	p.adjusted
## 1	WH.CL.BO vs	WH.CL.YO	1	0.6632405	4.879979	0.3788810	0.00829917	0.2987701
## 2	WH.CL.BO vs	WH.CY.BU	1	1.1433027	8.500806	0.5151752	0.00709929	0.2555744
## 3	WH.CL.BO vs	WH.CY.YO	1	1.1251093	7.533254	0.4849759	0.00679932	0.2447755
## 4	WH.CL.BO vs	WH.SC.HE	1	1.0304398	7.540431	0.4852138	0.00929907	0.3347665
## 5	WH.CL.BO vs	WH.SC.SH	1	0.6707283	3.742479	0.3187129	0.00709929	0.2555744
## 6	WH.CL.BO vs	WH.SL.AN	1	0.4921162	3.662869	0.3140624	0.00609939	0.2195780
## 7	WH.CL.BO vs	WH.SL.BE	1	0.5277124	3.485304	0.3323989	0.00849915	0.3059694
## 8	WH.CL.BO vs	WH.SL.SH	1	0.5729351	3.795038	0.3217487	0.00739926	0.2663734
## 9	WH.CL.YO vs	WH.CY.BU	1	1.2318530	9.056919	0.5309821	0.00869913	0.3131687
## 10	WH.CL.YO vs	WH.CY.YO	1	0.9002886	5.967262	0.4272321	0.00639936	0.2303770
## 11	WH.CL.YO vs	WH.SC.HE	1	1.1590697	8.388465	0.5118518	0.00919908	0.3311669
## 12	WH.CL.YO vs	WH.SC.SH	1	0.5929366	3.280619	0.2908190	0.00719928	0.2591741
## 13	WH.CL.YO vs	WH.SL.AN	1	0.5857893	4.311344	0.3501928	0.00769923	0.2771723
## 14	WH.CL.YO vs	WH.SL.BE	1	0.7612919	4.970999	0.4152535	0.00829917	0.2987701
## 15	WH.CL.YO vs	WH.SL.SH	1	0.5537116	3.631171	0.3121931	0.00849915	0.3059694
## 16	WH.CY.BU vs	WH.CY.YO	1	1.2662987	8.472820	0.5143515	0.00879912	0.3167683
## 17	WH.CY.BU vs	WH.SC.HE	1	0.8697659	6.359930	0.4428942	0.00789921	0.2843716
## 18	WH.CY.BU vs	WH.SC.SH	1	0.9066775	5.056137	0.3872613	0.00889911	0.3203680
## 19	WH.CY.BU vs	WH.SL.AN	1	0.9651275	7.178095	0.4729246	0.00839916	0.3023698
## 20	WH.CY.BU vs	WH.SL.BE	1	1.0162083	6.706443	0.4892913	0.00619938	0.2231777
## 21	WH.CY.BU vs	WH.SL.SH	1	0.9931700	6.574176	0.4510839	0.00849915	0.3059694
## 22	WH.CY.YO vs	WH.SC.HE	1	1.5272711	10.073279	0.5573576	0.00879912	0.3167683
## 23	WH.CY.YO vs	WH.SC.SH	1	0.8337784	4.293818	0.3492665	0.00749925	0.2699730
## 24	WH.CY.YO vs	WH.SL.AN	1	1.2311524	8.245421	0.5075535	0.00799920	0.2879712
## 25	WH.CY.YO vs	WH.SL.BE	1	1.0555054	6.263800	0.4722478	0.00859914	0.3095690
## 26	WH.CY.YO vs	WH.SL.SH	1	0.8949815	5.393719	0.4027051	0.00779922	0.2807719
## 27	WH.SC.HE vs	WH.SC.SH	1	0.9505140	5.237452	0.3956541	0.00889911	0.3203680
## 28	WH.SC.HE vs	WH.SL.AN	1	0.8816640	6.453574	0.4465037	0.00889911	0.3203680
## 29	WH.SC.HE vs	WH.SL.BE	1	0.9356846	6.075959	0.4646664	0.00809919	0.2915708
## 30	WH.SC.HE vs	WH.SL.SH	1	0.9782871	6.384301	0.4438381	0.00809919	0.2915708
## 31	WH.SC.SH vs	WH.SL.AN	1	0.6084657	3.395808	0.2979875	0.00479952	0.1727827

```
## 32 WH.SC.SH vs WH.SL.BE 1 0.6448442 3.182159 0.3125230 0.01609839 0.5795420
## 33 WH.SC.SH vs WH.SL.SH 1 0.3204480 1.636623 0.1698337 0.03729627 1.0000000
## 34 WH.SL.AN vs WH.SL.BE 1 0.6978229 4.610161 0.3970798 0.00949905 0.3419658
## 35 WH.SL.AN vs WH.SL.SH 1 0.6009452 3.981599 0.3323095 0.00869913 0.3131687
## 36 WH.SL.BE vs WH.SL.SH 1 0.6206904 3.643469 0.3423197 0.00909909 0.3275672
## sig
## 1
## 2
## 3
## 4
## 5
## 6
## 7
## 8
## 9
## 10
## 11
## 12
## 13
## 14
## 15
## 16
## 17
## 18
## 19
## 20
## 21
## 22
## 23
## 24
## 25
## 26
## 27
## 28
## 29
## 30
## 31
## 32
## 33
## 34
## 35
## 36
```

```
# Clean up by removing unnecessary objects
rm(metdat, dat)
```

Upset plot using UpsetR

When it comes to representing sets visually, the go-to option is usually a Venn diagram. These diagrams work well when dealing with up to five sets, providing a clear visualisation. However, as the dataset expands, such as when dealing with five sets, deriving the desired insights from the diagram becomes more complex. As a result, considering an UpSet graph for data visualisation becomes an appealing choice. UpSet graphs offer a more streamlined way to display intersections and complements, particularly when dealing with larger

datasets or multiple sets. This option ensures a more intuitive and informative representation of the data.

```
# BiocManager::install("microbiome")
library("microbiome")

# devtools::install_github("mikemc/speedyseq")
library("speedyseq")

library("UpSetR")
library("plyr")
library("reshape2")

# Aggregate taxa at the genus level
B <- aggregate_taxa(physeq.wheat.group, "Genus", verbose = TRUE)
```

```
## [1] "Remove taxonomic information below the target level"
## [1] "Mark the potentially ambiguous taxa"
## [1] "--- split"
## [1] "--- sum"
## [1] "Create phyloseq object"
## [1] "Remove ambiguous levels"
## [1] "--- unique"
## [1] "--- Rename the lowest level"
## [1] "--- rownames"
## [1] "--- taxa"
## [1] "Convert to taxonomy table"
## [1] "Combine OTU and Taxon matrix into Phyloseq object"
## [1] "Add the metadata as is"
```

```
# Remove undesired genera
# B2 <- subset_taxa(B, !get("Genus") %in% c("uncultured", "Unknown"))

# Remove unwanted taxon names
taxa_to_remove <- c("uncultured", "Unknown")
B2 <- subset_taxa(B, !get("Genus") %in% taxa_to_remove)
```

```
# Convert to tibble, rename columns, select relevant columns, group by Sample, and keep top 100 abundances
# Convert data to a tibble and perform necessary operations
```

```
D <- as_tibble(B2) %>%
  mutate(Sample = .sample, ASV = .otu, Abundance = .abundance) %>%
  select(Sample, Abundance, Genus) %>%
  group_by(Sample) %>%
  filter(rank(desc(Abundance)) <= 100) %>% # Filter the top 100
  ungroup()
```

```
# Remove the Abundance column
D$Abundance <- NULL
```

```
# Rename the second column to "ASV"
names(D)[2] <- "ASV"
```

```
# Convert data from long to wide format
```

```

E <- dcast(D, ASV ~ Sample)


# Define a binary function
binary_fun <- function(x) {
  x[is.na(x)] <- 0
  ifelse(x > 0, 1, 0)
}

# Apply the binary function to columns 2 to 10
temp_df <- apply(E[2:10], 2, binary_fun)
temp_df <- as.data.frame(temp_df)

# Create an UpSet plot
upset_plot <- upset(temp_df, sets = colnames(temp_df), sets.bar.color = "#56B4E9",
  order.by = "freq", empty.intersections = "on")

# Print the UpSet plot
print(upset_plot)

```



```

# Clean up by removing unnecessary objects
# rm(B, B2, D, E, binary_fun, temp_df, upset_plot, physeq.wheat.group)

# https://github.com/joey711/phyloseq/issues/901
# Filter taxa with mean abundance > 0.1
physeq2 = filter_taxa(physeq.wheat.vs, function(x) mean(x) > 0.1, TRUE)

```



```
# Transform sample counts to relative abundances
physeq3 = transform_sample_counts(physeq2, function(x) x / sum(x))

# Aggregate taxa at the Phylum level
glom <- tax_glom(physeq3, taxrank = 'Phylum')

# Convert the aggregated phyloseq object to a dataframe
data <- psmelt(glom)

# Convert Phylum column to character
data$Phylum <- as.character(data$Phylum)

# Rename phyla with abundance < 1% to "< 1% abund."
data$Phylum[data$Abundance < 0.01] <- "< 1% abund."


# Calculate medians of abundance for each Phylum
medians <- ddply(data, ~Phylum, function(x) c(median = median(x$Abundance)))

# Extract phyla with median abundance <= 0.01
remainder <- medians[medians$median <= 0.01, ]$Phylum

# Rename selected phyla to "Phyla < 1% abund."
data[data$Phylum %in% remainder, ]$Phylum <- "Phyla < 1% abund."

# Rename phyla with abundance < 1% to "Phyla < 1% abund."
data$Phylum[data$Abundance < 0.01] <- "Phyla < 1% abund."

# Plot the data with condensed phyla categories
p <- ggplot(data = data, aes(x = Sample, y = Abundance, fill = Phylum))
p + geom_bar(aes(), stat = "identity", position = "stack") +
  scale_fill_manual(values = c("darkblue", "darkgoldenrod1", "darkseagreen", "darkorchid", "darkolivegreen"),
    theme(text = element_text(size=15, colour = "black"),
      axis.ticks = element_line(colour = "black", size = 1.25),
      axis.line = element_line(colour = 'black', size = 1.25),
      axis.text.x = element_text(angle=45, hjust=1, colour = "black", size = 13),
      axis.text.y = element_text(angle=0, hjust=0.5, colour = "black",size = 13),
      axis.title.y = element_text(color="black", size=15,face="bold"), legend.position = "right") # +
```



```
rm(p, data, remainder, medians,glom, physeq3, physeq2)
```


Linear discriminant analysis effect size (LEFSe) Linear discriminant analysis effect size (LEFSe) is a powerful computational tool commonly used in microbiome analysis to identify microbial biomarkers that are differentially abundant between different experimental groups. Microbiome analysis involves studying the composition and diversity of microbial communities present in various environments, such as the human gut, soil, water, and more. LEFSe is specifically designed to address the challenge of identifying microbial taxa that drive differences between different experimental conditions or groups.

LEFSe combines linear discriminant analysis (LDA), which is a statistical technique used for classification and feature selection, with the concept of effect size. Effect size refers to the magnitude of a difference between two groups and provides insights into the biological relevance of the observed differences.

P,S,ALDEx2 and ANCOM-II produce the most consistent results (<https://www.nature.com/articles/s41467-022-28034-z>)

```
# BiocManager::install("microbiomeMarker")
library("microbiomeMarker")
```

```
lef_out <- run_lefse(physeq.wheat.vs, group = "Group",
  norm = "CPM",
  # taxa_rank = "Genus",
  kw_cutoff = 0.01, lda_cutoff = 4)
plot_ef_bar(lef_out)
```



Visualising the marker table as cladogram The default is 4 which is showing the kingdom, Phylum, Class on the cladogram.

```
# plot_cladogram(lef_out, color = c("red", "blue"),
#                 clade_label_level = 6)
# rm(lef_out)
```

```
sessionInfo()
```

```
## R version 4.3.1 (2023-06-16 ucrt)
## Platform: x86_64-w64-mingw32/x64 (64-bit)
## Running under: Windows 10 x64 (build 19045)
##
## Matrix products: default
##
##
## locale:
## [1] LC_COLLATE=English_United Kingdom.utf8
## [2] LC_CTYPE=English_United Kingdom.utf8
## [3] LC_MONETARY=English_United Kingdom.utf8
## [4] LC_NUMERIC=C
## [5] LC_TIME=English_United Kingdom.utf8
##
## time zone: Europe/London
```

```

## tzcode source: internal
##
## attached base packages:
## [1] parallel stats      graphics grDevices utils      datasets methods
## [8] base
##
## other attached packages:
## [1] microbiomeMarker_1.6.0 reshape2_1.4.4      plyr_1.8.8
## [4] UpSetR_1.4.0           speedyseq_0.5.3.9018  microbiome_1.22.0
## [7] pairwiseAdonis_0.4.1  cluster_2.1.4        vegan_2.6-4
## [10] lattice_0.21-8        permute_0.9-7        lubridate_1.9.2
## [13] forcats_1.0.0         stringr_1.5.0        dplyr_1.1.2
## [16] purrr_1.0.1           readr_2.1.4          tidyr_1.3.0
## [19] tibble_3.2.1          ggplot2_3.4.2        tidyverse_2.0.0
## [22] doParallel_1.0.17     iterators_1.0.14     foreach_1.5.2
## [25] ConQuR_2.0            phyloseq_1.44.0      qiime2R_0.99.6
##
## loaded via a namespace (and not attached):
## [1] IRanges_2.34.1          gld_2.6.6
## [3] nnet_7.3-19             DT_0.28
## [5] Biostrings_2.68.1      TH.data_1.1-2
## [7] vctrs_0.6.3            energy_1.7-11
## [9] digest_0.6.33          png_0.1-8
## [11] shape_1.4.6            proxy_0.4-27
## [13] Exact_3.2              ggrepel_0.9.3
## [15] MASS_7.3-60            BiocGenerics_0.46.0
## [17] withr_2.5.0            xfun_0.39
## [19] ggfun_0.1.2            survival_3.5-5
## [21] doRNG_1.8.6            memoise_2.0.1
## [23] ggbeeswarm_0.7.2       MatrixModels_0.5-2
## [25] gmp_0.7-2              tidytree_0.4.5
## [27] zoo_1.8-12             GlobalOptions_0.1.2
## [29] gtools_3.9.4           DEoptimR_1.1-1
## [31] Formula_1.2-5          httr_1.4.6
## [33] rhdf5filters_1.12.1    rhdf5_2.44.0
## [35] rstudioapi_0.15.0      generics_0.1.3
## [37] base64enc_0.1-3        S4Vectors_0.38.1
## [39] zlibbioc_1.46.0        ScaledMatrix_1.8.1
## [41] randomForest_4.7-1.1   statip_0.2.3
## [43] GenomeInfoDbData_1.2.10 ade4_1.7-22
## [45] evaluate_0.21          S4Arrays_1.0.5
## [47] hms_1.1.3              glmnet_4.1-7
## [49] GenomicRanges_1.52.0   irlba_2.3.5.1
## [51] colorspace_2.1-0       ROCR_1.0-11
## [53] readxl_1.4.3           magrittr_2.0.3
## [55] viridis_0.6.4          ggtree_3.8.2
## [57] robustbase_0.99-0      SparseM_1.81
## [59] DECIPHER_2.28.0        scuttle_1.10.2
## [61] matrixStats_1.0.0      class_7.3-22
## [63] Hmisc_5.1-0            pillar_1.9.0
## [65] nlme_3.1-162           decontam_1.20.0
## [67] plotROC_2.3.0          caTools_1.18.2
## [69] compiler_4.3.1         beachmat_2.16.0
## [71] stringi_1.7.12         biomformat_1.28.0

```

## [73] DescTools_0.99.49	stabledist_0.7-1
## [75] minqa_1.2.5	SummarizedExperiment_1.30.2
## [77] crayon_1.5.2	abind_1.4-5
## [79] scater_1.28.0	truncnorm_1.0-9
## [81] timeSeries_4030.106	gridGraphics_0.5-1
## [83] cqrReg_1.2.1	locfit_1.5-9.8
## [85] bit_4.0.5	mia_1.8.0
## [87] rootSolve_1.8.2.3	sandwich_3.0-2
## [89] codetools_0.2-19	multcomp_1.4-25
## [91] BiocSingular_1.16.0	e1071_1.7-13
## [93] lmom_2.9	GetoptLong_1.0.5
## [95] multtest_2.56.0	MultiAssayExperiment_1.26.0
## [97] splines_4.3.1	metagenomeSeq_1.42.0
## [99] circlize_0.4.15	Rcpp_1.0.11
## [101] fastDummies_1.7.3	quantreg_5.96
## [103] sparseMatrixStats_1.12.2	cellranger_1.1.0
## [105] knitr_1.43	blob_1.2.4
## [107] utf8_1.2.3	clue_0.3-64
## [109] lme4_1.1-34	fBasics_4022.94
## [111] checkmate_2.2.0	DelayedMatrixStats_1.22.5
## [113] Rdpack_2.4	expm_0.999-7
## [115] gsl_2.1-8	ggplotify_0.1.2
## [117] Matrix_1.6-0	statmod_1.5.0
## [119] tzdb_0.4.0	pkgconfig_2.0.3
## [121] tools_4.3.1	cachem_1.0.8
## [123] rbibutils_2.2.14	RSQLite_2.3.1
## [125] viridisLite_0.4.2	NADA_1.6-1.1
## [127] DBI_1.1.3	numDeriv_2016.8-1.1
## [129] rmutil_1.1.10	fastmap_1.1.1
## [131] rmarkdown_2.23	scales_1.2.1
## [133] grid_4.3.1	stable_1.1.6
## [135] patchwork_1.1.2	rpart_4.1.19
## [137] farver_2.1.1	mgcv_1.8-42
## [139] yaml_2.3.7	MatrixGenerics_1.12.3
## [141] spatial_7.3-16	foreign_0.8-84
## [143] bayesm_3.1-5	cli_3.6.1
## [145] stats4_4.3.1	lifecycle_1.0.3
## [147] Biobase_2.60.0	mvtnorm_1.2-2
## [149] backports_1.4.1	modeest_2.4.0
## [151] BiocParallel_1.34.2	timechange_0.2.0
## [153] gtable_0.3.3	rjson_0.2.21
## [155] ANCOMBC_2.2.1	ape_5.7-1
## [157] limma_3.56.2	CVXR_1.0-11
## [159] jsonlite_1.8.7	bitops_1.0-7
## [161] bit64_4.0.5	Rtsne_0.16
## [163] yulab.utils_0.0.7	BiocNeighbors_1.18.0
## [165] TreeSummarizedExperiment_2.8.0	highr_0.10
## [167] timeDate_4022.108	lazyeval_0.2.2
## [169] zCompositions_1.4.0-1	htmltools_0.5.5
## [171] glue_1.6.2	Wrench_1.18.0
## [173] XVector_0.40.0	RCurl_1.98-1.12
## [175] treeio_1.24.3	gridExtra_2.3
## [177] boot_1.3-28.1	igraph_1.5.0.1
## [179] R6_2.5.1	DESeq2_1.40.2

## [181] SingleCellExperiment_1.22.0	gplots_3.1.3
## [183] Rmpfr_0.9-3	labeling_0.4.2
## [185] rngtools_1.5.2	Rhdf5lib_1.22.0
## [187] aplot_0.2.0	GenomeInfoDb_1.36.1
## [189] nloptr_2.0.3	compositions_2.0-6
## [191] DirichletMultinomial_1.42.0	DelayedArray_0.26.7
## [193] tidyselect_1.2.0	vipor_0.4.5
## [195] htmlTable_2.4.1	tensorA_0.36.2
## [197] GUniFrac_1.7	rsvd_1.0.5
## [199] munsell_0.5.0	KernSmooth_2.23-21
## [201] data.table_1.14.8	htmlwidgets_1.6.2
## [203] ComplexHeatmap_2.16.0	RColorBrewer_1.1-3
## [205] rlang_1.1.1	lmerTest_3.1-3
## [207] fansi_1.0.4	beeswarm_0.4.0