



National Taiwan University

nontoi

FHVirus, nathanlee726, hhhhaura

2022-11-18

- 1 Contest
- 2 Mathematics
- 3 Data structures
- 4 Numerical
- 5 Number theory
- 6 Combinatorial
- 7 Graph
- 8 Geometry
- 9 Strings
- 10 Various

Contest (1)

.bashrc
.vimrc
template.cpp
troubleshoot.txt
cmp.sh

Mathematics (2)

2.1 Equations

$$ax^2+bx+c=0\Rightarrow x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$$

The extremum is given by $x=-b/2a$.

$$\begin{aligned}ax+by&=e & x&=\frac{ed-bf}{ad-bc} \\ cx+dy&=f & y&=\frac{af-ec}{ad-bc}\end{aligned}$$

In general, given an equation $Ax=b$, the solution to a variable x_i is given by

$$x_i=\frac{\det A'_i}{\det A}$$

where A'_i is A with the i 'th column replaced by b .

2.2 Recurrences

If $a_n=c_1a_{n-1}+\cdots+c_ka_{n-k}$, and r_1,\ldots,r_k are distinct roots of $x^k+c_1x^{k-1}+\cdots+c_k$, there are d_1,\ldots,d_k s.t.

$$a_n=d_1r_1^n+\cdots+d_kr_k^n.$$

Non-distinct roots r become polynomial factors, e.g. $a_n=(d_1n+d_2)r^n$.

2.3 Trigonometry

$$\begin{aligned}\sin(v+w)&=\sin v\cos w+\cos v\sin w \\ \cos(v+w)&=\cos v\cos w-\sin v\sin w\end{aligned}$$

$$\begin{aligned}\tan(v+w)&=\frac{\tan v+\tan w}{1-\tan v\tan w} \\ \sin v+\sin w&=2\sin\frac{v+w}{2}\cos\frac{v-w}{2} \\ \cos v+\cos w&=2\cos\frac{v+w}{2}\cos\frac{v-w}{2}\end{aligned}$$

$$(V+W)\tan(v-w)/2=(V-W)\tan(v+w)/2$$

where V,W are lengths of sides opposite angles v,w .

$$\begin{aligned}a\cos x+b\sin x&=r\cos(x-\phi) \\ a\sin x+b\cos x&=r\sin(x+\phi)\end{aligned}$$

where $r=\sqrt{a^2+b^2},\phi=\operatorname{atan2}(b,a)$.

2.4 Geometry

2.4.1 Triangles

Side lengths: a,b,c

Semiperimeter: $p=\frac{a+b+c}{2}$

Area: $A=\sqrt{p(p-a)(p-b)(p-c)}$

Circumradius: $R=\frac{abc}{4A}$

Inradius: $r=\frac{A}{p}$

Length of median (divides triangle into two equal-area triangles): $m_a=\frac{1}{2}\sqrt{2b^2+2c^2-a^2}$

Length of bisector (divides angles in two):

$$s_a=\sqrt{bc\left[1-\left(\frac{a}{b+c}\right)^2\right]}$$

Law of sines: $\frac{\sin\alpha}{a}=\frac{\sin\beta}{b}=\frac{\sin\gamma}{c}=\frac{1}{2R}$

Law of cosines: $a^2=b^2+c^2-2bc\cos\alpha$

Law of tangents: $\frac{a+b}{a-b}=\frac{\tan\frac{\alpha+\beta}{2}}{\tan\frac{\alpha-\beta}{2}}$

2.4.2 Quadrilaterals

With side lengths a,b,c,d , diagonals e,f , diagonals angle θ , area A and magic flux $F=b^2+d^2-a^2-c^2$:

$$4A=2ef\cdot\sin\theta=F\tan\theta=\sqrt{4e^2f^2-F^2}$$

For cyclic quadrilaterals the sum of opposite angles is 180° , $ef=ac+bd$, and $A=\sqrt{(p-a)(p-b)(p-c)(p-d)}$.

2.4.3 Spherical coordinates

$$\begin{aligned}x&=r\sin\theta\cos\phi & r&=\sqrt{x^2+y^2+z^2} \\ y&=r\sin\theta\sin\phi & \theta&=\operatorname{acos}(z/\sqrt{x^2+y^2+z^2}) \\ z&=r\cos\theta & \phi&=\operatorname{atan2}(y,x)\end{aligned}$$

2.5 Derivatives/Integrals

$$\begin{aligned}\frac{d}{dx}\arcsin x&=\frac{1}{\sqrt{1-x^2}} & \frac{d}{dx}\arccos x&=-\frac{1}{\sqrt{1-x^2}} \\ \frac{d}{dx}\tan x&=1+\tan^2x & \frac{d}{dx}\arctan x&=\frac{1}{1+x^2} \\ \int\tan ax&=-\frac{\ln|\cos ax|}{a} & \int x\sin ax&=\frac{\sin ax-ax\cos ax}{a^2} \\ \int e^{-x^2}&=\frac{\sqrt{\pi}}{2}\operatorname{erf}(x) & \int xe^{ax}dx&=\frac{e^{ax}}{a^2}(ax-1)\end{aligned}$$

Integration by parts:

$$\int_a^bf(x)g(x)dx=[F(x)g(x)]_a^b-\int_a^bF(x)g'(x)dx$$

2.6 Sums

$$c^a+c^{a+1}+\cdots+c^b=\frac{c^{b+1}-c^a}{c-1},c\neq1$$

$$1+2+3+\cdots+n=\frac{n(n+1)}{2}$$

$$1^2+2^2+3^2+\cdots+n^2=\frac{n(2n+1)(n+1)}{6}$$

$$1^3+2^3+3^3+\cdots+n^3=\frac{n^2(n+1)^2}{4}$$

$$1^4+2^4+3^4+\cdots+n^4=\frac{n(n+1)(2n+1)(3n^2+3n-1)}{30}$$

2.7 Series

$$e^x=1+x+\frac{x^2}{2!}+\frac{x^3}{3!}+\ldots,(-\infty<x<\infty)$$

$$\ln(1+x)=x-\frac{x^2}{2}+\frac{x^3}{3}-\frac{x^4}{4}+\ldots,(-1<x\leq1)$$

$$\begin{aligned}\sqrt{1+x} &= 1 + \frac{x}{2} - \frac{x^2}{8} + \frac{2x^3}{32} - \frac{5x^4}{128} + \dots, \quad (-1 \leq x \leq 1) \\ \sin x &= x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots, \quad (-\infty < x < \infty) \\ \cos x &= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots, \quad (-\infty < x < \infty)\end{aligned}$$

2.8 Probability theory

Let X be a discrete random variable with probability $p_X(x)$ of assuming the value x . It will then have an expected value (mean) $\mu = \mathbb{E}(X) = \sum_x x p_X(x)$ and variance $\sigma^2 = V(X) = \mathbb{E}(X^2) - (\mathbb{E}(X))^2 = \sum_x (x - \mathbb{E}(X))^2 p_X(x)$ where σ is the standard deviation. If X is instead continuous it will have a probability density function $f_X(x)$ and the sums above will instead be integrals with $p_X(x)$ replaced by $f_X(x)$.

Expectation is linear:

$$\mathbb{E}(aX + bY) = a\mathbb{E}(X) + b\mathbb{E}(Y)$$

For independent X and Y ,

$$V(aX + bY) = a^2V(X) + b^2V(Y).$$

2.8.1 Discrete distributions

Binomial distribution

The number of successes in n independent yes/no experiments, each which yields success with probability p is $\text{Bin}(n, p)$, $n = 1, 2, \dots$, $0 \leq p \leq 1$.

$$\begin{aligned}p(k) &= \binom{n}{k} p^k (1-p)^{n-k} \\ \mu &= np, \sigma^2 = np(1-p)\end{aligned}$$

$\text{Bin}(n, p)$ is approximately $\text{Po}(np)$ for small p .

First success distribution

The number of trials needed to get the first success in independent yes/no experiments, each wich yields success with probability p is $\text{Fs}(p)$, $0 \leq p \leq 1$.

$$\begin{aligned}p(k) &= p(1-p)^{k-1}, \quad k = 1, 2, \dots \\ \mu &= \frac{1}{p}, \sigma^2 = \frac{1-p}{p^2}\end{aligned}$$

Poisson distribution

The number of events occurring in a fixed period of time t if these events occur with a known average rate κ and independently of the time since the last event is $\text{Po}(\lambda)$, $\lambda = t\kappa$.

$$\begin{aligned}p(k) &= e^{-\lambda} \frac{\lambda^k}{k!}, \quad k = 0, 1, 2, \dots \\ \mu &= \lambda, \sigma^2 = \lambda\end{aligned}$$

2.8.2 Continuous distributions

Uniform distribution

If the probability density function is constant between a and b and 0 elsewhere it is $\text{U}(a, b)$, $a < b$.

$$\begin{aligned}f(x) &= \begin{cases} \frac{1}{b-a} & a < x < b \\ 0 & \text{otherwise} \end{cases} \\ \mu &= \frac{a+b}{2}, \sigma^2 = \frac{(b-a)^2}{12}\end{aligned}$$

Exponential distribution

The time between events in a Poisson process is $\text{Exp}(\lambda)$, $\lambda > 0$.

$$\begin{aligned}f(x) &= \begin{cases} \lambda e^{-\lambda x} & x \geq 0 \\ 0 & x < 0 \end{cases} \\ \mu &= \frac{1}{\lambda}, \sigma^2 = \frac{1}{\lambda^2}\end{aligned}$$

Normal distribution

Most real random values with mean μ and variance σ^2 are well described by $\mathcal{N}(\mu, \sigma^2)$, $\sigma > 0$.

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

If $X_1 \sim \mathcal{N}(\mu_1, \sigma_1^2)$ and $X_2 \sim \mathcal{N}(\mu_2, \sigma_2^2)$ then

$$aX_1 + bX_2 + c \sim \mathcal{N}(\mu_1 + \mu_2 + c, a^2\sigma_1^2 + b^2\sigma_2^2)$$

2.9 Markov chains

A *Markov chain* is a discrete random process with the property that the next state depends only on the current state. Let X_1, X_2, \dots be a sequence of random variables generated by the Markov process. Then there is a transition matrix $\mathbf{P} = (p_{ij})$, with $p_{ij} = \Pr(X_n = i | X_{n-1} = j)$, and $\mathbf{p}^{(n)} = \mathbf{P}^n \mathbf{p}^{(0)}$ is the probability distribution for X_n (i.e., $p_i^{(n)} = \Pr(X_n = i)$), where $\mathbf{p}^{(0)}$ is the initial distribution.

π is a stationary distribution if $\pi = \pi \mathbf{P}$. If the Markov chain is *irreducible* (it is possible to get to any state from any state), then $\pi_i = \frac{1}{\mathbb{E}(T_i)}$ where $\mathbb{E}(T_i)$ is the expected time between two visits in state i . π_j/π_i is the expected number of visits in state j between two visits in state i .

For a connected, undirected and non-bipartite graph, where the transition probability is uniform among all neighbors, π_i is proportional to node i 's degree.

A Markov chain is *ergodic* if the asymptotic distribution is independent of the initial distribution. A finite Markov chain is ergodic iff it is irreducible and *aperiodic* (i.e., the gcd of cycle lengths is 1). $\lim_{k \rightarrow \infty} \mathbf{P}^k = \mathbf{1}\pi$.

A Markov chain is an A-chain if the states can be partitioned into two sets \mathbf{A} and \mathbf{G} , such that all states in \mathbf{A} are absorbing ($p_{ii} = 1$), and all states in \mathbf{G} leads to an absorbing state in \mathbf{A} . The probability for absorption in state $i \in \mathbf{A}$, when the initial state is j , is $a_{ij} = p_{ij} + \sum_{k \in \mathbf{G}} a_{ik} p_{kj}$. The expected time until absorption, when the initial state is i , is $t_i = 1 + \sum_{k \in \mathbf{G}} p_{ki} t_k$.

Data structures (3)

OrderStatisticTree.h

HashMap.h

UnionFindRollback.h

SubMatrix.h

Matrix.h

LineContainer.h

Treap.h

RMQ.h

MoQueries.h

Numerical (4)

4.1 Polynomials and recurrences

Polynomial.h

PolyRoots.h

PolyInterpolate.h

BerlekampMassey.h

LinearRecurrence.h

4.2 Optimization

GoldenSectionSearch.h

HillClimbing.h

Integrate.h

IntegrateAdaptive.h

Simplex.h

Duality.h

4.3 Matrices

Determinant.h

IntDeterminant.h

SolveLinear.h

SolveLinear2.h

SolveLinearBinary.h

MatrixInverse.h

Tridiagonal.h

4.4 Fourier transforms

FastFourierTransform.h

FastFourierTransformMod.h

NumberTheoreticTransform.h

FastSubsetTransform.h

Number theory (5)

5.1 Modular arithmetic

ModularArithmetic.h

ModInverse.h

ModPow.h

ModLog.h

ModSum.h

ModMulLL.h

ModSqrt.h

5.2 Primality

FastEratosthenes.h

MillerRabin.h

Factor.h

5.3 Divisibility

euclid.h

CRT.h

5.3.1 Bézout’s identity

For $a \neq 0, b \neq 0$, then $d = gcd(a,b)$ is the smallest positive integer for which there are integer solutions to

$$ax + by = d$$

If (x,y) is one solution, then all solutions are given by

$$\left(x + \frac{kb}{gcd(a,b)}, y - \frac{ka}{gcd(a,b)}\right), \quad k \in \mathbb{Z}$$

phiFunction.h

5.4 Fractions

ContinuedFractions.h

FracBinarySearch.h

5.5 Pythagorean Triples

The Pythagorean triples are uniquely generated by

$$a = k \cdot (m^2 - n^2), \quad b = k \cdot (2mn), \quad c = k \cdot (m^2 + n^2),$$

with $m > n > 0, k > 0, m \perp n$, and either m or n even.

5.6 Primes

$p = 962592769$ is such that $2^{21} \mid p - 1$, which may be useful. For hashing use 970592641 (31-bit number), 31443539979727 (45-bit), 3006703054056749 (52-bit). There are 78498 primes less than 1 000 000.

Primitive roots exist modulo any prime power p^a , except for $p = 2, a > 2$, and there are $\phi(\phi(p^a))$ many. For $p = 2, a > 2$, the group $\mathbb{Z}_{2^a}^\times$ is instead isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_{2^{a-2}}$.

5.7 Estimates

$$\sum_{d \mid n} d = O(n \log \log n).$$

The number of divisors of n is at most around 100 for $n < 5e4$, 500 for $n < 1e7$, 2000 for $n < 1e10$, 200 000 for $n < 1e19$.

5.8 Mobius Function

$$\mu(n) = \begin{cases} 0 & n \text{ is not square free} \\ 1 & n \text{ has even number of prime factors} \\ -1 & n \text{ has odd number of prime factors} \end{cases}$$

Mobius Inversion:

$$g(n) = \sum_{d \mid n} f(d) \Leftrightarrow f(n) = \sum_{d \mid n} \mu(d) g(n/d)$$

Other useful formulas/forms:

$$\sum_{d \mid n} \mu(d) = [n = 1] \text{ (very useful)}$$

$$g(n) = \sum_{n \mid d} f(d) \Leftrightarrow f(n) = \sum_{n \mid d} \mu(d/n) g(d)$$

$$g(n) = \sum_{1 \leq m \leq n} f(\lfloor \frac{n}{m} \rfloor) \Leftrightarrow f(n) = \sum_{1 \leq m \leq n} \mu(m) g(\lfloor \frac{n}{m} \rfloor)$$

Combinatorial (6)

6.1 Permutations

6.1.1 Factorial

n	1	2	3	4	5	6	7	8	9	10
$n!$	1	2	6	24	120	720	5040	40320	362880	3628800
n	11	12	13	14	15	16	17			
$n!$	4.0e7	4.8e8	6.2e9	8.7e10	1.3e12	2.1e13	3.6e14			
n	20	25	30	40	50	100	150	171		
$n!$	2e18	2e25	3e32	8e47	3e64	9e157	6e262	>DBL.MAX		

IntPerm.h

6.1.2 Cycles

Let $g_S(n)$ be the number of n -permutations whose cycle lengths all belong to the set S . Then

$$\sum_{n=0}^\infty g_S(n) \frac{x^n}{n!} = \exp \left(\sum_{n \in S} \frac{x^n}{n} \right)$$

6.1.3 Derangements

Permutations of a set such that none of the elements appear in their original position.

$$D(n) = (n-1)(D(n-1) + D(n-2)) = nD(n-1) + (-1)^n = \left\lfloor \frac{n!}{e} \right\rfloor$$

6.1.4 Burnside’s lemma

Given a group G of symmetries and a set X , the number of elements of X up to *symmetry* equals

$$\frac{1}{|G|} \sum_{g \in G} |X^g|,$$

where X^g are the elements fixed by g ($g.x = x$).

If $f(n)$ counts “configurations” (of some sort) of length n , we can ignore rotational symmetry using $G = \mathbb{Z}_n$ to get

$$g(n) = \frac{1}{n} \sum_{k=0}^{n-1} f(\gcd(n,k)) = \frac{1}{n} \sum_{k \mid n} f(k) \phi(n/k).$$

6.2 Partitions and subsets

6.2.1 Partition function

Number of ways of writing n as a sum of positive integers, disregarding the order of the summands.

$$p(0) = 1, \quad p(n) = \sum_{k \in \mathbb{Z} \setminus \{0\}} (-1)^{k+1} p(n - k(3k-1)/2)$$

$$p(n) \sim 0.145/n \cdot \exp(2.56\sqrt{n})$$

n	0	1	2	3	4	5	6	7	8	9	20	50	100
$p(n)$	1	1	2	3	5	7	11	15	22	30	627	$\sim 2e5$	$\sim 2e8$

6.2.2 Lucas’ Theorem

Let n, m be non-negative integers and p a prime. Write $n = n_kp^k + \dots + n_1p + n_0$ and $m = m_kp^k + \dots + m_1p + m_0$. Then $\binom{n}{m} \equiv \prod_{i=0}^k \binom{n_i}{m_i} \pmod{p}$.

6.2.3 Binomials

multinomial.h

6.3 General purpose numbers

6.3.1 Bernoulli numbers

EGF of Bernoulli numbers is $B(t) = \frac{t}{e^t - 1}$ (FFT-able).
 $B[0, \dots] = [1, -\frac{1}{2}, \frac{1}{6}, 0, -\frac{1}{30}, 0, \frac{1}{42}, \dots]$

Sums of powers:

$$\sum_{i=1}^n n^m = \frac{1}{m+1} \sum_{k=0}^m \binom{m+1}{k} B_k \cdot (n+1)^{m+1-k}$$

Euler-Maclaurin formula for infinite sums:

$$\begin{aligned} \sum_{i=m}^\infty f(i) &= \int_m^\infty f(x)dx - \sum_{k=1}^\infty \frac{B_k}{k!} f^{(k-1)}(m) \\ &\approx \int_m^\infty f(x)dx + \frac{f(m)}{2} - \frac{f'(m)}{12} + \frac{f'''(m)}{720} + O(f^{(5)}(m)) \end{aligned}$$

6.3.2 Stirling numbers of the first kind

Number of permutations on n items with k cycles.

$$\begin{aligned} c(n, k) &= c(n-1, k-1) + (n-1)c(n-1, k), \quad c(0, 0) = 1 \\ \sum_{k=0}^n c(n, k)x^k &= x(x+1)\dots(x+n-1) \end{aligned}$$

$$\begin{aligned} c(8, k) &= 8, 0, 5040, 13068, 13132, 6769, 1960, 322, 28, 1 \\ c(n, 2) &= 0, 0, 1, 3, 11, 50, 274, 1764, 13068, 109584, \dots \end{aligned}$$

6.3.3 Eulerian numbers

Number of permutations $\pi \in S_n$ in which exactly k elements are greater than the previous element. k j :s s.t. $\pi(j) > \pi(j+1)$, $k+1$ j :s s.t. $\pi(j) \geq j$, k j :s s.t. $\pi(j) > j$.

$$E(n, k) = (n-k)E(n-1, k-1) + (k+1)E(n-1, k)$$

$$E(n, 0) = E(n, n-1) = 1$$

$$E(n, k) = \sum_{j=0}^k (-1)^j \binom{n+1}{j} (k+1-j)^n$$

6.3.4 Stirling numbers of the second kind

Partitions of n distinct elements into exactly k groups.

$$S(n, k) = S(n-1, k-1) + kS(n-1, k)$$

$$S(n, 1) = S(n, n) = 1$$

$$S(n, k) = \frac{1}{k!} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} j^n$$

6.3.5 Bell numbers

Total number of partitions of n distinct elements. $B(n) = 1, 1, 2, 5, 15, 52, 203, 877, 4140, 21147, \dots$ For p prime,

$$B(p^m + n) \equiv mB(n) + B(n+1) \pmod{p}$$

6.3.6 Labeled unrooted trees

on n vertices: n^{n-2}
on k existing trees of size n_i : $n_1n_2\dots n_kn^{k-2}$
with degrees d_i : $(n-2)!/((d_1-1)!\dots(d_n-1)!)$

6.3.7 Catalan numbers

$$C_n = \frac{1}{n+1} \binom{2n}{n} = \binom{2n}{n} - \binom{2n}{n+1} = \frac{(2n)!}{(n+1)n!}$$

$$C_0 = 1, \quad C_{n+1} = \frac{2(2n+1)}{n+2} C_n, \quad C_{n+1} = \sum C_i C_{n-i}$$

$$C_n = 1, 1, 2, 5, 14, 42, 132, 429, 1430, 4862, 16796, 58786, \dots$$

- sub-diagonal monotone paths in an $n \times n$ grid.
- strings with n pairs of parenthesis, correctly nested.
- binary trees with with $n+1$ leaves (0 or 2 children).
- ordered trees with $n+1$ vertices.
- ways a convex polygon with $n+2$ sides can be cut into triangles by connecting vertices with straight lines.
- permutations of $[n]$ with no 3-term increasing subseq.

Graph (7)

7.1 Fundamentals

BellmanFord.h

FloydWarshall.h

TopoSort.h

7.2 Network flow

PushRelabel.h

MinCostMaxFlow.h

EdmondsKarp.h

Dinic.h

MinCut.h

GlobalMinCut.h

GomoryHu.h

7.3 Matching

hopcroftKarp.h

DFSMatching.h

MinimumVertexCover.h

WeightedMatching.h

GeneralMatching.h

7.4 DFS algorithms

SCC.h

BiconnectedComponents.h

2sat.h

EulerWalk.h

7.5 Coloring

EdgeColoring.h

7.6 Heuristics

MaximalCliques.h

MaximumClique.h

MaximumIndependentSet.h

7.7 Trees

BinaryLifting.h

LCA.h

CompressTree.h

HLD.h

LinkCutTree.h

DirectedMST.h

7.8 Math

7.8.1 Number of Spanning Trees

Create an $N \times N$ matrix `mat`, and for each edge $a \rightarrow b \in G$, do `mat[a][b]--`, `mat[b][b]++` (and `mat[b][a]--`, `mat[a][a]++` if G is undirected). Remove the i th row and column and take the determinant; this yields the number of directed spanning trees rooted at i (if G is undirected, remove any row/column).

7.8.2 Erdős–Gallai theorem

A simple graph with node degrees $d_1 \geq \dots \geq d_n$ exists iff $d_1 + \dots + d_n$ is even and for every $k = 1 \dots n$,

$$\sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min(d_i, k).$$

Geometry (8)

8.1 Geometric primitives

- Point.h
- lineDistance.h
- SegmentDistance.h
- SegmentIntersection.h
- lineIntersection.h
- sideOf.h
- OnSegment.h
- linearTransformation.h
- Angle.h

8.2 Circles

- CircleIntersection.h
- CircleTangents.h
- CirclePolygonIntersection.h
- circumcircle.h
- MinimumEnclosingCircle.h

8.3 Polygons

- InsidePolygon.h
- PolygonArea.h
- PolygonCenter.h
- PolygonCut.h
- PolygonUnion.h

8.4 Convex Hull

- ConvexHull.h
- HullDiameter.h
- PointInsideHull.h
- LineHullIntersection.h

- minkowskiSum.h
- halfPlaneInter.h

8.5 Misc. Point Set Problems

- ClosestPair.h
- ManhattanMST.h
- kdTree.h
- FastDelaunay.h

8.6 3D

- PolyhedronVolume.h
- Point3D.h
- 3dHull.h
- sphericalDistance.h

Strings (9)

- KMP.h
- Zfunc.h
- Manacher.h
- MinRotation.h
- SuffixArray.h
- SuffixAutomaton.h
- AhoCorasick.h

Various (10)

10.1 Intervals

- IntervalContainer.h
- IntervalCover.h
- ConstantIntervals.h

10.2 Misc. algorithms

- TernarySearch.h
- LIS.h
- FastKnapsack.h

10.3 Dynamic programming

- KnuthDP.h
- DivideAndConquerDP.h

10.4 Debugging tricks

- `signal(SIGSEGV, [](int) { _Exit(0); });` converts segfaults into Wrong Answers. Similarly one can catch SIGABRT (assertion failures) and SIGFPE (zero divisions). `_GLIBCXX_DEBUG` failures generate SIGABRT (or SIGSEGV on gcc 5.4.0 apparently).
- `feenableexcept(29);` kills the program on NaNs (1), 0-divs (4), infinities (8) and denormals (16).

10.5 Optimization tricks

`__builtin_ia32_ldmxcsr(40896);` disables denormals (which make floats 20x slower near their minimum value).

10.5.1 Bit hacks

- `x & -x` is the least bit in `x`.
- `for (int x = m; x;) { --x &= m; ... }` loops over all subset masks of `m` (except `m` itself).
- `c = x&-x, r = x+c; (((r^x) >> 2)/c) | r` is the next number after `x` with the same number of bits set.
- `rep(b,0,K) rep(i,0,(1 << K)) if (i & 1 << b) D[i] += D[i^(1 << b)];` computes all sums of subsets.

10.5.2 Pragmas

- `#pragma GCC optimize ("Ofast")` will make GCC auto-vectorize loops and optimizes floating points better.
- `#pragma GCC target ("avx2")` can double performance of vectorized code, but causes crashes on old machines.
- `#pragma GCC optimize ("trapv")` kills the program on integer overflows (but is really slow).

- FastMod.h
- FastInput.h
- BumpAllocator.h
- SmallPtr.h
- BumpAllocatorSTL.h
- SIMD.h

techniques.txt