

Local minimality for the generalized Neumann Hénon problem

Henry Zweiman

September 15, 2026

Abstract

We prove strict local minimality, modulo scaling, of radial minimizers for the generalized Neumann Hénon quotient when the exponent of the radial weight is large. The parameter range is determined by the first nonlinear Steklov eigenvalue. It contains a right neighborhood of the homogeneity exponent for every $2 < p < n$, and the entire subcritical range when $n \geq 4$ and $p > 2$ is sufficiently close to 2. This settles the local-minimality question left open after the second-variation analysis of Nazarov and Shcheglova. The proof combines a common weighted space, compact boundary traces, and a nonlinear coercivity lemma that retains the energy of concentrating perturbations. We also give an analytic proof that the admissible interval is nonempty. Above the Steklov threshold, a first angular mode yields instability; the threshold itself is not decided.

1. Introduction and main result

Let B be the unit ball in $\mathbb{R}^n$, let $\Sigma = \partial B$, and write $S = |\Sigma|$. All function spaces are real. For

$$n \geq 3, \quad 2 < p < n, \quad p < q < p^* := \frac{np}{n-p},$$

consider

$$\mathcal{Q}_\alpha(u) = \frac{\int_B (|\nabla u|^p + |u|^p) dx}{\left(\int_B |x|^\alpha |u|^q dx \right)^{p/q}}, \quad u \in W^{1,p}(B) \setminus \{0\}, \quad \alpha > 0. \quad (1)$$

A positive critical point, after multiplication by a constant, solves

$$-\Delta_p u + u^{p-1} = |x|^\alpha u^{q-1} \quad \text{in } B, \quad \partial_\nu u = 0 \quad \text{on } \Sigma. \quad (2)$$

We study positive minimizers of (1) among radial functions. Their existence and limiting profile are established in [Sh18]. The issue is whether these radial minimizers also minimize locally against arbitrary, possibly nonradial, perturbations.

Define the first nonlinear Steklov eigenvalue

$$\lambda = \inf_{\substack{u \in W^{1,p}(B) \\ \int_\Sigma |u|^p > 0}} \frac{\int_B (|\nabla u|^p + |u|^p)}{\int_\Sigma |u|^p}. \quad (3)$$

Its positive eigenfunction ϕ is radial and unique up to scaling. We use the normalization

$$\int_B (|\nabla \phi|^p + \phi^p) = 1, \quad S\lambda\phi(1)^p = 1. \quad (4)$$

Set

$$q_c(n, p) = 1 + \lambda^{-p/(p-1)}(1 - (n-1)\lambda). \quad (5)$$

Theorem 1.1. If $p < q < \min\{p^*, q_c(n, p)\}$, there is α_0 such that, for every $\alpha \geq \alpha_0$, each positive radial minimizer v_α of (1) is a strict $W^{1,p}(B)$ local minimizer modulo its positive scalar ray. More precisely, there is a neighborhood U of v_α such that

$$\mathcal{Q}_\alpha(u) > \mathcal{Q}_\alpha(v_\alpha) \quad \text{for } u \in U \setminus \{tv_\alpha : t > 0\}. \quad (6)$$

If $q_c(n, p) < q < p^*$, every such radial minimizer has a negative nonradial second-variation direction for all sufficiently large α .

The threshold and its angular origin already occur in [NS26, Lemma 1 and equation (25)]. The advance in the positive range is the passage from directional second-variation positivity to a local minimum in the full Sobolev topology. The negative-range assertion is the corresponding angular test, included to identify the scope of the result. We make no assertion when $q = q_c$.

Corollary 1.2. For every $n \geq 3$ and $2 < p < n$, one has $q_c(n, p) > p$. For every $n \geq 4$, there exists $p_0(n) > 2$ such that $q_c(n, p) > p^*$ whenever $2 < p < p_0(n)$. Thus Theorem 1.1 applies to every subcritical $q > p$ in the latter range.

The local-minimum property need not be global. In the interval $p(n-1)/(n-p) < q < p^*$, the global minimizer is nonradial for large α [Sh18, Theorem 3.1]. In dimensions at least four and for p close to two, this gives radial strict local minima coexisting with lower nonradial states.

For $p = 2$, local minimality was proved by Gazzini and Serra [GS08, Proposition 3.6 and Theorem 3.8]. Nazarov and Shcheglova [NS26] obtain directional positivity for $p > 2$ and explicitly leave local minimality open in the introduction. Their latest listed arXiv version on September 15, 2026 is v2, dated May 11, 2026. Our proof uses the radial convergence from [Sh18], but does not infer Sobolev coercivity from directional positivity.

Weighted completions and compact embeddings are established tools in the spectral theory of linearized p -Laplace operators [CES11]. The local Morse theory surveyed in [Ci13], and the critical-group estimates in [CDS24], address the substantial difference between Hilbert and Sobolev Banach spaces. The latter work treats Dirichlet problems and compactness for varying regularized solutions. Here a common space controls a degeneracy at the center and a degeneracy at the Neumann boundary. Its compact trace transfers the concentrating potential to a Steklov form. A separate convexity argument then controls perturbations whose gradients concentrate near the degenerate set. The non-degeneracy theorem of Asselle, Cingolani, and Starostka [ACS25] assumes a positive regularization parameter in its PDE application. Its abstract criterion requires a nearby directional Hessian estimate, which is not supplied by positivity at the critical point. We claim neither the introduction of weighted spectral spaces nor a general Morse theory.

2. From a weighted gap to a local minimum

This section isolates the nonlinear argument. For a bounded smooth domain $\Omega \subset \mathbb{R}^n$, use the norm

$$\|h\|_{1,p}^p = \int_{\Omega} (|\nabla h|^p + |h|^p).$$

Lemma 2.1 (nonlinear coercivity). Suppose $2 < p < q < p^*$, $0 \leq b \in L^\infty(\Omega)$, and $b \not\equiv 0$. Let $v \in C^1(\overline{\Omega})$ be a positive critical point of

$$\frac{pF(u)}{D(u)^{p/q}}, \quad F(u) = \frac{1}{p}\|u\|_{1,p}^p, \quad D(u) = \int_{\Omega} b|u|^q,$$

normalized by $D(v) = 1$. Assume $\inf_{\Omega} v > 0$. Put $\Lambda = pF(v)$,

$$\ell(h) = \int_{\Omega} b v^{q-1} h, \quad a = |\nabla v|^{p-2},$$

and define the Hessian form

$$A(h) = \int_{\Omega} \left[a |\nabla h|^2 + (p-2) |\nabla v|^{p-4} (\nabla v \cdot \nabla h)^2 + (p-1) v^{p-2} h^2 \right]. \quad (7)$$

The middle integrand is zero where $\nabla v = 0$. Suppose that $a > 0$ almost everywhere, $a^{-1} \in L^1(\Omega)$, and the completion H of $C^\infty(\overline{\Omega})$ in the norm $A^{1/2}$ embeds compactly into $L^2(\Omega)$. If, for some $\kappa > 0$,

$$A(h) - \Lambda(q-1) \int_{\Omega} b v^{q-2} h^2 \geq \kappa A(h) \quad (h \in H, \ell(h) = 0), \quad (8)$$

then there exist $c, \rho > 0$ such that

$$F(v+h) - F(v)D(v+h)^{p/q} \geq c(A(h) + \|h\|_{1,p}^p) \quad (9)$$

whenever $h \in W^{1,p}(\Omega)$, $\ell(h) = 0$, and $\|h\|_{1,p} < \rho$.

Proof. Boundedness of v and ∇v gives a continuous inclusion $W^{1,p} \subset H$. The reciprocal-weight assumption identifies the weighted gradients as distributional gradients: weighted Cauchy--Schwarz bounds their L^1 norm. The value term in (7) controls L^2 .

Consider any sequence of nonzero tangent perturbations h_j with $\varepsilon_j = \|h_j\|_{1,p} \rightarrow 0$. Write $A(h, k)$ for the polarization of A , and set

$$t_j^2 = A(h_j) + \varepsilon_j^p, \quad z_j = h_j/t_j. \quad (10)$$

Then $\varepsilon_j^{p/2} \leq t_j \leq C\varepsilon_j$. On a subsequence,

$$\begin{aligned} z_j &\rightharpoonup z \text{ in } H, & z_j &\rightarrow z \text{ in } L^2, & \ell(z) &= 0, \\ A(z_j) &\rightarrow a_0, & \varepsilon_j^p/t_j^2 &\rightarrow b_0, & a_0 + b_0 &= 1, & A(z) &\leq a_0. \end{aligned} \quad (11)$$

Let $\mathcal{B}_F(u, v) = F(u) - F(v) - F'(v)[u - v]$. We first prove

$$\liminf_j \frac{\mathcal{B}_F(v + h_j, v)}{t_j^2} \geq \frac{1}{2}A(z) + c_p(1 - A(z)) \quad (12)$$

with $c_p > 0$. The second term accounts for energy lost in the weak limit.

Fix $w \in C^\infty(\bar{\Omega})$ and put $k_j = h_j - t_j w$. The exact identity

$$\begin{aligned} \mathcal{B}_F(v + h_j, v) &= \mathcal{B}_F(v + t_j w, v) \\ &\quad + (F'(v + t_j w) - F'(v))[k_j] \\ &\quad + \mathcal{B}_F(v + t_j w + k_j, v + t_j w) \end{aligned} \quad (13)$$

gives the appropriate expansion. Dividing by t_j^2 , the first term tends to $A(w)/2$, and the second tends to $A(w, z - w)$. To justify the latter assertion in H^* , the difference quotient of $|X|^{p-2}X$ at $X = \nabla v$ in the fixed direction ∇w is bounded by $C_w(a + 1)$ for $0 < t \leq 1$. Its square divided by a is bounded by

$$C_w(a + 2 + a^{-1}) \in L^1(\Omega).$$

The pointwise derivative is the matrix in (7). Dominated convergence gives the derivative expansion in the weighted dual norm. The value term is simpler because v is bounded away from zero.

For $f_p(X) = |X|^p/p$, uniform convexity gives

$$f_p(X + Y) - f_p(X) - Df_p(X)Y \geq c_p(|X|^{p-2}|Y|^2 + |Y|^p). \quad (14)$$

For completeness, when $|Y| \leq |X|/2$, the integral Hessian formula supplies the quadratic bound and absorbs $|Y|^p$. After scaling to $|Y| = 1$, strict convexity and compactness supply a positive minimum when $|X| \leq 2$. These two regions cover all pairs. The constant depends only on p .

Apply (14) to both gradient and value terms in the last line of (13). Since

$$|X|^{p-2} \leq C_p(|X + tW|^{p-2} + t^{p-2}|W|^{p-2}),$$

we may replace its shifted quadratic weights by those at v , at a fixed multiplicative cost and an error at most

$$C_w t_j^{p-2} (\varepsilon_j + t_j)^2.$$

After division by t_j^2 , this is $o(1)$. Indeed it is bounded by $C_w \varepsilon_j^{p-2}$ if $p \geq 4$, and by $C_w \varepsilon_j^{(p-2)^2/2}$ if $2 < p < 4$, using the two bounds following (10). The anisotropic Hessian (7) is comparable, with constants depending only on p , to its isotropic weighted part. Moreover,

$$\frac{|||k_j|||_{1,p}^p - \varepsilon_j^p}{t_j^2} \leq C_w \left(\frac{\varepsilon_j^{p-1}}{t_j} + t_j^{p-2} \right) \rightarrow 0. \quad (15)$$

Consequently the lower limit of the last line of (13), divided by t_j^2 , is at least

$$c_p(a_0 - 2A(z, w) + A(w) + b_0).$$

Combine the three terms, then let smooth w converge to z in H . This proves (12). The order of these limits is fixed: first $j \rightarrow \infty$ for each smooth w , then approximation in H .

We next expand the constraint. For any $\eta > 0$, uniform positive upper and lower bounds on v imply

$$\left| |v + s|^q - v^q - qv^{q-1}s - \frac{q(q-1)}{2}v^{q-2}s^2 \right| \leq \eta s^2 + C_\eta |s|^q. \quad (16)$$

This follows from Taylor expansion for small s and the q -power bound away from zero. Sobolev embedding and (10) give

$$\frac{\|h_j\|_q^q}{t_j^2} \leq C\varepsilon_j^{q-p} \longrightarrow 0.$$

Integrate (16) against b , use the tangent constraint, and let $\eta \downarrow 0$. Strong L^2 convergence in (11) yields

$$D(v + h_j) = 1 + \frac{q(q-1)}{2} \int_{\Omega} b v^{q-2} h_j^2 + o(t_j^2). \quad (17)$$

Criticality gives $F'(v)[h_j] = \Lambda \ell(h_j) = 0$. Expanding the scalar power in (17) and using (8), (12), we obtain

$$\begin{aligned} \liminf_j \frac{F(v + h_j) - F(v)D(v + h_j)^{p/q}}{t_j^2} &\geq \frac{1}{2} \left(A(z) - \Lambda(q-1) \int_{\Omega} b v^{q-2} z^2 \right) + c_p(1 - A(z)) \\ &\geq \min\{\kappa/2, c_p\} > 0. \end{aligned} \quad (18)$$

A sequence contradicting (9), with the ratio there tending to zero, is therefore impossible. This proves the lemma. $\square$

The tangent slice suffices for local minimality of the quotient. For u sufficiently close to v , set $\tilde{u} = u/\ell(u)$. Since $\ell(v) = 1$, this factor is positive and tends to one, while $\ell(\tilde{u} - v) = 0$. Homogeneity preserves the quotient, and (9) gives a strict increase unless $\tilde{u} = v$.

3. Radial profiles and a common weight

Normalize radial minimizers by $\|v_{\alpha}\|_{1,p} = 1$. The radial convergence theorem [Sh18, Theorem 2.2] gives

$$v_{\alpha} \longrightarrow \phi \quad \text{in } C(\overline{B}) \text{ and } W^{1,p}(B). \quad (19)$$

These statements apply along every choice of positive normalized radial minimizers. They are C^1 up to the boundary. The Euler equation and its multiplier are

$$-\Delta_p v_{\alpha} + v_{\alpha}^{p-1} = c_{\alpha} r^{\alpha} v_{\alpha}^{q-1}, \quad c_{\alpha} = \left(\int_B r^{\alpha} v_{\alpha}^q \right)^{-1}. \quad (20)$$

The equation holds against all Sobolev tests, by radial averaging of an arbitrary test. Uniform convergence in (19), positivity of ϕ , and concentration of $(\alpha + n)r^{\alpha+n-1} dr$ at 1 imply

$$0 < c \leq v_{\alpha} \leq C, \quad \frac{c_{\alpha}}{\alpha + n} \longrightarrow \frac{1}{S\phi(1)^q}. \quad (21)$$

No boundary expansion is needed to obtain the weight estimates below.

Lemma 3.1. Set $\beta = (p-2)/(p-1) \in (0,1)$. Uniformly over large α and normalized radial minimizers,

$$|v'_\alpha(r)|^{p-2} \asymp r^\beta \min\{1, \alpha(1-r)\}^\beta, \quad 0 < r < 1. \quad (22)$$

In particular $v'_\alpha > 0$ in $(0,1)$.

Proof. Integrating (20) from either endpoint gives the two exact formulas

$$\begin{aligned} r^{n-1}|v'_\alpha|^{p-2}v'_\alpha &= \int_0^r s^{n-1}v_\alpha^{p-1} ds - c_\alpha \int_0^r s^{\alpha+n-1}v_\alpha^{q-1} ds \\ &= c_\alpha \int_r^1 s^{\alpha+n-1}v_\alpha^{q-1} ds - \int_r^1 s^{n-1}v_\alpha^{p-1} ds. \end{aligned} \quad (23)$$

For fixed small $\delta > 0$ and $r \leq 1 - \delta$, the first formula is bounded above by Cr^n and below by $cr^n - Cr^{\alpha+n}$. For large α , it is therefore comparable to r^n .

On $1 - \delta \leq r < 1$, the second formula is bounded below by

$$c(1 - r^{\alpha+n}) - C(1 - r).$$

Here $1 - r^{\alpha+n} \asymp \min\{1, \alpha(1-r)\}$. Choose δ small and then α large. The negative term can be absorbed, both when $\alpha(1-r) \leq 1$ and when it exceeds 1. Dropping that term gives the matching upper bound. Divide by r^{n-1} and raise to the power β . The two regions yield (22). $\square$

Define

$$\omega(r) = r^\beta(1-r)^\beta, \quad E(h) = \int_B (\omega|\nabla h|^2 + h^2), \quad (24)$$

and let H_ω be the completion of $C^\infty(\overline{B})$ in this norm. If $a_\alpha = |v'_\alpha|^{p-2}$, then (22) gives

$$c\omega \leq a_\alpha \leq C, \quad a_\alpha \leq C_\alpha \omega. \quad (25)$$

The first two constants are uniform for large α ; the last may depend on α . Also $\omega^{-1} \in L^1(B)$. Formula (23) and (19) imply convergence of v'_α to ϕ' uniformly on every closed subinterval of $(0,1)$.

4. Compact traces and boundary concentration

Lemma 4.1. The maps $H_\omega \rightarrow L^2(B)$ and $T : H_\omega \rightarrow L^2(\Sigma)$ are compact, where T extends the ordinary boundary trace of smooth functions. If $h_j \rightarrow h$ in H_ω and $\alpha_j \rightarrow \infty$, then

$$(\alpha_j + n) \int_B r^{\alpha_j} v_{\alpha_j}^{q-2} h_j^2 \longrightarrow \phi(1)^{q-2} \int_\Sigma (Th)^2. \quad (26)$$

The corresponding linear limit is

$$(\alpha_j + n) \int_B r^{\alpha_j} v_{\alpha_j}^{q-1} h_j \longrightarrow \phi(1)^{q-1} \int_\Sigma Th. \quad (27)$$

Proof. On every annulus separated from 0 and 1, the norm $E^{1/2}$ controls the ordinary H^1 norm. For $1/2 \leq r < 1$, radial Cauchy-Schwarz gives

$$\|h(1, \cdot) - h(r, \cdot)\|_{L^2(\Sigma)}^2 \leq C(1-r)^{1-\beta} E(h). \quad (28)$$

An average over a fixed interior annulus bounds the trace itself by $CE(h)^{1/2}$, so it extends by completion. Averages over an annulus at distance δ from Σ are compact in $L^2(\Sigma)$ by interior Rellich compactness. By (28), they approximate the trace in operator norm with error $O(\delta^{(1-\beta)/2})$. Thus T is compact. The same estimate gives

$$\int_{1-\delta < |x| < 1} h^2 \leq C\delta E(h). \quad (29)$$

At the center, comparison along rays with the interior trace at $r = 1/2$ uses

$$\int_r^{1/2} \frac{ds}{s^{n-1}\omega(s)} \leq Cr^{2-n-\beta}.$$

The interior trace is bounded by $CE(h)^{1/2}$. Integrating the ray estimate therefore gives

$$\int_{|x| < \delta} h^2 \leq C(\delta^n + \delta^{2-\beta})E(h). \quad (30)$$

Equations (29), (30), and Rellich compactness on interior annuli prove compactness in $L^2(B)$.

For (26), first replace $h_j(r, \cdot)$ by Th_j on $r \geq 1/2$. Both have uniformly bounded $L^2(\Sigma)$ norm there. By (28) the error in their squared norms is at most $C(1-r)^{(1-\beta)/2}$. Its integral against the probability density $(\alpha_j + n)r^{\alpha_j+n-1}$ is $O(\alpha_j^{-(1-\beta)/2})$. The contribution from $r < 1/2$ is exponentially small by the interior L^2 bound. Uniform convergence in (19) and continuity of ϕ replace $v_{\alpha_j}^{q-2}$ by $\phi(1)^{q-2}$, with an error tending to zero. Finally compactness gives $Th_j \rightarrow Th$ in $L^2(\Sigma)$. This proves (26). The same argument, without squaring h_j , proves (27). $\square$

5. The limiting angular inequality and the weighted gap

For a radial profile v_α , write its Hessian energy as

$$A_\alpha(h) = \int_B \left[a_\alpha((p-1)|\partial_r h|^2 + r^{-2}|\nabla_\theta h|^2) + (p-1)v_\alpha^{p-2}h^2 \right]. \quad (31)$$

Let A_∞ denote the same expression with ϕ in place of v_α . In particular $A_\alpha \geq cE$, uniformly for large α , and A_α is equivalent to E for every fixed such α .

Lemma 5.1. If g has zero spherical average at almost every radius and finite A_∞ energy, then

$$A_\infty(g) \geq L \int_\Sigma (Tg)^2, \quad L = \phi(1)^{p-2} \left[\lambda^{-1/(p-1)} - (n-1)\lambda^\beta \right]. \quad (32)$$

Proof. This is the angular Steklov calculation in [NS26, Lemma 1]. We give a verification in the finite-energy class needed here. The radial equation is

$$(r^{n-1}(\phi')^{p-1})' = r^{n-1}\phi^{p-1}, \quad \phi'(1) = \lambda^{1/(p-1)}\phi(1). \quad (33)$$

Set $f = \phi' > 0$. Differentiation on $(0, 1)$ yields

$$-((p-1)r^{n-1}(\phi')^{p-2}f')' + \left[(n-1)r^{n-3}(\phi')^{p-2} + (p-1)r^{n-1}\phi^{p-2} \right] f = 0. \quad (34)$$

Apply the spherical Poincaré inequality with eigenvalue $n-1$, and complete the radial square with g/f on an annulus. The resulting lower bound is the radial boundary expression

$$\left[(p-1)r^{n-1}(\phi')^{p-2} \frac{f'}{f} \|g(r, \cdot)\|_{L^2(\Sigma)}^2 \right]_{r=\varepsilon}^{r=1}. \quad (35)$$

Indeed the discarded term is the integral of $(p-1)r^{n-1}(\phi')^{p-2}f^2|\partial_r(g/f)|^2$.

At the origin, (33) implies $\phi'(r) \sim \phi(0)(r/n)^{1/(p-1)}$ and $f'/f \sim 1/((p-1)r)$. Finite angular energy gives

$$\int_0^{1/2} r^{n-3+\beta} \|g(r, \cdot)\|_2^2 dr < \infty.$$

Hence the lower boundary term in (35), of order $r^{n-2+\beta}\|g(r, \cdot)\|_2^2$, tends to zero along a sequence $r \downarrow 0$. At 1, (33) makes its coefficient equal to L . Apply the square identity first on interior annuli, then pass to the boundary trace and to that sequence at the center. Ordinary Sobolev approximation on annuli justifies the argument for finite-energy g . This proves (32). $\square$

Define the tangent functional and the constrained Hessian by

$$\begin{aligned} \ell_\alpha(h) &= \int_B r^\alpha v_\alpha^{q-1} h, \\ Q_\alpha(h) &= A_\alpha(h) - (q-1)c_\alpha \int_B r^\alpha v_\alpha^{q-2} h^2. \end{aligned} \quad (36)$$

Both extend continuously to H_ω for each fixed α .

Proposition 5.2. If $q < q_c(n, p)$, there exist $\delta > 0$ and α_0 such that

$$Q_\alpha(h) \geq \delta E(h) \quad \text{whenever } \alpha \geq \alpha_0, \ h \in H_\omega, \ \ell_\alpha(h) = 0. \quad (37)$$

The constants apply to every positive normalized radial minimizer.

Proof. Otherwise choose $\alpha_j \rightarrow \infty$ and tangent h_j with $E(h_j) = 1$ and $Q_{\alpha_j}(h_j) \leq 1/j$. By weak compactness and Lemma 4.1, after a subsequence $h_j \rightharpoonup h$ in H_ω , with strong convergence in both L^2 spaces. Since $A_{\alpha_j}(h_j) \geq c > 0$, (21) and (26) imply

$$\int_\Sigma (Th)^2 > 0. \quad (38)$$

The linear limit (27) and the tangent constraint give $\int_\Sigma Th = 0$.

On each interior annulus the gradient coefficients converge uniformly and are uniformly positive. Weak lower semicontinuity there, exhaustion, and strong L^2 convergence for the value term give

$$\begin{aligned} 0 &\geq \liminf_j Q_{\alpha_j}(h_j) \\ &\geq A_\infty(h) - (q-1)\lambda\phi(1)^{p-2} \int_\Sigma (Th)^2. \end{aligned} \quad (39)$$

Here we used $1/(S\phi(1)^2) = \lambda\phi(1)^{p-2}$ from (4). The energy $A_\infty(h)$ is finite; otherwise (39) is already a contradiction.

Decompose $h = h_0 + g$ into its radial average and a function of zero spherical average. The energy splits orthogonally. The trace of h_0 is zero, so $Tg = Th$. Discard $A_\infty(h_0) \geq 0$ and apply (32). The last line of (39) is at least

$$\lambda\phi(1)^{p-2}(q_c - q) \int_\Sigma (Th)^2 > 0,$$

contradicting (38)–(39). $\square$

6. Proof of the main theorem and the parameter range

Fix α as in Proposition 5.2. By (25), (31), and (37), there is $\kappa_\alpha > 0$ such that $Q_\alpha \geq \kappa_\alpha A_\alpha$ on $\ker \ell_\alpha$. Multiply v_α by a positive constant so that $\int_B r^\alpha v_\alpha^q = 1$. Both the Hessian energy and its negative potential scale by the same power $p - 2$, so the gap persists. The weighted completion is still H_ω , with an equivalent norm. Its embedding into L^2 is compact. Equation (22) gives reciprocal-gradient-weight integrability, and (21) supplies strict positivity. Thus Lemma 2.1 applies with $b = r^\alpha$. Its coercivity and the scalar retraction following the lemma prove (6).

The neighborhood and nonlinear coercivity constant may depend on α and on the minimizer. Proposition 5.2 provides a uniform weighted quadratic gap, but we do not claim a uniform $W^{1,p}$ neighborhood.

For the negative-range assertion, let Y be a first spherical harmonic and set $g(r, \theta) = \phi'(r)Y(\theta)$. Equality holds in (32). This function has finite A_∞ energy, since its gradient energy near zero is of order

$$\int_0^\varepsilon r^{n-1-\beta} dr < \infty.$$

Cut it off smoothly for $r < \varepsilon$, using a cutoff equal to one for $r > 2\varepsilon$. The cutoff error in A_∞ is $O(\varepsilon^{n-\beta})$, and its boundary trace is unchanged. This produces smooth functions g_ε on $\overline{B}$, each with zero spherical average. If $q > q_c$, choose ε so small that

$$A_\infty(g_\varepsilon) - (q - 1)\lambda\phi(1)^{p-2} \int_\Sigma g_\varepsilon^2 < 0. \quad (40)$$

For this fixed smooth test, (21), (25), interior convergence of the gradient coefficients, and dominated convergence give $Q_\alpha(g_\varepsilon) < 0$ for large α . The test is tangent for every radial v_α . Since the quotient is C^2 on $W^{1,p} \setminus \{0\}$ for the exponents under consideration, its second derivative in this direction is a positive multiple of $Q_\alpha(g_\varepsilon)$. Local minimality is impossible. This proves Theorem 1.1.

To prove Corollary 1.2, introduce

$$z(r) = \left(\frac{\phi'(r)}{\phi(r)} \right)^{p-1}.$$

Equation (33) gives

$$z' = 1 - \frac{n-1}{r}z - (p-1)z^{p/(p-1)}, \quad z(r) = r/n + o(r), \quad z'(r) \rightarrow 1/n. \quad (41)$$

Differentiating for $r > 0$ yields

$$z'' + \left(\frac{n-1}{r} + pz^{1/(p-1)} \right) z' = \frac{n-1}{r^2} z > 0. \quad (42)$$

An integrating factor starting at a sufficiently small positive radius shows that $z' > 0$ throughout $(0, 1]$. Since $z(1) = \lambda$, (41) implies

$$1 - (n-1)\lambda - (p-1)\lambda^{p/(p-1)} > 0,$$

which is exactly $q_c > p$. This provides a direct analytic proof of the positivity underlying [NS26, Lemmas 6–7].

Also, integrating (33) and using strict increase of ϕ gives

$$z(r) = \frac{r^{1-n}}{\phi(r)^{p-1}} \int_0^r s^{n-1} \phi(s)^{p-1} ds < \frac{r}{n}.$$

Thus $\lambda < 1/n$, and (5) implies

$$q_c > 1 + n^{1/(p-1)}. \quad (43)$$

For $n \geq 4$, one has $1+n > 2n/(n-2)$. Continuity of the explicit functions in (43) and $p^* = np/(n-p)$ shows that $1 + n^{1/(p-1)} > p^*$ for p in a right neighborhood of 2. This proves Corollary 1.2. $\square$

7. Scope and further questions

The equality case $q = q_c(n, p)$ requires higher-order analysis or more precise dependence on α . The present limiting quadratic form has a first-angular-mode zero direction there, so Proposition 5.2 does not decide that case. A uniform nonlinear neighborhood as $\alpha \rightarrow \infty$ is also beyond the result: the norm-equivalence constant in (25) can grow with α .

Lemma 2.1 is formulated independently of radial geometry. It may be useful when a degenerate variational problem supplies a compact weighted completion and a strictly positive constrained Hessian. Its hypotheses are part of the conclusion's scope: pointwise positivity of second variations alone is not asserted to suffice. The common weight in this application has integrable reciprocal and admits a compact boundary trace; problems with stronger boundary degeneracy require a different argument.

References

- [ACS25] L. Asselle, S. Cingolani, and M. Starostka, *Morse homology for a class of elliptic partial differential equations*, arXiv:2504.19721v1 (2025), Theorems 1.2 and 3.7. [Author preprint](#); [related journal DOI](#).
- [CDS24] S. Cingolani, M. Degiovanni, and B. Sciunzi, *Weighted Sobolev spaces and Morse estimates for quasilinear elliptic equations*, Journal of Functional Analysis **286** (2024), no. 8, 110346. [DOI](#).
- [CES11] D. Castorina, P. Esposito, and B. Sciunzi, *Spectral theory for linearized p -Laplace equations*, Nonlinear Analysis **74** (2011), 3606–3613. [DOI](#).
- [Ci13] S. Cingolani, *On local Morse theory for p -area functionals, $p > 2$* , Journal of Fixed Point Theory and Applications **14** (2013), 355–373; published online April 7, 2014. [DOI](#).
- [GS08] M. Gazzini and E. Serra, *The Neumann problem for the Hénon equation, trace inequalities and Steklov eigenvalues*, Annales de l'Institut Henri Poincaré C, Analyse non linéaire **25** (2008), 281–302. [DOI](#).
- [NS26] A. I. Nazarov and A. P. Shcheglova, *The Neumann problem for the generalized Hénon equation. Local analysis*, arXiv:2604.26286v2, May 11, 2026. [Author preprint](#).
- [Sh18] A. P. Shcheglova, *The Neumann problem for the generalized Hénon equation*, Journal of Mathematical Sciences **235** (2018), no. 3, 360–373. [DOI](#).