

Sharp logarithmic concavity on balls and its instability under convex deformation

Henry Zweiman

September 16, 2026

Abstract

We study positive Dirichlet solutions of $-\Delta u = u \log u^2$ on a ball in any dimension. We prove that $-\log(u(r)/u(0))$ has a power series in r^2 with strictly positive coefficients and convergence radius equal to the square of the ball radius. The coefficients follow an explicit positive recurrence and are polynomials with nonnegative coefficients in the excess of the central logarithmic height above the Gaussian threshold. This structure proves concavity of $-\sqrt{-\log(u/\|u\|_\infty)}$, answering a question of Gallo, Mosconi and Squassina. The exponent $1/2$ is sharp among positive powers of the negative normalized logarithm. We also obtain a quantitative lower bound for its Hessian before taking the square root, positive power concavity of u on every ball, and a comparison of normalized profiles. The argument gives an alternative proof of the known shooting order and ball uniqueness theorem. The essential step is to pass from a local positive recurrence to a convergent representation throughout the domain. We also show that the normalized square-root-log conclusion is unstable under smooth uniformly convex domain deformations. In every dimension at least two and near every ball, an explicit degree-three boundary deformation produces positive solutions that remain strictly log-concave but fail the stronger transformed concavity arbitrarily close to their unique maximum. A cubic-jet obstruction and an invertible ball linearization give a hand proof, including the boundary nonlinearity in the shape argument.

Mathematics Subject Classification (2020). 35B45, 35J61, 34B15, 26A51.

Keywords. Logarithmic Schrödinger equation; concavity; radial solutions; positive power series; shooting method.

1. Introduction and main results

Let $N \geq 1$, $R > 0$, and $B_R = \{x \in \mathbb{R}^N : |x| < R\}$. We consider

$$-\Delta u = u \log u^2 \quad \text{in } B_R, \quad u > 0 \quad \text{in } B_R, \quad u = 0 \quad \text{on } \partial B_R, \quad (1.1)$$

where $u \in C^2(B_R) \cap C(\overline{B_R})$. All concavity statements about a logarithmic transform concern the open ball. Statements about u^α extend to the closed ball by continuity.

Gallo, Mosconi and Squassina [GMS] establish existence of a solution with locally negative-definite logarithmic Hessian on every bounded convex domain. Their Theorem 6.5 gives radial symmetry and radial decrease of every positive solution on a ball. In Sections 6.2–6.3 they ask for stronger concavity in

balls: positive power concavity of u , and concavity of the negative square root of its negative normalized logarithm. Their Theorem 6.6 proves both properties in dimension one. The second question remains explicitly stated on page 44 of the final article.

Our main result is a positive-series representation that answers both questions in every dimension. Write $u(x) = U(r)$, $r = |x|$, and set

$$M = \log U(0), \quad w(r) = -\log \frac{U(r)}{U(0)}, \quad A = \frac{2M}{N}. \quad (1.2)$$

Theorem 1.1 (Positive radial series). Every solution of (1.1) satisfies $A > 1$. There are strictly positive coefficients c_n , $n \geq 0$, such that

$$\frac{w'(r)}{r} = \sum_{n=0}^{\infty} c_n r^{2n}, \quad w(r) = \sum_{j=1}^{\infty} \frac{c_{j-1}}{2j} r^{2j}, \quad 0 \leq r < R. \quad (1.3)$$

The quotient at $r = 0$ is interpreted by continuity. Both series, regarded as series in r^2 , have radius of convergence exactly R^2 . Their coefficients are determined by

$$c_0 = A, \quad c_{n+1} = \frac{(n+1) \sum_{i=0}^n c_i c_{n-i} - c_n}{(n+1)(2n+N+2)}. \quad (1.4)$$

In particular, $Z(t) = w'(\sqrt{t})/\sqrt{t}$ is absolutely monotone on $[0, R^2]$: every derivative is strictly positive there. The same is true of every derivative of positive order of $W(t) = w(\sqrt{t})$. In addition,

$$D^2 w(|x|) \succeq A I_N, \quad (1.5)$$

with strict inequality as quadratic forms when $x \neq 0$. Here $\succeq$ denotes the order on symmetric matrices.

Theorem 1.2 (Sharp transformed concavity). For every solution of (1.1),

$$x \mapsto - \left(-\log \frac{u(x)}{\|u\|_{\infty}} \right)^{\beta} \text{ is concave on } B_R \iff \beta \geq \frac{1}{2}, \quad \beta > 0. \quad (1.6)$$

For $\beta = 1/2$ its Hessian is negative definite away from the center. At the center it is continuous and has a norm-type cusp; no differentiability assertion there is intended.

Theorem 1.3 (Positive power concavity). There exists a number $\alpha_*(U) \in (0, 1)$ such that, for $\alpha > 0$,

$$u^{\alpha} \text{ is concave on } \overline{B_R} \iff \alpha \leq \alpha_*(U). \quad (1.7)$$

It is characterized by an attained interior minimum,

$$\alpha_*(U) = \min_{0 < r < R} \frac{w''(r)}{w'(r)^2}. \quad (1.8)$$

The formula is implicit. We do not assert a positive exponent uniform over all radii, or monotonicity of this optimal exponent as a function of the radius.

Theorem 1.4 (Instability under convex deformation). Let $N \geq 2$, $R > 0$, and let U be the positive Dirichlet solution on B_R . Define $Y_3(\theta) = \theta_1^3 - 3\theta_1\theta_2^2$ and

$$\Omega_\varepsilon = \{r\theta : 0 \leq r < R + \varepsilon Y_3(\theta), \theta \in S^{N-1}\}.$$

For all sufficiently small $|\varepsilon|$, these are smooth uniformly convex domains and there is a smooth branch of positive solutions u_ε , with $u_0 = U$. For each sufficiently small nonzero ε , the solution is strictly log-concave and has a unique maximum, but

$$-\sqrt{-\log(u_\varepsilon/\|u_\varepsilon\|_\infty)}$$

is not concave in any neighborhood of that maximum. The domains converge to B_R in every C^k boundary norm.

The normalization in this statement is essential. The scale-invariant notion of 1/2-log-concavity permits a sufficiently small multiplicative factor in front of u ; see [IST, Section 4.2]. Theorem 1.4 concerns the specific factor $1/\|u\|_\infty$, where the transform has its cusp at the maximum. It does not disprove the weaker rescaled property. It answers the controlled-deformation question posed in the earlier version of Section 7 and leaves the ball theorem unchanged.

1.1. Prior results and the scope of the contribution

The amplitude threshold, uniqueness in balls, and strict ordering of first zeros are prior results. For $k = 1$ and $\beta = 0$, Ben Chrouda [BC, Theorem 1.2 and Corollaries 1.3 and 1.5] proves these facts for $-\Delta v = v \log v$. The change $v(y) = u(y/\sqrt{2})$ gives the normalization (1.1). The equality case in the amplitude threshold is excluded by the Gaussian solution and uniqueness of the central initial-value problem. We give new proofs of these particular consequences because they fit the recurrence argument, but do not claim their first discovery.

Liu, Sun and Zou [LSZ] prove a broader shooting classification of whole-space nodal solutions; in particular, their argument establishes strict motion of zeros with the initial value. Gallo, Moraschi and Squassina [GMoS] develop parabolic concavity principles whose logarithmic example yields preservation of logarithmic concavity. Neither of these results supplies the positive radial series used here. Our priority claim concerns that structure and the resulting stronger concavity, rather than existence, radial symmetry, Gaussian ground-state classification, or shooting uniqueness. Current versions of these sources were compared on September 15, 2026.

The proof of Theorem 1.1 has three stages. An integrating factor puts the logarithmic radial derivative above its Gaussian value. Its equation then produces positive Taylor coefficients. Finally, positivity and a real-axis singularity argument show that the series represents the solution all the way to its first zero. Convexity of a Hilbert-space norm proves Theorem 1.2. A separate boundary argument, which accommodates the singular derivative of $u \log u^2$ at zero, completes Theorem 1.3. For Theorem 1.4, the ball Hessian bound and the classical weighted Poincaré inequality make the linearized Dirichlet operator invertible. A degree-three shape derivative then violates a necessary cubic-jet condition at the moving maximum.

2. The radial equation and the Gaussian threshold

We use radial symmetry and radial decrease from [GMS, Theorem 6.5]. These are nontrivial inputs: the derivative of the nonlinearity is unbounded below near zero, so an unqualified appeal to the usual locally Lipschitz moving-plane theorem would be inappropriate. Existence follows from [GMS, Theorem 1.1], with its one-dimensional result covering $N = 1$. The rest of the argument applies to every positive radial decreasing solution.

Put $f(s) = s \log s^2$ for $s > 0$, and $f(0) = 0$. The radial equation and its integral form are

$$U'' + \frac{N-1}{r}U' + f(U) = 0, \quad U'(r) = -r^{1-N} \int_0^r s^{N-1} f(U(s)) ds. \quad (2.1)$$

The central initial-value problem is unique as long as its solution stays positive. Indeed, integrating (2.1) once more gives a Volterra equation. On a sufficiently small interval the Lipschitz bound for f near a prescribed positive central value makes its integral operator a contraction, since the norm of the double integral is bounded by a constant times r^2 . Ordinary uniqueness then continues the solution away from zero. The same identity, using smoothness of f near a positive central value, gives the usual even expansions at the center.

Since U has its maximum at zero, (2.1) implies $f(U(0)) \geq 0$, hence $M \geq 0$. If $M = 0$, central uniqueness would give $U \equiv 1$, which violates the boundary condition. Thus $M > 0$.

In fact $U'(r) < 0$ for $0 < r < R$. If U' vanished there, its nonpositivity would imply $U'' = 0$ at that point. Equation (2.1) would give $U = 1$, and ordinary uniqueness with data $(U, U') = (1, 0)$ would again force the constant solution.

In terms of (1.2), equation (2.1) becomes

$$w'' + \frac{N-1}{r}w' - (w')^2 = 2M - 2w. \quad (2.2)$$

Define $z(r) = w'(r)/r$ for $r > 0$ and extend it to the center. The central expansion gives

$$z(0) = \frac{2M}{N} = A, \quad z'(r) = O(r). \quad (2.3)$$

Differentiation of (2.2) yields

$$z'' + \left(\frac{N+1}{r} - 2rz \right) z' = 2z(z-1). \quad (2.4)$$

Equivalently,

$$\left(r^{N+1} e^{-2w} z' \right)' = 2r^{N+1} e^{-2w} z(z-1). \quad (2.5)$$

Lemma 2.1. We have $A > 1$ and $z'(r) > 0$ for all $0 < r < R$.

Proof. Suppose first that $0 < A < 1$. Integrating (2.5) from zero shows that $z' < 0$ while $0 < z < 1$. The boundary term at zero vanishes by (2.3). The function cannot first reach 1, since it is decreasing. It cannot first reach zero either: the integral in (2.5) would then give a strictly negative derivative, whereas $z = w'/r > 0$ throughout the interior. Thus $0 < z \leq A$ on $[0, R)$, giving $w(r) \leq Ar^2/2$. This contradicts $w(r) \rightarrow \infty$ as $r \uparrow R$.

If $A = 1$, the central solution is

$$U(r) = \exp\left(\frac{N}{2} - \frac{r^2}{2}\right), \quad (2.6)$$

by the uniqueness following (2.1). It has no finite zero. Hence $A > 1$. Now (2.5) gives $z' > 0$ as long as $z > 1$, and this strict increase prevents any first exit from that region. $\square$

The eigenvalues of $D^2w(|x|)$ are $w'' = z + rz'$ in the radial direction and z in the tangential directions. Lemma 2.1 proves (1.5), including its strict form away from the center. In dimension one only the radial eigenvalue is present.

3. A positive recurrence valid throughout the ball

Let $t = r^2$ and $Z(t) = z(\sqrt{t})$. Equation (2.4) becomes

$$2tZ'' + (N+2)Z' - 2tZZ' = Z(Z-1). \quad (3.1)$$

Lemma 3.1 (Local construction). For each $A > 1$, recurrence (1.4) defines strictly positive coefficients satisfying

$$0 < c_n \leq A^{n+1}, \quad n \geq 0. \quad (3.2)$$

The resulting series solves (3.1) near zero and agrees with the transformed radial solution.

Proof. The coefficient of t^n in the linear derivative terms of (3.1) is $(n+1)(2n+N+2)c_{n+1}$. In the nonlinear terms, symmetry of convolution gives

$$[t^n](Z^2 + 2tZZ') = (n+1) \sum_{i=0}^n c_i c_{n-i}, \quad (3.3)$$

which proves (1.4). In particular $c_1 = A(A-1)/(N+2) > 0$. For $n \geq 1$, its numerator can be written

$$(2(n+1)A-1)c_n + (n+1) \sum_{i=1}^{n-1} c_i c_{n-i}, \quad (3.4)$$

so positivity follows by induction. Dropping the negative term in (1.4) and using the inductive upper bounds gives

$$c_{n+1} \leq \frac{(n+1)A^{n+2}}{2n+N+2} \leq A^{n+2}. \quad (3.5)$$

Consequently the series converges for $|t| < 1/A$, and termwise differentiation verifies (3.1).

Integrate $w'(r) = rZ(r^2)$ with $w(0) = 0$. The derivative of the difference between the two sides of (2.2) vanishes by (3.1). Its limiting value at zero is $NA - 2M = 0$. Thus (2.2) holds. Setting $U(r) = e^{M-w(r)}$ recovers the radial equation with the prescribed central data. The uniqueness argument after (2.1) identifies this constructed solution with the given one. $\square$

We next state the positive-series continuation principle, including its short proof.

Lemma 3.2 (Positive real singularity). Suppose $F(t) = \sum_{n \geq 0} a_n t^n$ has $a_n \geq 0$ and finite positive radius of convergence ρ . It cannot admit a holomorphic continuation to a neighborhood of the positive real point ρ .

Proof. If such a continuation existed, continuity of its derivatives and monotone convergence as $t \uparrow \rho$ would give, for every $k \geq 0$,

$$F^{(k)}(\rho) = \sum_{n \geq k} \frac{n!}{(n-k)!} a_n \rho^{n-k}. \quad (3.6)$$

Choose a sufficiently small $h > 0$ in the convergence disk of the Taylor series centered at ρ . Its coefficients are nonnegative by (3.6). Interchanging nonnegative sums gives

$$\begin{aligned} \sum_{k \geq 0} \frac{F^{(k)}(\rho)}{k!} h^k &= \sum_{n \geq 0} a_n \sum_{k=0}^n \binom{n}{k} \rho^{n-k} h^k \\ &= \sum_{n \geq 0} a_n (\rho + h)^n < \infty, \end{aligned} \quad (3.7)$$

contradicting the definition of ρ . $\square$

Proof of Theorem 1.1. By Lemma 3.1 the series for Z represents the actual solution near zero. Denote its convergence radius by ρ . If $\rho < R^2$, the actual solution is regular at the positive point ρ . On an interval around that point, (3.1) is an ordinary differential equation with analytic right-hand side after solving for Z'' . It therefore has a locally holomorphic solution with the actual values of Z and Z' there. Ordinary uniqueness identifies this solution with the continuation from zero. This contradicts Lemma 3.2. Thus $\rho \geq R^2$.

Conversely, $z(r)$ tends to infinity as $r \uparrow R$: by Lemma 2.1 it is increasing, and a finite limit would make $w(r) = \int_0^r sz(s) ds$ bounded. Therefore $\rho > R^2$ is impossible. We have proved $\rho = R^2$. Integration gives the second series in (1.3). Dividing its coefficients by a factor linear in the index leaves the convergence radius unchanged. All derivative assertions now follow by differentiating convergent positive series on compact subintervals of $[0, R^2)$. The Hessian assertion was proved after Lemma 2.1. $\square$

The continuation step is essential: positivity of finitely many coefficients, or even a formal positive recurrence without convergence control, would not imply a global concavity theorem.

4. Convexity of the square root and sharpness

Put $a_j = c_{j-1}/(2j) > 0$. Theorem 1.1 gives

$$\sqrt{w(r)} = \left\| (\sqrt{a_j} r^j)_{j \geq 1} \right\|_{\ell^2}, \quad 0 \leq r < R. \quad (4.1)$$

Every coordinate on the right is nonnegative, nondecreasing and convex as a function of r . Coordinatewise convexity, monotonicity of the ℓ^2 norm on nonnegative sequences, and the triangle inequality show that $r \mapsto \sqrt{w(r)}$ is convex and nondecreasing. Equivalently one may first prove this for the finite partial sums and pass to the finite pointwise limit on $[0, R)$.

For strictness it is useful to give an exact identity. At $r > 0$ let

$$p_j = \frac{a_j r^{2j}}{w(r)}, \quad m = \sum_{j \geq 1} j p_j, \quad v = \sum_{j \geq 1} (j - m)^2 p_j. \quad (4.2)$$

These are a probability distribution, its mean and its variance; all moments are finite inside the convergence interval. Direct differentiation gives

$$\frac{r^2(2ww'' - (w')^2)}{4w^2} = 2v + m(m-1) > 0. \quad (4.3)$$

The strict inequality holds because $m > 1$, as $a_2 > 0$. Thus $(\sqrt{w})'' > 0$ for $r > 0$. Its tangential Hessian eigenvalue is $(\sqrt{w})'/r > 0$.

Proof of Theorem 1.2. A convex nondecreasing function of the Euclidean norm is convex. Applying this fact to (4.1) proves convexity of $x \mapsto \sqrt{w(|x|)}$ across the center as well. The Hessian assertions away from the center follow from (4.3). For $\beta \geq 1/2$, the function $s \mapsto s^{2\beta}$ is convex and nondecreasing on $[0, \infty)$; composing it with $\sqrt{w}$ proves the forward implication of (1.6).

For necessity, the central expansion is

$$w(r) = \frac{A}{2}r^2 + O(r^4). \quad (4.4)$$

If $0 < \beta < 1/2$, the restriction $w(r)^\beta$ to a positive radial segment has strictly negative second derivative for sufficiently small $r > 0$. Indeed its leading term is $(A/2)^\beta r^{2\beta}$, whose second derivative has the negative factor $2\beta(2\beta - 1)$; the differentiated remainder has higher order. It cannot be convex. Finally, (4.4) gives $\sqrt{w(|x|)} = \sqrt{A/2}|x| + O(|x|^3)$, which describes the stated cusp. $\square$

The same series gives more than a Hessian bound. For every integer $k \geq 1$, the radial differential operator $\mathcal{D} = (2r)^{-1}d/dr$ satisfies $\mathcal{D}^k w(r) > 0$ on $0 < r < R$, with a positive limit at zero. These infinitely many sign inequalities refer to differentiation with respect to squared radius, not arbitrary mixed spatial derivatives.

5. Boundary behavior and the optimal positive power

Because f is bounded on the compact range of U , (2.1) gives a finite limit of U' at R . The differential equation then gives a continuous limit of U'' there. The next argument rules out a zero boundary slope without assuming a finite value of $f'(0)$.

Lemma 5.1 (Simple boundary zero). There is $b > 0$ such that

$$U(r) = b(R - r) + O((R - r)^2), \quad U'(r) = -b + O(R - r), \quad (5.1)$$

and

$$z(r) \sim \frac{1}{R(R - r)} \quad \text{as } r \uparrow R. \quad (5.2)$$

Proof. Write $v = -U' > 0$. Near R we have $0 < U < 1$, and (2.1) gives $U'' > 0$. Thus v is decreasing there. Set

$$F(s) = \frac{s^2}{2}(\log s^2 - 1), \quad E(r) = \frac{v(r)^2}{2} + F(U(r)). \quad (5.3)$$

The energy identity is $E' = -(N - 1)v^2/r$. Suppose $U'(R) = 0$. Then $E(R) = 0$, and for r sufficiently close to R ,

$$E(r) = \int_r^R \frac{N-1}{s} v(s)^2 ds \leq C(R-r)v(r)^2. \quad (5.4)$$

Choose this neighborhood so that $C(R-r) \leq 1/4$. Equation (5.4) implies

$$\frac{v(r)^2}{4} \leq -F(U(r)), \quad v(r) \leq C_1 U(r) \sqrt{\log(1/U(r))}. \quad (5.5)$$

Since U is strictly decreasing, it can be used as the variable of integration on this interval. The remaining time to reach zero would satisfy

$$R-r = \int_0^{U(r)} \frac{ds}{v(U^{-1}(s))} \geq \frac{1}{C_1} \int_0^{U(r)} \frac{ds}{s \sqrt{\log(1/s)}} = \infty, \quad (5.6)$$

a contradiction. When $N = 1$, the integral in (5.4) is zero and the same argument applies. Thus $b = -U'(R) > 0$. The bounded second derivative gives (5.1), and $z = -U'/(rU)$ gives (5.2). $\square$

Proof of Theorem 1.3. For $r > 0$,

$$(U^\alpha)'' = \alpha U^\alpha (\alpha (w')^2 - w''), \quad \frac{(U^\alpha)'}{r} = -\alpha U^\alpha \frac{w'}{r} < 0. \quad (5.7)$$

Thus the radial Hessian condition is precisely $\alpha \leq q(r)$, where $q = w''/(w')^2$. The condition at the center holds automatically, because $D^2 u^\alpha(0) = -\alpha U(0)^\alpha A I_N$. By Lemma 2.1, $q > 0$. Equation (4.4) gives $q(r) \rightarrow \infty$ at zero, while Lemma 5.1 gives

$$q(r) = 1 - \frac{U(r)U''(r)}{U'(r)^2} \rightarrow 1 \quad \text{as } r \uparrow R. \quad (5.8)$$

Moreover $U'' > 0$ near R , so $q < 1$ there. Choose any such point. The limits at both endpoints and continuity imply that the global infimum is achieved on a compact interior interval; its value is strictly between zero and one. Equations (5.7) prove (1.7)–(1.8), and continuity extends concavity to the closed ball. $\square$

6. Coefficient comparison and two consequences

The recurrence contains information about variation with the central height. Fix N and put $\delta = A - 1 > 0$. Write $c_n(\delta)$ for its coefficients.

Proposition 6.1 (Positive parameter dependence). For every $n \geq 1$, $c_n(\delta)$ is a polynomial in δ with nonnegative coefficients and zero constant term. If two solutions have $0 < \delta_1 < \delta_2$ and $\theta = \delta_2/\delta_1$, then on their common radial interval,

$$z_2(r) - 1 \geq \theta(z_1(r) - 1), \quad (6.1)$$

and

$$w_2(r) - \frac{r^2}{2} \geq \theta \left(w_1(r) - \frac{r^2}{2} \right). \quad (6.2)$$

Proof. We have $c_1 = (\delta + \delta^2)/(N+2)$. For $n \geq 1$, rewrite (1.4) as

$$c_{n+1} = \frac{(2n+1+2(n+1)\delta)c_n + (n+1)\sum_{i=1}^{n-1} c_i c_{n-i}}{(n+1)(2n+N+2)}. \quad (6.3)$$

The polynomial assertion follows by induction. Every monomial of positive degree satisfies $\delta_2^k \geq \theta \delta_1^k$. Apply this coefficientwise, including the constant-in- r term $c_0 - 1 = \delta$, and sum the convergent series to obtain (6.1). Integrating $r(z_i(r) - 1)$ gives (6.2). $\square$

Corollary 6.2 (Known shooting order and uniqueness). If $U_2(0) > U_1(0)$, their first zero radii satisfy $R_2 < R_1$. Consequently (1.1) has exactly one positive solution for every $R > 0$.

Proof. Suppose $R_2 \geq R_1$. By (6.1) and divergence of z_1 at R_1 , strict inequality $R_2 > R_1$ is impossible. If $R_2 = R_1$, (5.2) gives $z_2/z_1 \rightarrow 1$, contradicting (6.1) and $\theta > 1$. This proves the strict shooting order. Two distinct positive solutions on the same ball are radial and have either different central values, contradicted by this order, or equal central values, contradicted by initial-value uniqueness. Existence is the prior input stated in Section 2. $\square$

As emphasized in Section 1.1, this is an alternative proof of a prior theorem. The coefficientwise parameter comparison (6.1)–(6.2) is the additional conclusion of the present method.

Corollary 6.3 (Products of balls). Let u_i be positive solutions of (1.1) on $B_{R_i} \subset \mathbb{R}^{N_i}$, for $1 \leq i \leq k$. Then

$$u(x_1, \dots, x_k) = \prod_{i=1}^k u_i(x_i) \quad (6.4)$$

solves the same equation on the product domain, and $-\sqrt{-\log(u/\|u\|_\infty)}$ is concave there.

Proof. Dividing the Laplacian of the product by the product gives $-\Delta u/u = \sum_i \log u_i^2 = \log u^2$. The negative normalized logarithm is $\sum_i w_i$. Its square root is the Euclidean norm of the vector $(\sqrt{w_i})_{i=1}^k$. Each component is nonnegative and convex in its corresponding variables by Theorem 1.2. The argument of (4.1) proves convexity of this norm. $\square$

This includes the interval-product construction already treated in [GMS]; the factorization itself is elementary and is not a separate novelty claim.

7. Instability under convex deformation

We prove Theorem 1.4. All boundary-distance and function-space statements below refer to a fixed smooth ball before pulling back the domain.

7.1. A necessary cubic-jet condition

Let $w \in C^4$ near x_0 , with $w(x_0) = 0$, $Dw(x_0) = 0$ and positive-definite Hessian $H = D^2w(x_0)$. If $\sqrt{w}$ is convex in a neighborhood of x_0 , then

$$D^3w(x_0) = 0. \quad (D1)$$

Indeed, on any unit direction e , write

$$w(x_0 + te) = at^2 + bt^3 + O(t^4), \quad a = \frac{1}{2}H[e, e] > 0, \quad b = \frac{1}{6}D^3w(x_0)[e, e, e].$$

For $t > 0$, the second derivative of the square root tends to $b/\sqrt{a}$; for $t < 0$ it tends to $-b/\sqrt{a}$. To justify differentiating the remainder, write $w(x_0 + te) = t^2 h(t)$ with h twice continuously differentiable and $h(0) = a$, $h'(0) = b$. Convexity on both half intervals forces $b = 0$. A symmetric cubic tensor whose diagonal polynomial vanishes is zero. Conversely, if the tensor is nonzero, one half of a line has strictly negative square-root second derivative at points arbitrarily close to x_0 . The vanishing condition is necessary, not sufficient.

7.2. The ball linearization is invertible

Use Theorem 1.1 and its estimate

$$-D^2 \log U \succeq AI_N, \quad A = \frac{2 \log U(0)}{N} > 1. \quad (\text{D2})$$

The linearized Dirichlet operator is

$$L = -\Delta - (2 \log U + 2). \quad (\text{D3})$$

It satisfies $LU = -2U$. Since $U > 0$, -2 is its simple first eigenvalue. For every $\phi \in H_0^1(B_R)$ perpendicular to U in L^2 ,

$$\langle L\phi, \phi \rangle \geq (2A - 2) \|\phi\|_2^2. \quad (\text{D4})$$

Here U has a strictly negative outward derivative and U is comparable to the boundary distance d . Consequently the logarithmic potential is form-bounded with relative bound zero on H_0^1 : use $|\log d| \leq \eta d^{-2} + C_\eta$ and Hardy's inequality. The operator has compact resolvent and its form domain is H_0^1 .

For a compactly supported smooth ϕ , set $h = \phi/U$. Integration by parts gives the ground-state identity

$$\langle L\phi, \phi \rangle = \int U^2 |Dh|^2 - 2 \int U^2 h^2. \quad (\text{D5})$$

The classical strongly log-concave weighted Poincare inequality (see [KS, Theorem 2.1] for the Brascamp-Lieb form on a convex domain), applied to the density U^2 , gives

$$\int U^2 |h - \bar{h}|^2 \leq \frac{1}{2A} \int U^2 |Dh|^2, \quad \bar{h} = \frac{\int U^2 h}{\int U^2}. \quad (\text{D6})$$

The relevant boundary version can be obtained without any assumption on a positive limiting density at ∂B_R . On each smaller concentric ball, the density is smooth and positive. For its weighted Neumann operator $\mathcal{L} = \Delta + 2D \log U \cdot D$, integration of the Bochner identity for an eigenfunction gives

$$\int (\mathcal{L}h)^2 U^2 = \int (|D^2 h|^2 - 2D^2 \log U [Dh, Dh]) U^2 + \int_{\partial B_r} \text{II}[D_T h, D_T h] U^2. \quad (\text{D7})$$

The outward second fundamental form II of the sphere is positive. Equation (D2) therefore bounds each nonconstant weighted Neumann eigenvalue below by $2A$. The spectral expansion yields (D6) on B_r ; let $r \uparrow R$. First apply this to the smooth compactly supported quotient. Ground-state identity and form density then extend the result to all H_0^1 functions. Perpendicularity to U is exactly $\bar{h} = 0$, proving (D4). Thus L has no zero eigenvalue and has exactly one negative eigenvalue.

For use in a shape argument, fix $0 < \gamma < 1$ and set $X = \{v \in C^{2,\gamma}(\overline{B}_R) : v|_{\partial B_R} = 0\}$, $Y = C^{0,\gamma}(\overline{B}_R)$. Then

$$L : X \longrightarrow Y \text{ is an isomorphism.} \quad (\text{D8})$$

To see the functional-analytic point at the vanishing boundary, choose a smooth positive defining function ρ equal to d near the boundary. Division by ρ maps X continuously to $C^{1,\gamma}$, and $U = \rho a$ with $a > 0$ on the closed ball. Hence multiplication by $\log U$ maps X to Y . It is compact: for $\gamma < \beta < 1$, a bounded sequence in X has a subsequence converging in $C^{1,\beta}$; its quotients by ρ converge in $C^{0,\beta}$, and multiplication by $\rho \log \rho$ maps that convergence to $C^{0,\gamma}$. Thus L is a compact perturbation of the Dirichlet Laplacian isomorphism. Its kernel is zero by (D4) and $LU = -2U$; the Fredholm alternative proves (D8).

7.3. A genuine branch on deformed domains

Let

$$H_3(x) = x_1^3 - 3x_1x_2^2, \quad Y_3 = H_3|_{S^{N-1}},$$

and define

$$\Omega_\varepsilon = \{r\theta : 0 \leq r < R + \varepsilon Y_3(\theta), \theta \in S^{N-1}\}. \quad (\text{D9})$$

For all sufficiently small $|\varepsilon|$, this is a smooth uniformly convex domain. Indeed the radial graph converges to the sphere in C^2 and its principal curvatures converge uniformly to $1/R > 0$; its outward normal parametrization, or the usual support-plane criterion, gives convexity. It converges in every C^k norm because Y_3 is smooth.

Choose a smooth radial cutoff χ zero on $B_{R/2}$ and one near the boundary, and use

$$T_\varepsilon(x) = x + \varepsilon \chi(|x|) Y_3(x/|x|) \frac{x}{|x|}, \quad (\text{D10})$$

with the second term zero near the origin. For small parameter this is a smooth diffeomorphism from the ball to (D9), equal to the identity near zero. The pulled-back Laplacian has coefficients smooth in x and real analytic in ε .

Although $s \log s^2$ is not differentiable at $s = 0$ as a scalar function, the Dirichlet nonlinear map is smooth in the spaces just specified near U . On the open positive-Hopf cone write $v = \rho b$, $b > 0$; then

$$v \log v^2 = 2\rho b \log \rho + 2\rho b \log b. \quad (\text{D11})$$

The first term is a bounded linear map of b into Y , and the second is real analytic for b bounded away from zero in $C^{1,\gamma}$. Division by ρ is bounded on X . Thus this is a real-analytic map from that open cone in X to Y . At parameter zero, its equation has derivative (D8). The Banach implicit-function theorem produces a unique nearby branch $v_\varepsilon \in X$ with $v_0 = U$ satisfying the pulled-back equation. It remains in the positive-Hopf cone. Set $u_\varepsilon = v_\varepsilon \circ T_\varepsilon^{-1}$ on Ω_ε ; these are genuine positive Dirichlet solutions. Interior elliptic regularity gives smooth dependence in every local C^k norm. This does not assert global uniqueness among all positive solutions on the deformed domains.

7.4. The shape derivative has a nonzero cubic term

Let $V = \partial_\varepsilon T_\varepsilon|_0$ and let z be the Eulerian shape derivative, $z = \dot{v}_0 - DU \cdot V$. Differentiation of the equation in each fixed compact subset and of the transported zero boundary value gives

$$Lz = 0 \text{ in } B_R, \quad z|_{\partial B_R} = -U'(R)Y_3. \quad (\text{D12})$$

This is a weak Dirichlet problem with a smooth nonzero boundary trace; z need not be C^2 up to the boundary, since the potential is logarithmic there. It is $C^{1,\gamma}$ up to the boundary and smooth in the interior. Uniqueness follows from (D8), or from the form kernel exclusion. Rotation invariance and spherical harmonic projection therefore give

$$z(r\theta) = \psi(r)Y_3(\theta),$$

where

$$-\psi'' - \frac{N-1}{r}\psi' + \frac{3(N+1)}{r^2}\psi - (2\log U + 2)\psi = 0, \quad \psi(R) = -U'(R) > 0. \quad (\text{D13})$$

Near zero, the regular solutions of this equation form a one-dimensional space and have the expansion

$$\psi(r) = cr^3 + O(r^5). \quad (\text{D14})$$

This follows by inserting $\psi = r^3a(r)$: a satisfies a regular radial integral equation in dimension $N+6$, with smooth even potential near zero. Its initial value $a(0) = c$ determines the solution uniquely. In particular $c \neq 0$, since $c = 0$ would force $\psi = 0$, contradicting its boundary value. In fact $c > 0$: test the radial equation with its negative part times the same degree-three harmonic; (D4) is positive on this angular subspace, so that negative part vanishes, and uniqueness excludes a zero leading value. Thus

$$z(x) = cH_3(x) + O(|x|^5), \quad c > 0. \quad (\text{D15})$$

7.5. Move the maximum before computing its jet

On a small fixed ball at zero, D^2U is negative definite. The nearby solutions have the same property there. Outside a smaller neighborhood of zero, U has a strict value gap below $U(0)$, which persists under the uniform pullback convergence. Hence each nearby u_ε has a unique global maximum x_ε near zero.

The implicit-function theorem applied to the gradient gives smooth x_ε . Since $Dz(0) = 0$, differentiation gives $x'_0 = 0$, and therefore $x_\varepsilon = O(\varepsilon^2)$. Because the base solution is radial, $D^3U(0) = 0$. Interior differentiability and (D15) imply

$$D^3u_\varepsilon(x_\varepsilon) = \varepsilon c D^3H_3 + O(\varepsilon^2). \quad (\text{D16})$$

Set

$$w_\varepsilon(x) = -\log \frac{u_\varepsilon(x)}{u_\varepsilon(x_\varepsilon)}.$$

At the maximum the logarithmic third derivative has no lower-derivative cross terms, since $Du_\varepsilon = 0$. In particular,

$$D^3 w_\varepsilon(x_\varepsilon)[e_1, e_1, e_1] = -\frac{6c}{U(0)}\varepsilon + O(\varepsilon^2) \neq 0 \quad (\text{D17})$$

for every sufficiently small nonzero parameter. Its Hessian at the maximum is positive definite. Apply Section 7.1 to conclude that $\sqrt{w_\varepsilon}$ is not convex in any neighborhood of the maximum. Equivalently, $-\sqrt{w_\varepsilon}$ is not concave.

The expansion centers at the actual maximum, rather than keeping the old center fixed; this is essential for the obstruction to apply.

7.6. Ordinary strict log-concavity persists

This additional statement keeps the two concavity properties distinct. On a fixed interior compact set, (D2) and C^2 convergence preserve positive definiteness of $-D^2 \log u_\varepsilon$. Near the boundary, uniform convexity, uniform Hopf bounds and uniform $C^{2,\gamma}$ bounds supply a collar on which the same is true, uniformly for small parameter. Here is the local estimate.

In normal coordinates $x = y - d\nu(y)$, write $u_\varepsilon = a(y)d + O(d^2)$, where a is uniformly positive. At the boundary, for a unit tangent τ , $D^2 u_\varepsilon[\tau, \tau] = -a \text{II}[\tau, \tau]$. The $C^{2,\gamma}$ bounds and the positive lower curvature bound therefore give, in the tangent/normal splitting at y ,

$$(-D^2 \log u_\varepsilon)_{TT} \succeq \frac{k}{d}I, \quad (-D^2 \log u_\varepsilon)_{NN} \geq \frac{k}{d^2}, \quad |(-D^2 \log u_\varepsilon)_{TN}| \leq \frac{K}{d}, \quad (\text{D18})$$

for fixed positive constants and a sufficiently small common collar. For the first estimate use that the tangential gradient term in $-D^2 \log u = -D^2 u/u + Du \otimes Du/u^2$ is nonnegative. For the mixed estimate use $|D_T u| = O(d)$ and bounded $D^2 u$; for the normal estimate use the Hopf bound and $u = ad + O(d^2)$. The Schur complement is bounded below by $k/d - K^2/k$, hence is positive for smaller d . Combining collar and interior proves strict log-concavity throughout Ω_ε .

8. Further questions

Theorem 1.4 resolves the earlier question about persistence under arbitrary small smooth uniformly convex deformations: the normalized square-root conclusion fails along the explicit cubic deformation near every ball in dimensions at least two. Classifying the deformations that preserve it remains open. Vanishing of the cubic jet is necessary but is not asserted to be sufficient. We do not classify all positive solutions on the deformed domains.

The positive recurrence still depends on the radial equation and its particular coefficient structure. A classification of nonlinearities or radial potentials admitting a comparable recurrence would be useful. Formula (1.8) does not evaluate the optimal positive concavity exponent. Determining its dependence on the radius in dimensions $N \geq 2$, especially its sharp limiting behavior, remains a further problem. All these questions belong to the same paper.

References

- [BC] M. Ben Chrouda, *Uniqueness and Liouville type results for radial solutions of some classes of k -Hessian equations*, Electronic Journal of Qualitative Theory of Differential Equations **2022**, no. 62, 1–17. [doi:10.14232/ejqtde.2022.1.62](https://doi.org/10.14232/ejqtde.2022.1.62).
- [GMS] M. Gallo, S. Mosconi and M. Squassina, *Power law convergence and concavity for the logarithmic Schrödinger equation*, Mathematische Annalen **395** (2026), article 21, 60 pp. [doi:10.1007/s00208-026-03452-2](https://doi.org/10.1007/s00208-026-03452-2). [arXiv:2411.01614v2](https://arxiv.org/abs/2411.01614v2).
- [GMoS] M. Gallo, R. Moraschi and M. Squassina, *Quantitative and exact concavity principles for parabolic and elliptic equations*, [arXiv:2504.09494v2](https://arxiv.org/abs/2504.09494v2). Also compared with the [41-page author manuscript](#), accessed September 15, 2026.
- [LSZ] T. Liu, X. Sun and W. Zou, *Uniqueness of bound states to the logarithmic Schrödinger equation*, [arXiv:2606.19077v2](https://arxiv.org/abs/2606.19077v2), June 22, 2026.
- [IST] K. Ishige, P. Salani and A. Takatsu, *New characterizations of log-concavity*, [arXiv:2004.13381](https://arxiv.org/abs/2004.13381), Section 4.2. The distinction between fixed normalization and concavity after small rescaling is used here; no heat-flow theorem is a proof dependency.
- [KS] D. J. Knezevic and E. Suli, *Spectral Galerkin approximation of Fokker-Planck equations with unbounded drift*, ESAIM: Mathematical Modelling and Numerical Analysis **43** (2009), 445–485. [Primary journal PDF](#), Theorem 2.1. The classical Brascamp-Lieb inequality is credited; the needed ball boundary version is also proved above.

Preparation note

This manuscript was prepared with OpenAI Codex assistance in literature searches, proof development and writing. It is a research preprint. No independent human verification or journal acceptance is claimed.