

Quasipolynomiality and eventual growth at fixed ordinarization number

Henry Zweiman

September 16, 2026; revised version 1.1

Abstract

Let $n_{g,r}$ count numerical semigroups of genus g and ordinarization number r . For every fixed positive integer r , we prove that $n_{g,r}$ is a quasipolynomial for all positive integers g , answering a question of Cyrusian and Kaplan. Its leading coefficient is $1/(2(2r)!)+1/(2 \cdot 4^r(r!)^2)$. We determine the complete next coefficient in closed form, including its constant part and its oscillation: its exact period is two, and its even value is smaller than its odd value by $1/(4^r r!(r-1)!)$. This gives explicit positive leading terms for $n_{g+1,r} - n_{g,r}$ in both parities and proves eventual strict monotonicity for every fixed r . The proof separates a binomial-sum bulk from exceptional additive relations. Those relations lie in linear hyperplanes through the origin, which prevents them from introducing periodicity in the next coefficient.

1 Introduction and main results

A numerical semigroup is an additive submonoid S of $\mathbb{Z}_{\geq 0}$ with finite complement. Its genus $g(S)$ is the number of gaps, and its Frobenius number $F(S)$ is the largest gap when $g(S) > 0$. The ordinary semigroup of genus g is $\{0, g+1, g+2, \dots\}$. Repeatedly removing the smallest positive nongap and inserting the largest gap leads to this ordinary semigroup. The number of steps is the ordinarization number $r(S)$.

The ordinarization number equals the number of positive elements of S at most $g(S)$; see [2, Proposition 1.4]. We write

$$n_{g,r} = \#\{S : g(S) = g, r(S) = r\}.$$

Cyrusian and Kaplan proved eventual quasipolynomiality of degree $2r$ for fixed r [2, Theorem 3.4]. They asked whether the quasipolynomial holds for every positive genus [2, Remark 3.7]. They also discuss the conjecture $n_{g+1,r} \geq n_{g,r}$ for every g, r [2, Conjecture 1.5], and prove it for $r = 2$. We answer their quasipolynomiality question and prove the eventual strict version of this conjecture for every fixed positive r .

Recall that a quasipolynomial on the positive integers is an expression $\sum_{j=0}^d c_j(g)g^j$ with rational-valued periodic functions c_j . Equivalently, one polynomial applies in each residue class modulo some positive integer. The degree is the largest j for which c_j is not identically zero. All asymptotic estimates below keep r fixed.

Theorem 1.1. *For every positive integer r , the function $g \mapsto n_{g,r}$ is a quasipolynomial of degree $2r$ on all positive integers. Put*

$$A_r = \frac{1}{2(2r)!} + \frac{1}{2 \cdot 4^r(r!)^2}, \quad C_r = \frac{1}{4^r r!(r-1)!}. \quad (1)$$

With the explicit rational number β_r given in Theorem 5.1, we have

$$n_{g,r} = A_r g^{2r} + \left(\beta_r - \frac{C_r}{2} (-1)^g \right) g^{2r-1} + O_r(g^{2r-2}). \quad (2)$$

Consequently, the leading coefficient is constant and the coefficient of g^{2r-1} has exact period two.

Corollary 1.2. For fixed $r \geq 1$,

$$n_{g+1,r} - n_{g,r} = \begin{cases} \left(\frac{r}{(2r)!} + 2C_r \right) g^{2r-1} + O_r(g^{2r-2}), & g \text{ even}, \\ \frac{r}{(2r)!} g^{2r-1} + O_r(g^{2r-2}), & g \text{ odd}. \end{cases} \quad (3)$$

In particular, $n_{g+1,r} > n_{g,r}$ for every sufficiently large g .

The conclusion is pointwise in r : no threshold uniform in a growing ordinarization number is asserted. We do not settle monotonicity at every genus, nor monotonicity of the total number of numerical semigroups. A positive constant leading coefficient alone would not prove Corollary 1.2, because a periodic next coefficient can contribute at the same order to successive differences. Determining that coefficient's oscillation is the essential additional step.

2 An exact homogeneous description

Fix positive integers g, r . Consider integer tuples satisfying

$$0 < b_1 < \dots < b_r \leq g < a_1 < \dots < a_r \leq 2g. \quad (4)$$

Write $B = \{b_1, \dots, b_r\}$ and $A = \{a_1, \dots, a_r\}$, and associate the set

$$S(A, B) = \{0\} \cup B \cup (\{g+1, g+2, \dots\} \setminus A). \quad (5)$$

It has precisely g gaps. We use $2g$ as the upper bound in (4); closure under addition will automatically exclude a gap at $2g$.

Lemma 2.1. The set (5) is a numerical semigroup if and only if both of the following conditions hold:

$$(b_i + b_j > g \text{ or } b_i + b_j \in B) \quad \text{and} \quad b_i + b_j \notin A \quad (1 \leq i \leq j \leq r), \quad (6)$$

$$a_h - b_i \leq g \text{ or } a_h - b_i \in A \quad (1 \leq h, i \leq r). \quad (7)$$

Every numerical semigroup of genus g and ordinarization number r arises from exactly one such tuple.

Proof. Condition (6) says exactly that sums of two elements of B lie in $S(A, B)$. For a mixed sum $b_i + x$ with $x \in S(A, B)$ and $x > g$, the only possible failure is $b_i + x = a_h$ for some h . Such a failure occurs precisely when $a_h - b_i > g$ and $a_h - b_i \notin A$. This is excluded exactly by (7). Two summands greater than g have sum greater than $2g$, and zero causes no obstruction. This proves the equivalence.

For completeness, if S has genus $g > 0$ and Frobenius number F , the map $s \mapsto F - s$ injects $S \cap [0, F]$ into the gaps: otherwise F would be a sum of two elements of S . Hence $F + 1 - g \leq g$, so $F \leq 2g - 1$. There are r positive nongaps at most g , leaving $g - r$ gaps there and therefore r gaps above g . These two sets give B and A uniquely, and the Frobenius bound places A in the required interval. $\square$

We recall the particular lattice-counting facts needed here. A rational polyhedral Boolean set means a set specified by a finite Boolean combination of rational linear equalities and strict or weak inequalities.

Lemma 2.2. *Let $D \subset \mathbb{R}^d$ be a bounded rational polyhedral Boolean set. Then*

$$t \mapsto \#(tD \cap \mathbb{Z}^d)$$

is a quasipolynomial for all positive integers t . If D is contained in a finite union of rational linear hyperplanes through the origin, then there is a rational constant κ such that

$$\#(tD \cap \mathbb{Z}^d) = \kappa t^{d-1} + O_D(t^{d-2}) \quad (8)$$

when $d \geq 2$. The same conclusion holds for a finite signed sum of such counts.

Proof. Take the arrangement of all hyperplanes appearing in a description of D , together with a rational bounding box. Its cells that belong to D are relative interiors of bounded rational polytopes, and they form a finite disjoint partition. Ehrhart's theorem gives quasipolynomial lattice counts for closed rational polytopes [1, Theorem 3.23]. Removing the boundary by inclusion-exclusion over proper faces gives the same conclusion for relative interiors and therefore for D . This argument holds at every positive dilation, including for lower-dimensional polytopes whose affine hulls may meet the lattice only at some dilations.

For the second assertion, refine the arrangement to include the stated hyperplanes. Every selected cell has dimension at most $d - 1$. A cell of dimension $d - 1$ lies in some hyperplane H , and its affine hull equals H . The lattice $\Lambda = H \cap \mathbb{Z}^d$ has full rank in H . After choosing a lattice basis of Λ , the closure P of this cell becomes a full-dimensional rational polytope in $\mathbb{R}^{d-1}$. Its lattice count is

$$\#(t \operatorname{relint} P \cap \Lambda) = \operatorname{vol}_\Lambda(P) t^{d-1} + O_P(t^{d-2}),$$

where volume is normalized so that a fundamental parallelepiped of Λ has volume one. One can see this directly by tiling H by such parallelepipeds: only those meeting the boundary contribute to the discrepancy, and their number is $O_P(t^{d-2})$. Boundary removal has the same error bound. The normalized volume is rational and does not depend on a residue class of t .

Cells of dimension at most $d - 2$ contribute $O_D(t^{d-2})$, by their Ehrhart quasipolynomials. Summing proves (8); signed sums follow by linearity. The requirement that the containing hyperplanes pass through the origin ensures that the full lattice in each top-dimensional affine hull is present at every dilation. $\square$

Proposition 2.3. *For fixed positive r , the function $n_{g,r}$ is a quasipolynomial on all positive integers g , of degree at most $2r$.*

Proof. Apply Lemma 2.1. Membership in a finite set such as B is a finite disjunction of linear equalities, and nonmembership is a conjunction of their negations. All conditions in (4), (6), and (7) are homogeneous in the variables $(a_1, \dots, a_r, b_1, \dots, b_r, g)$. In particular, strict inequalities are retained as strict inequalities, without introducing an additive constant.

Let $D_r \subset \mathbb{R}^{2r}$ be the set obtained by putting $g = 1$ in these conditions. It is bounded and rational polyhedral Boolean. Homogeneity gives

$$n_{g,r} = \#(gD_r \cap \mathbb{Z}^{2r}) \quad (g \geq 1).$$

Lemma 2.2 gives the desired quasipolynomial, with degree at most the ambient dimension. $\square$

3 The bulk and its exceptional hyperplanes

Let $\mathcal{N}_r(g)$ be the set of tuples counted by $n_{g,r}$. Define the bulk $\mathcal{M}_r(g)$ to consist of the integer tuples in (4) satisfying

$$2b_1 > g, \quad a_r \leq g + b_1. \quad (9)$$

These conditions do not themselves impose closure. Put $M_r(g) = \#\mathcal{M}_r(g)$.

Lemma 3.1. *For all positive integers g, r ,*

$$M_r(g) = \sum_{m=\lfloor g/2 \rfloor + 1}^g \binom{g-m}{r-1} \binom{m}{r}, \quad (10)$$

where a binomial coefficient is zero if its lower index exceeds its nonnegative upper index. Moreover, there is $\kappa_r \in \mathbb{Q}$ such that

$$n_{g,r} - M_r(g) = \kappa_r g^{2r-1} + O_r(g^{2r-2}). \quad (11)$$

Proof. Choose $m = b_1$. The remaining $r-1$ elements of B are chosen from $m+1, \dots, g$. The r elements of A are chosen from $g+1, \dots, g+m$. This proves (10).

We claim that the symmetric difference of $\mathcal{N}_r(g)$ and $\mathcal{M}_r(g)$ lies in the union of the proper linear hyperplanes

$$2b_1 = b_i \quad (1 \leq i \leq r), \quad (12)$$

$$a_r - b_1 = a_h \quad (1 \leq h \leq r), \quad (13)$$

$$b_i + b_j = a_h \quad (1 \leq i \leq j \leq r, 1 \leq h \leq r). \quad (14)$$

Indeed, if a tuple is valid but $2b_1 \leq g$, condition (6) forces $2b_1 \in B$. If it is valid but $a_r > g + b_1$, condition (7) forces $a_r - b_1 \in A$. These are (12) and (13).

Conversely, suppose a tuple belongs to the bulk. Every sum of two elements of B is greater than g . Also $a_h - b_i \leq a_r - b_1 \leq g$ for all h, i , so the mixed-sum condition is automatic. The only remaining failures of closure are therefore the equalities in (14). This proves the claim. The argument also holds for the real tuples defined by the same Boolean conditions, because it uses only those conditions and the ordering.

Set $g = 1$ in the real valid and bulk sets. Their two set differences are bounded rational polyhedral Boolean sets in the union of (12)–(14). These hyperplanes involve no g and pass through the origin in $\mathbb{R}^{2r}$. Apply the signed version of Lemma 2.2 with $d = 2r$ to obtain (11). $\square$

Remark 3.2. The constancy of κ_r is crucial. The weaker estimate $n_{g,r} - M_r(g) = O_r(g^{2r-1})$ would permit a correction of the same size as the main term in a successive difference. In contrast, (11) implies

$$(n_{g+1,r} - M_r(g+1)) - (n_{g,r} - M_r(g)) = O_r(g^{2r-2}).$$

No differentiation of an unspecified error term is used: subtracting two quantities individually of order g^{2r-2} remains of that order.

4 Two leading coefficients of the bulk

Lemma 4.1. *There is a rational number γ_r such that*

$$M_r(g) = A_r g^{2r} + \left(\gamma_r - \frac{C_r}{2} (-1)^g \right) g^{2r-1} + O_r(g^{2r-2}), \quad (15)$$

with A_r, C_r as in (1).

Proof. First suppose $r \geq 2$. Expanding the two binomial polynomials gives, uniformly for $g/2 \leq m \leq g$,

$$\binom{g-m}{r-1} \binom{m}{r} = g^{2r-1} f_r(m/g) + g^{2r-2} h_r(m/g) + O_r(g^{2r-3}), \quad (16)$$

where $h_r \in \mathbb{Q}[x]$ is a polynomial and

$$f_r(x) = \frac{x^r(1-x)^{r-1}}{r!(r-1)!}.$$

The polynomial interpretation of the binomial coefficients is valid even when a nonnegative upper index is smaller than the lower index, since the product defining the polynomial then vanishes. There is no endpoint exception in (16).

Write $\delta = 1$ for even g and $\delta = 1/2$ for odd g . The first summation point in (10) is $m = g/2 + \delta$. For any fixed polynomial f , the trapezoidal summation formula gives

$$\sum_{m=\lfloor g/2 \rfloor + 1}^g f(m/g) = g \int_{1/2}^1 f(x) dx + \frac{f(1)}{2} + \left(\frac{1}{2} - \delta\right) f(1/2) + O_f(g^{-1}). \quad (17)$$

To verify the endpoint term, apply the trapezoidal formula on the mesh from $1/2 + \delta/g$ to 1. Extending the integral to $1/2$ subtracts $\delta f(1/2)$, whereas the lower endpoint contributes $f(1/2)/2$, up to $O_f(g^{-1})$.

The h_r sum equals $g \int_{1/2}^1 h_r(x) dx + O_r(1)$. Substitution into (16) shows that the coefficient of g^{2r} is $\int_{1/2}^1 f_r$, and that the only parity dependence in the next coefficient comes from

$$\left(\frac{1}{2} - \delta\right) f_r(1/2).$$

Since $f_r(1/2) = 2C_r$, the even second coefficient is smaller than the odd second coefficient by C_r . Their average is rational.

It remains to evaluate the integral. Put $q(x) = x^{r-1}(1-x)^{r-1}$. Then

$$xq(x) = \frac{1}{2}q(x) + \frac{1}{2}(2x-1)q(x), \quad \frac{d}{dx}(x^r(1-x)^r) = -r(2x-1)q(x).$$

Symmetry of q and the beta integral therefore give

$$\begin{aligned} \int_{1/2}^1 x^r(1-x)^{r-1} dx &= \frac{1}{4} \int_0^1 x^{r-1}(1-x)^{r-1} dx + \frac{1}{2r4^r} \\ &= \frac{((r-1)!)^2}{4(2r-1)!} + \frac{1}{2r4^r}. \end{aligned}$$

Dividing by $r!(r-1)!$ gives A_r .

For $r = 1$ the summand in (10) is exactly m . Direct summation yields

$$M_1(g) = \begin{cases} 3g^2/8 + g/4, & g \text{ even,} \\ 3g^2/8 + g/2 + 1/8, & g \text{ odd.} \end{cases}$$

This has $A_1 = 3/8$, $C_1 = 1/4$, and $\gamma_1 = 3/8$, as required. $\square$

Proof of Theorem 1.1 and Corollary 1.2. Proposition 2.3 gives exact quasipolynomiality for positive g . Combine Lemmas 3.1 and 4.1, and set $\beta_r = \gamma_r + \kappa_r$. This proves (2). Because $A_r > 0$, the degree is $2r$. On each residue class of a common period, a polynomial that is $O(g^{2r-2})$ has zero coefficients in degrees $2r$ and $2r-1$. Thus (2) identifies those coefficients of the exact quasipolynomial. Since $C_r > 0$, the second coefficient has exact period two.

Subtract (2) at consecutive genera. The leading term contributes $2rA_rg^{2r-1} + O_r(g^{2r-2})$, the constant part β_rg^{2r-1} contributes only $O_r(g^{2r-2})$, and the alternating contribution is

$$C_r(-1)^g g^{2r-1} + O_r(g^{2r-2}).$$

Using $2rA_r = r/(2r)! + C_r$ gives (3). Both parity constants are strictly positive, so the error is eventually smaller than the leading term in each parity. $\square$

5 The complete second coefficient

We now evaluate the constant left implicit in the preceding argument. This also answers the question about β_r in the first version of this paper.

Theorem 5.1. *For every $r \geq 1$, the constant in (2) is*

$$\begin{aligned} \beta_r = & -\frac{r^2 + 7r - 10}{8(2r-1)!} - \frac{4r^2 - 4r + 3}{2(2r-1)}C_r \\ & - \frac{3(r-1)}{(2r-1)9^r r!(r-1)!}. \end{aligned} \tag{18}$$

In particular, $\beta_1 = -1/8$, $\beta_2 = -1193/5184$, and $\beta_r < 0$ for every positive r . Moreover,

$$\frac{\beta_r}{A_r} \sim -\frac{r^3}{2} \quad (r \rightarrow \infty). \tag{19}$$

The last statement concerns the coefficients as functions of r . It does not make the error in (2) uniform in r , and does not give a threshold for the counting conjecture.

5.1 The three exceptional contributions

All dimensions in the next argument refer to the $2r$ coordinates (A, B) at fixed real $g > 0$. Enlarge the equality arrangement used in Section 3 to include all equations

$$b_i + b_j = b_k, \quad a_h - b_i = a_j, \quad b_i + b_j = a_h.$$

Delete equations with no solutions in the strict ordering region, and identify equations defining the same hyperplane. Two distinct hyperplanes in this arrangement have an intersection of dimension at most $2r-2$. Their bounded integer sections consequently contribute $O_r(g^{2r-2})$: choose two coordinates with independent coefficients and fix the other $2r-2$ coordinates; at most one solution remains. We may therefore compute the coefficient of g^{2r-1} away from such intersections. Boundary faces of each resulting region do not affect its leading lattice volume.

Suppose first that a valid tuple is outside the bulk because $2m \leq g$, where $m = b_1$. Closure forces $2m \in B$. Away from the discarded intersections this is its only additive equality. Thus $3m > g$, every other element of $B \setminus \{m, 2m\}$ exceeds $g-m$, and $\max A \leq g+m$. For example, $3m \leq g$ would force a second equality for $m+2m$; any other $b \leq g-m$ would force a second equality for $m+b$. Conversely, these inequalities make all low sums except $m+m$ exceed g , and make all hole-minus-low-element

differences at most g . Excluding sums in A costs only the discarded intersections. After scaling $g = 1$, the leading contribution is therefore

$$\ell_r = \frac{1}{(r-2)!r!} \int_{1/3}^{1/2} m^{2r-2} dm = \frac{2^{-(2r-1)} - 3^{-(2r-1)}}{(2r-1)(r-2)!r!}, \quad r \geq 2. \quad (20)$$

Indeed, the $r-2$ free low elements lie in $(1-m, 1)$, of length m , and the r holes lie in $(1, 1+m)$, also of length m . The forced low element $2m$ is in the first interval. Dividing by the factorials counts unordered sets. Eliminating the coordinate equal to $2m$ preserves the integer lattice; no Euclidean surface-area factor enters this volume.

The second positive contribution comes from a valid tuple with $2m > g$ and $a_r > g + m$. Closure forces $a_r - m = a_h$ for some $h < r$. Write $a = a_h$. Away from multiple equalities, all the other holes are at most $g + m$ and all elements of $B \setminus \{m\}$ are at least $a + m - g$. These conditions are also sufficient away from the discarded intersections. At $g = 1$, $1/2 < m < 1$ and $1 < a < 2 - m$. The free holes occupy an interval of length m ; the free low elements occupy an interval of length $2 - a - m$. Hence

$$\begin{aligned} u_r &= \frac{1}{(r-2)!(r-1)!} \int_{1/2}^1 \int_1^{2-m} m^{r-2} (2-a-m)^{r-1} da dm \\ &= \frac{1}{(r-2)!r!} \int_{1/2}^1 m^{r-2} (1-m)^r dm, \quad r \geq 2. \end{aligned} \quad (21)$$

Here the eliminated coordinate is $a_r = a + m$, again an integral graph. Tuples violating both bulk inequalities have two distinct equalities and are already in the discarded set.

For the negative contribution, a bulk tuple is invalid exactly when a sum $b_i + b_j$ is a hole. Distinct such equalities overlap only on discarded intersections, so their leading contributions add. Fix $m \in (1/2, 1)$ and put $L = 1 - m$. For each low set B , the r unordered pairs involving its minimum, including (m, m) , always have sum in $(1, 1 + m)$. The other low elements lie in $(m, 1)$. Among their doubled elements, the permitted interval has half its full length. Among two distinct such elements, the permitted unordered region is a triangle of area $L^2/4$. Relative to the volume $L^{r-1}/(r-1)!$ of all choices of the free low elements, these three contributions have factors

$$r, \quad \frac{r-1}{2}, \quad \frac{(r-1)(r-2)}{4}.$$

Their sum is $r(r+3)/4$. Once a sum is forced to be a hole, the other $r-1$ holes have volume $m^{r-1}/(r-1)!$. Consequently the magnitude of the negative coefficient is

$$d_r = \frac{r(r+3)}{4((r-1)!)^2} \int_{1/2}^1 m^{r-1} (1-m)^{r-1} dm = \frac{r(r+3)}{8(2r-1)!}. \quad (22)$$

The last equality uses symmetry and the beta integral. Each sum-hole hyperplane eliminates a hole coordinate with coefficient one, so the same lattice normalization applies. We have proved, for $r \geq 2$,

$$\kappa_r = \ell_r + u_r - d_r. \quad (23)$$

5.2 Evaluation and the bulk average

For $r \geq 2$, reflection about $1/2$ and an elementary antiderivative give

$$\begin{aligned} \int_{1/2}^1 (m^r (1-m)^{r-2} + m^{r-2} (1-m)^r) dm &= \frac{r!(r-2)!}{(2r-1)!}, \\ \int_{1/2}^1 (m^r (1-m)^{r-2} - m^{r-2} (1-m)^r) dm &= \frac{1}{(r-1)4^{r-1}}. \end{aligned}$$

The second integrand is $(2m-1)m^{r-2}(1-m)^{r-2}$, the negative derivative of $m^{r-1}(1-m)^{r-1}/(r-1)$. Thus

$$u_r = \frac{1}{2(2r-1)!} - 2C_r.$$

Substituting this and (20) into (23) yields

$$\kappa_r = -\frac{(r+4)(r-1)}{8(2r-1)!} - \frac{2r}{2r-1}C_r - \frac{3(r-1)}{(2r-1)9^r r!(r-1)!}. \quad (24)$$

We also make the polynomial h_r in (16) explicit:

$$h_r(x) = -\frac{(r-1)(r-2)x^r(1-x)^{r-2} + r(r-1)x^{r-1}(1-x)^{r-1}}{2r!(r-1)!}.$$

The mean of the two lower-endpoint terms in (17) is $-f_r(1/2)/4 = -C_r/2$. Since $f_r(1) = 0$ for $r \geq 2$, the preceding integrals give

$$\gamma_r = \int_{1/2}^1 h_r(x) dx - \frac{C_r}{2} = -\frac{2r-3}{4(2r-1)!} - \left(r - \frac{3}{2}\right) C_r. \quad (25)$$

Adding (24) and (25) proves (18) for $r \geq 2$.

For $r = 1$, no valid tuple is outside the bulk: either violated bulk inequality would force $m = 0$. A bulk pair (a, m) fails only at $a = 2m$, so $n_{g,1} - M_1(g) = -\lceil g/2 \rceil$. Hence $\kappa_1 = -1/2$ and, using $\gamma_1 = 3/8$, we obtain $\beta_1 = -1/8$. Formulas (24), (25), and (18) have these same values at $r = 1$. For $r \geq 2$, all three summands of (18) are negative, proving the asserted sign.

Finally, $\binom{2r}{r}/4^r \sim 1/\sqrt{\pi r}$ implies

$$A_r \sim \frac{1}{4r(2r-1)!}, \quad \beta_r \sim -\frac{r^2}{8(2r-1)!}.$$

The remaining terms in (18) are smaller by the same central-binomial estimate and the exponential factor $(4/9)^r$. Taking the ratio proves (19) and completes the proof of Theorem 5.1.

5.3 Exact sums for reproducing the leading volumes

The three regions above admit convenient finite counting formulas. These count candidates, with multiplicity for the negative contribution; they are not exact formulas for the correction $n_{g,r} - M_r(g)$. For $r \geq 2$, put

$$L_r(g) = \sum_{\lfloor g/3 \rfloor < m \leq \lfloor g/2 \rfloor} \binom{m-1}{r-2} \binom{m}{r}, \quad (26)$$

$$U_r(g) = \sum_{\lfloor g/2 \rfloor < m \leq g} \binom{m-1}{r-2} \binom{g-m+1}{r}. \quad (27)$$

The first formula removes the forced element $2m$ from the m possible free-low positions. For the second, sum over the lower forced hole a : the number of available other low elements is $2g-a-m+1$. The hockey-stick identity then sums their binomial coefficients. For every $r \geq 1$, define

$$D_r(g) = \sum_{m=\lfloor g/2 \rfloor+1}^g \binom{m-1}{r-1} \left[r \binom{g-m}{r-1} + \left\lfloor \frac{g-m}{2} \right\rfloor \binom{g-m-1}{r-2} + \left\lfloor \frac{(g-m-1)^2}{4} \right\rfloor \binom{g-m-2}{r-3} \right]. \quad (28)$$

A binomial coefficient in these formulas is zero unless its two indices are integers with $0 \leq k \leq n$. The last term counts distinct unordered $u, v \in [1, g - m]$ with $u + v \leq g - m$. There are $\lfloor (g - m - 1)^2 / 4 \rfloor$ such pairs. Set $L_1 = U_1 = 0$.

The equality-arrangement argument proves

$$n_{g,r} - M_r(g) = L_r(g) + U_r(g) - D_r(g) + O_r(g^{2r-2}). \quad (29)$$

In particular their leading coefficients are ℓ_r, u_r, d_r , respectively. Formula (29) retains a remainder from multiple additive equalities. It must not be used as an exact identity or as a uniform estimate with r increasing.

6 Consequences, checks, and remaining questions

The first three leading coefficients are $A_1 = 3/8$, $A_2 = 11/384$, and $A_3 = 7/7680$. The first two agree with the exact formulas of [2, Proposition 2.1 and Theorem 3.5]. These small values illustrate the general coefficient formula; our contribution concerns arbitrary fixed r .

The homogeneous description is also effective in principle. For a specified r , form the finite arrangement in the proof of Proposition 2.3, enumerate its rational cells, and compute their Ehrhart quasipolynomials. A common multiple of the vertex denominators bounds a period. The argument gives no efficient bound for this enumeration. Once the finitely many constituents are obtained, eventual positivity of (3) reduces the full monotonicity question for that particular r to finitely many genera.

The accompanying Python script compares literal closure with Lemma 2.1 on all 66 187 tuples through genus 9, checking all 584 exceptional tuples. Independently generated semigroup trees through genus 16 contain 11 770 semigroups in total; their counts agree with the tuple counts and the known $r = 1, 2$ formulas on the overlapping ranges. Exact rational interpolation tests the bulk's two leading coefficients for $1 \leq r \leq 12$, with 936 additional evaluations beyond the interpolation data. For the revised second-coefficient formula, a separate script checks all 17 568 tuples through genus 8, verifies that all 223 nonzero signed residuals satisfy two independent additive equalities, and checks the candidate-sum formulas. Exact rational interpolation of those sums for $1 \leq r \leq 8$ has 864 holdout evaluations; the resulting coefficients also agree with every residue of the cited exact $r = 2$ formula. These finite diagnostics do not replace the general proofs.

Several questions remain. Theorem 5.1 evaluates the previously undetermined constant β_r . One can next determine the full period of $n_{g,r}$, evaluate the lower-dimensional multiple-equality contributions, and find bounds for the onset of strict growth. The stronger conjecture asks for monotonicity at every genus and every ordinarization number. The present argument now evaluates the top two coefficients completely for fixed r and leaves those further questions open. The coefficient ratio in (19) records growth of the correction with r ; it supplies no uniform remainder bound and does not settle either total monotonicity or the full Bras-Amorós Fibonacci inequality.

References

- [1] M. Beck and S. Robins, *Computing the Continuous Discretely: Integer-Point Enumeration in Polyhedra*, 2nd ed., Undergraduate Texts in Mathematics, Springer, 2015. doi:10.1007/978-1-4939-2969-6.
- [2] S. Cyrusian and N. Kaplan, Ordinarization numbers of numerical semigroups, *Communications in Algebra* (2026), 1–24. doi:10.1080/00927872.2026.2615092. Also arXiv:2506.10222v2; theorem numbering follows this version.