

Closed-shell parametrization

September 18, 2025

1 Notation

1.1 Excitation operators

Let $\hat{a}_{p,\sigma}^\dagger$ ($\hat{a}_{p,\sigma}$) be the creation (annihilation) operator associated with a spin-orbital

$$\psi_p(\mathbf{x}) \equiv \phi_p(\mathbf{r}) \otimes \sigma(m_s), \quad (1)$$

where $\phi_p(\mathbf{r})$ is a spatial orbital, and $\sigma(m_s)$ a spin eigenfunction. The spin functions are orthonormal in the sense

$$\int \sigma^*(m_s) \tau(m_s) dm_s = \delta_{\sigma,\tau}. \quad (2)$$

Following Helgaker et al. [1], the singlet excitation operators are defined as

$$\hat{E}_{pq} \equiv \hat{a}_{p,\alpha}^\dagger \hat{a}_{q,\alpha} + \hat{a}_{p,\beta}^\dagger \hat{a}_{q,\beta}, \quad (3)$$

and satisfy the commutation relations

$$[\hat{E}_{pq}, \hat{E}_{rs}] = \hat{E}_{ps} \delta_{rq} - \hat{E}_{rq} \delta_{ps}. \quad (4)$$

Moreover, we introduce the two-electron excitation operator

$$\hat{e}_{pqrs} \equiv \hat{E}_{pr} \hat{E}_{qs} - \delta_{rq} \hat{E}_{ps} = \sum_{\sigma\tau} \hat{a}_{p,\sigma}^\dagger \hat{a}_{q,\tau}^\dagger \hat{a}_{s,\tau} \hat{a}_{r,\sigma}, \quad (5)$$

which satisfy the following commutation relation with $\hat{E}_{t,u}$

$$[\hat{e}_{pqrs}, \hat{E}_{tu}] = \hat{e}_{pqr u} \delta_{ts} - \hat{e}_{ptrs} \delta_{qu} + \hat{e}_{pqus} \delta_{tr} - \hat{e}_{tqrs} \delta_{pu}. \quad (6)$$

1.2 The Hamiltonian

The second quantized representation of a nonrelativistic spin-free Hamiltonian containing at most two-particle interactions can now be written in terms of singlet and two-electron excitation operators as

$$\hat{H} = \sum_{pq} h_q^p \hat{E}_{pq} + \frac{1}{2} \sum_{pqrs} u_{rs}^{pq} \hat{e}_{pqrs}, \quad (7)$$

where the one- and two-body matrix elements h_q^p and u_{rs}^{pq} (physicist notation) are defined as

$$h_q^p \equiv \int \phi_p^*(\mathbf{r}) \hat{h} \phi_q(\mathbf{r}) d\mathbf{r}, \quad (8)$$

$$u_{rs}^{pq} \equiv \iint \phi_p^*(\mathbf{r}_1) \phi_q^*(\mathbf{r}_2) \hat{w}(\mathbf{r}_1, \mathbf{r}_2) \phi_r(\mathbf{r}_1) \phi_s(\mathbf{r}_2) d\mathbf{r}_1 d\mathbf{r}_2, \quad (9)$$

where \hat{h} is a one-body operator and $\hat{w}(\mathbf{r}_1, \mathbf{r}_2)$ is a two-particle interaction/operator (typically the electron-electron Coulomb interaction).

The expectation value of the (spin-free) Hamiltonian for a (closed-shell) reference determinant (usually the Hartree-Fock determinant) is given by

$$E_{\text{ref}} \equiv \langle \Phi_0 | \hat{H} | \Phi_0 \rangle = 2 \sum_i h_i^i + 2 \sum_{ij} u_{ij}^{ij} - \sum_{ij} u_{ji}^{ij}, \quad (10)$$

where i, j run over all (doubly) occupied orbitals $\{\phi_i(\mathbf{r})\}_{i=1}^{N_{\text{docc}}}$. More generally, the expectation value of the Hamiltonian (or some other one- plus two-body operator) where $\langle \tilde{\Psi} |, |\Psi \rangle$ are (possibly independent) bra and ket states can be written as

$$\langle \tilde{\Psi} | \hat{H} | \Psi \rangle = \sum_{pq} h_q^p \gamma_p^q + \frac{1}{2} \sum_{pqrs} u_{rs}^{pq} \Gamma_{pq}^{rs}, \quad (11)$$

where we have defined the one- and two-body reduced density matrices

$$\gamma_p^q \equiv \langle \tilde{\Psi} | \hat{E}_{pq} | \Psi \rangle, \quad (12)$$

$$\Gamma_{pq}^{rs} \equiv \langle \tilde{\Psi} | \hat{e}_{pqrs} | \Psi \rangle. \quad (13)$$

The expressions for the reduced density matrices depend on the particular wave-function parametrization used, i.e., Hartree-Fock, configuration interaction, coupled cluster, and approximate coupled cluster methods such as CC2 or OMP2.

1.3 Singlet configuration state functions

Single and double excited (ket/bra) (singlet) configuration state functions (CSFs) are defined as

$$|\Phi_i^a\rangle \equiv \hat{X}_i^a |\Phi_0\rangle, \quad |\Phi_{ij}^{ab}\rangle \equiv \hat{X}_{ij}^{ab} |\Phi_0\rangle, \quad (14)$$

$$\langle \Phi_i^a | \equiv \langle \Phi_0 | E_{ia}, \quad \langle \Phi_{ij}^{ab} | \equiv \langle \Phi_0 | E_{jb} E_{ia}, \quad (15)$$

where we have defined the single and double excitation (de-excitation) operators

$$\hat{X}_i^a \equiv \hat{E}_{ai}, \quad \hat{X}_{ij}^{ab} \equiv \hat{E}_{ai} \hat{E}_{bj}, \quad (16)$$

$$(\hat{X}_i^a)^\dagger = \hat{E}_{ia}, \quad (\hat{X}_{ij}^{ab})^\dagger = \hat{E}_{jb} \hat{E}_{ia}. \quad (17)$$

The CSFs defined this way are not orthogonal, and the overlaps are given by

$$\langle \Phi_i^a | \Phi_j^b \rangle = 2\delta_{ab}\delta_{ij}, \quad (18)$$

$$\langle \Phi_{ij}^{ab} | \Phi_{kl}^{cd} \rangle = 2\hat{P}_{ij}^{ab} (2\delta_{ac}\delta_{bd}\delta_{ik}\delta_{jl} - \delta_{ac}\delta_{bd}\delta_{il}\delta_{jk}), \quad (19)$$

where we have defined the permutation operator

$$\hat{P}_{rs}^{pq} A_{rs}^{pq} = A_{rs}^{pq} + A_{sr}^{qp}. \quad (20)$$

However, following Refs. [1, 2] a biorthogonal basis can be defined (for singles and doubles), such that,

$$\langle \tilde{\Phi}_i^a | \Phi_k^c \rangle = \delta_{ac} \delta_{ik}, \quad (21)$$

$$\langle \tilde{\Phi}_{ij}^{ab} | \Phi_{kl}^{cd} \rangle = P_{ij}^{ab} \delta_{ac} \delta_{bd} \delta_{ik} \delta_{jl}, \quad (22)$$

by choosing

$$\langle \tilde{\Phi}_i^a | = \frac{1}{2} \langle \Phi_i^a | = \langle \Phi_0 | \hat{Y}_a^i, \quad (23)$$

$$\langle \tilde{\Phi}_{ij}^{ab} | = \frac{1}{3} \langle \Phi_{ij}^{ab} | + \frac{1}{6} \langle \Phi_{ji}^{ab} | = \langle \Phi_0 | \hat{Y}_{ab}^{ij}, \quad (24)$$

where we have defined the single and double de-excitation operators

$$\hat{Y}_a^i = \frac{1}{2} \hat{E}_{ia}, \quad (25)$$

$$\hat{Y}_{ab}^{ij} = \frac{1}{3} \hat{E}_{jb} \hat{E}_{ia} + \frac{1}{6} \hat{E}_{ib} \hat{E}_{ja} \quad (26)$$

2 Configuration interaction

2.1 CISD

We use the following parametrization of the CISD ket and bra wavefunction

$$|\Psi\rangle = C_0 |\Phi_0\rangle + \sum_{ai} C_i^a |\Phi_i^a\rangle + \frac{1}{2} \sum_{abij} C_{ij}^{ab} |\Phi_{ij}^{ab}\rangle, \quad (27)$$

$$\langle \tilde{\Psi} | = \langle \tilde{\Phi}_0 | \tilde{C}_0 + \sum_{ia} \tilde{\Phi}_i^a \tilde{C}_a^i + \frac{1}{2} \sum_{ijab} \langle \tilde{\Phi}_{ij}^{ab} | \tilde{C}_{ab}^{ij}, \quad (28)$$

where the left expansion coefficients \tilde{C} are related to the right coefficients C by

$$\tilde{C}_0 = C_0^*, \quad (29)$$

$$\tilde{C}_a^i = 2C_i^{a*}, \tilde{C}_{ab}^{ij} = (4C_{ij}^{ab} - 2C_{ji}^{ab})^*. \quad (30)$$

The norm/overlap of the bra and ket states is given by

$$\langle \tilde{\Psi} | \Psi \rangle = \tilde{C}_0 C_0 + \sum_{ai} \tilde{C}_a^i C_i^a + \frac{1}{2} \sum_{abij} \tilde{C}_{ab}^{ij} C_{ij}^{ab}, \quad (31)$$

Right and left sigma-vectors are defined as

$$\sigma_\mu \equiv \sum_\nu \langle \tilde{\Phi}_\mu | \hat{H} | \Phi_\nu \rangle C_\nu, \quad (32)$$

$$\tilde{\sigma}_\nu \equiv \sum_\mu \tilde{C}_\mu \langle \tilde{\Phi}_\mu | \hat{H} | \Phi_\nu \rangle. \quad (33)$$

Working equations/algebraic expressions for the (closed-shell) sigma-vectors and the one- and two-body density matrices can be generated symbolically with the `drudge/gristmill` package [3, 4].

The CIS and CID approximation are obtained by setting the doubles or singles coefficients to zero, respectively.

3 Coupled cluster

3.1 CCSD

The right and left CCSD wavefunctions are parametrized as

$$|\Psi\rangle = e^{\hat{T}} |\Phi_0\rangle, \quad \langle\tilde{\Psi}| = \langle\Phi_0| (1 + \hat{\Lambda}) e^{-\hat{T}}, \quad (34)$$

where we have defined the cluster excitation (de-excitation) operators

$$\hat{T} = \hat{T}_1 + \hat{T}_2, \quad \hat{\Lambda} = \hat{\Lambda}_1 + \hat{\Lambda}_2 \quad (35)$$

$$\hat{T}_1 = \sum_{ai} \tau_i^a \hat{X}_i^a, \quad \hat{T}_2 = \frac{1}{2} \sum_{abij} \tau_{ij}^{ab} \hat{X}_{ij}^{ab}, \quad (36)$$

$$\hat{\Lambda}_1 = \sum_{ia} \lambda_a^i \hat{Y}_a^i, \quad \hat{\Lambda}_2 = \frac{1}{2} \sum_{ijab} \lambda_{ab}^{ij} \hat{Y}_{ab}^{ij}, \quad (37)$$

and the CCSD energy functional is given by

$$\mathcal{H} = \langle\tilde{\Psi}|\hat{H}|\Psi\rangle. \quad (38)$$

The CCSD-amplitudes (λ, τ) (for the ground state) are determined by solving the (non-linear) amplitude equations

$$\frac{\partial \mathcal{H}}{\partial \lambda_\mu} = \langle\Phi_0|\hat{Y}^\mu e^{-\hat{T}} \hat{H} e^{\hat{T}} |\Phi_0\rangle = 0, \quad (39)$$

$$\frac{\partial \mathcal{H}}{\partial \tau^\mu} = \langle\Phi_0| (1 + \hat{\Lambda}) e^{-\hat{T}} [\hat{H}, \hat{X}_\mu] e^{\hat{T}} |\Phi_0\rangle = 0, \quad (40)$$

where μ denotes the excitation level (singles or doubles).

$$\gamma_p^q \equiv \langle\tilde{\Psi}|\hat{E}_{pq}|\Psi\rangle, \quad (41)$$

$$\Gamma_{pq}^{rs} \equiv \langle\tilde{\Psi}|\hat{e}_{pqrs}|\Psi\rangle \quad (42)$$

Excited states can be determined with the EOM-CCSD approach, where the left and right excited states are parametrized as

$$|\Psi_k\rangle = \hat{R}_k e^{\hat{T}} |\Phi_0\rangle, \quad (43)$$

$$\langle\tilde{\Psi}_k| = \langle\tilde{\Phi}_0| \hat{L}_k e^{-\hat{T}}, \quad (44)$$

where

$$\hat{R}_k = \sum_{\mu=0} r_k^\mu \hat{X}_\mu, \quad \hat{L}_k = \sum_{\mu=0} l_{k,\mu} \hat{Y}^\mu. \quad (45)$$

The EOM coefficients l, r are determined by solving the left and right eigenvalue problems

$$\sigma_\mu = E_k r_k^\mu, \quad (46)$$

$$\tilde{\sigma}_\nu = E_k l_{\nu,k} \quad (47)$$

where we have defined the EOM sigma-vectors

$$\sum_{\nu} \bar{H}_{\mu,\nu} r_k^\nu \equiv \sigma_\mu, \quad (48)$$

$$\sum_{\mu} l_{k,\mu} \bar{H}_{\mu,\nu} \equiv \tilde{\sigma}_\nu, \quad (49)$$

and the matrix elements of the similarity transformed Hamiltonian $\bar{H} \equiv e^{-\hat{T}} \hat{H} e^{\hat{T}}$

$$\bar{H}_{\mu,\nu} \equiv \langle \Phi_\mu | e^{-\hat{T}} \hat{H} e^{\hat{T}} | \Phi_\nu \rangle. \quad (50)$$

Moreover, the right and left eigenvectors can be normalized such that they form a bi-orthonormal set, i.e.,

$$\langle \tilde{\Psi}_k | \Psi_l \rangle = \delta_{k,l}. \quad (51)$$

Working equations for the (closed-shell) amplitude equations and the sigma vectors can be generated with the **drudge/gristmill** package.

References

- ¹T. Helgaker, P. Jorgensen, and J. Olsen, *Molecular electronic-structure theory* (John Wiley & Sons, 2013).
- ²P. Pulay, S. Saebo/, and W. Meyer, “An efficient reformulation of the closed-shell self-consistent electron pair theory”, *The Journal of chemical physics* **81**, 1901–1905 (1984).
- ³J. Zhao, G. P. Chen, G. Harsha, M. Wholey, T. M. Henderson, and G. E. Scuseria, *Drudge: a symbolic algebra system for tensorial and noncommutative algebras*, GitHub repository, 2016–2025.
- ⁴J. Zhao, “Symbolic solution for computational quantum many-body theory development”, PhD thesis (Rice University, Houston, Texas, USA, Apr. 2018).