GraphPathology enables geometry-aware reconstruction of spatial transcriptomics from sparse molecular measurements

Truong-Son Hy[1,*]

[1]Department of Computer Science, University of Alabama at Birmingham, Birmingham, Alabama, United States

[*]Correspondence: thy@uab.edu

Abstract

Reconstructing spatial transcriptomics from limited molecular measurements requires accounting for where observations are collected. GraphPathology combines observed molecular anchors with spatial representations of tissue morphology. We investigated different forms of graph neural networks (GNNs). In two breast-cancer cohorts comprising 25 evaluation patients, reconstruction from 20% distributed molecular observations achieved median patient Pearson correlations of 0.359 and 0.416. We then examined measurement placement in 22 previously evaluated patients, holding the hidden locations, reconstruction method and number of observed sites fixed. Excluding anchors from a central region reduced median paired patient correlation by 0.154 and increased mean squared error by 45.0% for inverse-distance interpolation; all 22 patients deteriorated. Across matched external comparisons, the tested GNN predictors did not consistently improve reconstruction over the same-information molecular baseline. These findings identify anchor placement as an important determinant of sparse reconstruction for the specified sampling design and delimit the contribution of morphology-guided modeling under cross-cohort conditions. Measuring one fifth of locations offers a conditional opportunity to reduce location-dependent costs; total assay savings require prospective validation. Code: https://github.com/HySonLab/GraphPathology.

Keywords: spatial transcriptomics; sparse reconstruction; geometry-aware graph learning; spatial transcriptomics imputation; histopathology; breast cancer

1 Introduction

Spatial transcriptomics (ST) links expression measurements to tissue organization[1], but dense profiling can limit study scale. Sparse measurement designs raise two related questions: how the placement of measured sites affects reconstruction, and whether histology supplies useful information beyond the molecular observations already available. Answering these questions requires separating the information supplied to a predictor from the spatial design on which it is evaluated.

We study reconstruction of a fixed gene panel within a tissue section. Complete panel measurements are available at observed anchors; expression at hidden sites is withheld, while their coordinates and histology remain known. This setting differs from predicting an entirely

unassayed panel or inferring expression from images alone. Breast-cancer ST studies provide paired data for examining these distinctions across tissue sections and platforms[2–5].

1.1 Related work

Histology-based prediction ranges from local convolutional methods such as ST-Net[6] and DeepSpaCE[7] to models with contextual representations. Hist2ST combines Transformer and graph components[8]; THItoGene[9] and FmH2ST[10] further examine visual context and feature-informed links between distant regions. CMRCNet[11] and cross-cohort benchmarking[12] highlight the importance of transportability. Graph convolutional networks (GCNs)[13], GraphSAGE[14], GraphGPS[15] and equivariant graph networks[16] offer complementary representation principles; their application does not by itself establish a new biological interaction or a reconstruction advantage.

Some methods combine measured expression with images, including XFuse[17], TESLA[18] and iStar[19]. Sparse-acquisition and tissue-wide mapping systems include Sparser2Sparse[20], SCR^2-ST[21] and PanoSpace[22], with differing modalities and sampling assumptions. SPACEL addresses spatial-domain and multi-section tasks[23]. Their training requirements, target resolutions and available inputs differ from ours. We therefore do not infer superiority or equivalence to these systems from comparisons with a frozen image predictor or simple molecular interpolation.

1.2 Framework and study objectives

GraphPathology distinguishes two graph roles. A coordinate-defined tissue-context graph aggregates histological representations, whereas an observation-dependent molecular-support graph connects hidden sites to measured anchors. The latter transports measured expression and supplies the strongest external reconstruction signal in this study. Keeping these roles separate permits same-information comparisons of molecular estimators and tests of the incremental contribution of image context.

We evaluate distributed reconstruction across two breast-cancer studies, isolate anchor placement in a controlled 22-patient comparison with identical hidden targets and equal anchor counts, and investigate several GNN representations against molecular and conventional neural baselines. Coverage, contiguous missing regions and distance-resolved errors characterize failure regimes. The purpose is to establish what the available measurements support, including negative results, rather than to equate architectural complexity with novelty or measured-location reduction with total assay savings.

2 Results

2.1 Study design separated information access and dataset roles

The task used a fixed 50-gene panel and paired spatial expression and histology (Fig. 1; Table 1). Four HEST patients supplied 35,536 locations: two patients for fitting, one for validation and one for development evaluation. Eight HER2ST patients (3,455 locations) supported transportability and calibration development; their reuse precludes treating them as

untouched external validation. The original external evaluation comprised three held-out Wu patients and the first numbered section from each of 22 Bassiouni patients, totaling 25 patients and 38,484 locations. All 50 genes were retained externally; HER2ST used a fixed 40-gene intersection.

The additional placement controls and neural comparisons reused these already evaluated cohorts and one development patient. They are exploratory follow-up analyses, not new blind replication. Image-only predictors receive histology and permitted geometry, molecular estimators receive coordinates and same-section anchor vectors, and fusion or residual predictors receive both. Thus image-only versus molecular comparisons quantify the value of measured anchors under unequal information access; incremental image value requires comparison with the same anchors supplied to both methods. The public external images were downsampled, containing only 15/25 native pixels in Wu and 24–29 in Bassiouni across the nominal 112-μm patch.

Table 1: Cohorts and their analytical roles. Counts refer to retained locations and patient-level study roles, not independent spot replicates. The same eight HER2ST patients were first inspected for transportability and subsequently used for calibration development. The Wu study contains 15,601 locations in all six patients; only 8,481 locations from its three evaluation patients enter the original external evaluation. All 22 Bassiouni patients were evaluated without new training or calibration. "Pixels per patch" refers to the native image information in a nominal 112-μm field of view, before resizing. The primary panel was fixed in development.

Cohort	Platform	Patients, total / evaluation	Locations, total / evaluation	Genes	Analytical role and image quality
Development (HEST)	Targeted Xenium	4 / 1	35,536 / 20,761	50/50	Two training, one validation, one primary evaluation patient; conditional four-patient sensitivity analyses. Estimated 0.21–0.36 μm/pixel.
HER2ST	Sequencing ST	8 / 8	3,455 / 3,455	40/50	Initial transportability, then calibration development on the same patients. Approximately 0.69–0.92 μm/pixel.
Wu	Visium	6 / 3	15,601 / 8,481	50/50	Three calibration and three original evaluation patients. Only 15 or 25 native pixels per patch.
Bassiouni	Visium	22 / 22	30,003 / 30,003	50/50	First numbered section per patient; original external evaluation with no cohort-specific fitting. Only 24–29 native pixels per patch.




Figure 1: Study design and neural-image/molecular reconstruction architecture. a, Cohort roles and illustrative observation geometries. Teal locations are observed molecular anchors; gray locations are hidden; purple locations form the contiguous scored region. The synthetic spatial examples illustrate 20% coverage and are not patient measurements. The Wu calibration and evaluation groups contain different patients. **b,** The frozen ImageNet ResNet18 maps histology to a 512-dimensional vector after global average pooling; the ImageNet classifier is removed. External nominal 112-μm patches are first resized to 224×224 with LANCZOS, then processed by the official resize-to-256, central-crop-to-224 and channel-normalization transform. Feature scaling fitted on the training patients precedes each $512 \to 256 \to 50$ MLP, with GELU activation and dropout 0.1 after the hidden linear layer. The three heads use seeds 0, 1 and 2. Their predictions are inverse-scaled to the common target units and averaged. **c,** Molecular interpolation uses up to 16 observed neighbors, with inverse-distance-squared weighting for distributed evaluation (V1) and equal weighting for contiguous regional completion (V2). Fixed fusion combines this output with the calibrated image prediction using weights 0.8 and 0.2. The uncalibrated image ensemble is also evaluated separately. Calibration uses the development-training and Wu calibration patients and is transported unchanged to Bassiouni. Evaluation uses hidden locations only. V1 and V2 differ in weighting and scoring support as well as observation geometry.




Figure 2: Geometry-aware graph construction and contextual representations. a–d, The same 36 synthetic tissue locations are connected by (a) a symmetrized six-nearest-neighbor graph with a 250-μm distance cap, (b,c) 150- and 250-μm radius neighborhoods, or (d) a 250-μm-capped Delaunay graph. Coordinates and axes are identical across panels and all retained edges are displayed. Symmetrization means that a node can have more than six neighbors. **e,** Geometry determines a fixed adjacency/aggregation operator acting on the frozen image embeddings before an expression-regression head. The molecular-reconstruction stage uses a separate observation-dependent graph joining hidden locations to up to 16 measured molecular anchors. Synthetic points illustrate the architecture only, not a fitted patient graph or a performance result.

2.2 Distributed molecular observations supported reconstruction across cohorts

At 20% distributed coverage in TENX99, median gene Pearson correlations were 0.695 for image-only prediction, 0.792 for interpolation and 0.822 for fixed fusion; corresponding mean squared errors (MSEs) were 0.911, 0.268 and 0.279 (Fig. 3; Table 2). The correlation increase from fusion therefore did not imply an error reduction. These are one-patient development results.




Figure 3: Development reconstruction depends on molecular coverage and masking regime. Results are from the single primary development patient, TENX99, using the frozen 50-gene panel and ResNet18 representation. **a,c,** Distributed random observation. **b,d,** The earlier development block-mask regime. Panels a and b show median gene Pearson correlation; panels c and d show mean gene MSE across measurement fractions. The image-only, selected molecular interpolation and fixed fusion curves summarize three frozen mask replicates per condition; curves are means over masks, not patient confidence intervals. Targets and error units are unchanged across methods. The development block masks differ from the external V2 forbidden-region protocol and should not be treated as directly equivalent geometry tests. At 20% distributed coverage, fusion has higher correlation than interpolation but slightly higher MSE, illustrating that spatial association and calibration are distinct properties. The image-only comparator receives no observed molecular anchors.

In the original Wu evaluation, the median of patient-level median gene correlations was 0.359 for interpolation and 0.059 for image-only prediction. Corresponding MSEs were 0.0987 and 2.9155, and mean absolute errors (MAEs) were 0.2179 and 1.5578 (Fig. 4). Bassiouni interpo-

lation achieved correlation 0.416, MSE 0.0923 and MAE 0.2033, compared with image-only values 0.075, 1.9673 and 1.2120 (Fig. 5). Every evaluation patient favored interpolation in correlation, MSE and MAE at each tested budget. All 22 Bassiouni patients met the predefined joint support criterion, which was descriptive rather than a significance test.

Table 2: Reconstruction at 20% distributed measurement coverage. Each entry is a cohort median of patient-level summaries after averaging mask seeds within geometry and geometries within patient, except that the development row represents the single primary patient. Pearson r is the median correlation across panel genes; MSE and MAE are patient means across genes and hidden locations before the cohort median. All methods within a cohort use identical target units and observation masks. Different gene panels and assay count distributions preclude treating these values as a pooled cross-cohort ranking. Fusion rules and their calibration are frozen but differ between the development and external protocols; see Methods and Supplementary Information. The image-only branch has no same-section molecular anchors.

Cohort	n	Genes	Method	Pearson r	MSE	MAE
Development	1	50	Image only	0.695	0.9110	0.7606
			Spatial interpolation	0.792	0.2684	0.3904
			Fixed fusion	0.822	0.2787	0.4173
HER2ST, zero-shot	8	40	Image only	0.143	4.3665	1.9376
			Spatial interpolation	0.194	0.0680	0.1730
			Fixed fusion	0.255	0.3481	0.5153
Wu, original	3	50	Image only	0.059	2.9155	1.5578
			Spatial interpolation	0.359	0.0987	0.2179
			Fixed fusion	0.335	0.1002	0.2321
Bassiouni, original	22	50	Image only	0.075	1.9673	1.2120
			Spatial interpolation	0.416	0.0923	0.2033
			Fixed fusion	0.407	0.1022	0.2288

Increasing distributed coverage from 10% to 40% improved correlation and MSE in every patient. Bassiouni correlations at 10/20/40% were 0.380/0.416/0.465, with MSEs 0.1001/0.0923/0.0852; Wu values were 0.313/0.359/0.393 and 0.1080/0.0987/0.0899 (Supplementary Table S9). These budget-specific protocols use nested anchor sets but different hidden sets. They neither identify a universally sufficient budget nor compare an identical target set across coverages. Earlier sampling comparisons, including label-free spatial coverage sampling, are retained in Supplementary Fig. S8.

2.3 Controlled anchor placement altered reconstruction in all 22 patients

The controlled experiment fixed a central region R and a hidden subset H, then changed only the eligible placement of an equal number of anchors. Distributed anchors could occupy any non-H location; region-excluded anchors were restricted outside R. Within each comparison, the estimator, gene panel, hidden targets and total anchor count were identical. This design isolates placement for the specified central-region sampling rule.

At 20% coverage, region exclusion worsened Pearson, MSE and MAE in all 22 Bassiouni patients for both fixed estimators (Fig. 6; Table 3). For inverse-distance weighting (IDW), the median paired correlation change was −0.1541 (descriptive 95% patient-bootstrap inter-




Figure 4: Fresh evaluation in three held-out Wu patients supports interpolation but not robust image fusion. The frozen 50-gene panel was evaluated in patients 1160920F, CID4290 and CID44971, none of whom contributed to Wu calibration. **a,** Patient-level correlation summaries at 20% distributed coverage. **b,c,** Median patient correlation and MSE across 10%, 20% and 40% distributed coverage. **d,** Paired patient correlation effects for interpolation versus image-only prediction and fusion versus interpolation. **e,** Paired distributed (V1) and contiguous-region (V2) molecular correlations at 20%. **f,** Distance-resolved molecular MSE at 20%. Patient summaries average ten random mask seeds within geometry and then geometry means; cohort summaries are medians, not pooled spot estimates. Three fixed contiguous geometries are used for V2. Each patient is a biological replicate; seeds are not. V1 uses inverse-distance-squared interpolation and V2 uses a nearest-anchor mean, so panel e is a protocol contrast. The public image inputs contained only 15 or 25 native pixels per 112-μm patch, limiting conclusions about morphology. Error units are common development-standardized log1p counts.




Figure 5: Independent 22-patient replication confirms the coverage response and weaker contiguous completion. One first-numbered section per Bassiouni patient was selected before evaluation, yielding 30,003 locations and the same 50 primary genes. No patient from this study was used for new training or calibration. **a,** Patient correlation summaries at 20% distributed coverage. **b,c,** Correlation and MSE across measurement fractions. **d,** Paired interpolation–image and fusion–interpolation correlation effects. **e,** Paired V1/V2 molecular correlations at 20%. **f,** Distance-resolved molecular MSE at 20%. Each patient summary averages ten mask seeds within geometry and then the three V2 geometry means where applicable; cohort curves use patient medians. All 22 patients improved in correlation and MSE between 10% and 40% V1 coverage and deteriorated in correlation, MSE and MAE under V2 relative to V1 at 20%. V1/V2 differ in scoring support and interpolation weighting as well as missing-region design. Image fusion uses the frozen Wu calibration and 0.2 image weight. Public histology contained only 24–29 native pixels per 112-μm patch. Distance curves have varying occupied-bin support and are not calibrated uncertainty intervals. No p-values or patient confidence intervals are implied.

val −0.2056 to −0.1096), relative MSE increased 45.0% (29.8–52.2%) and MAE increased 0.0335 (0.0266–0.0508). Uniform interpolation gave a correlation change of −0.1502 (−0.1805 to −0.1094), relative MSE increase 39.9% (29.2–53.5%) and MAE increase 0.0326 (0.0234–0.0462). All three budgets and all patient effects are retained. These intervals describe uncertainty across the 22 patients; they are not p-values or evidence of a universal geometry effect.




Figure 6: Identical-target placement effects in all 22 patients. Each point is one patient, averaged over ten mask seeds; shapes distinguish fixed IDW and uniform k16. Both anchor placements score the same H with equal anchor counts. Panels retain all three budgets. These are not historical V2-minus-V1 effects.

Table 3: Controlled Bassiouni placement effects. Region-exclusion minus distributed anchors at 20% coverage, identical H and anchor counts. Median paired patient effects with descriptive 95% percentile intervals from 10,000 patient resamples. Positive error changes deteriorate. All 22 patients are retained; masks are averaged within patient. The full 20-estimator and three-budget exports accompany the paper.

Operator	Effect	Median [95% interval]
IDW	Δr	-0.1541 [-0.2056, -0.1096]
IDW	MSE%	+44.9864 [+29.8116, +52.2347]
IDW	Δ MAE	+0.0335 [+0.0266, +0.0508]
Uniform	Δr	-0.1502 [-0.1805, -0.1094]
Uniform	MSE%	+39.9128 [+29.2387, +53.4974]
Uniform	Δ MAE	+0.0326 [+0.0234, +0.0462]

The three previously evaluated Wu patients corroborated the direction: at 20% coverage, region exclusion reduced IDW correlation by 0.106–0.245 and increased relative MSE by 20.1–45.4%; uniform reconstruction lost 0.093–0.188 in correlation and increased MSE by 21.7–38.9% (Fig. 7b; Supplementary Table S12). Each patient is reported individually. The original distributed-versus-regional protocol comparison is distinct: it changes geometry, estimator and scored support, rather than placement alone.

2.4 Matched spatial baselines separated estimator choice from sampling design

We compared twenty fixed molecular estimators on identical distributed masks in Wu and Bassiouni at each of three coverage levels: 10%, 20% and 40%. The grid included nearest-anchor, uniform, IDW and Gaussian reconstruction; all declared estimators are shown, without choosing an outcome-selected winner (Fig. 8; Supplementary Table S10 and source tables). Separately, IDW and uniform interpolation were crossed with the original distributed and regional masks (Fig. 7a; Supplementary Table S11). This separates the effect of changing the estimator from the controlled placement comparison, while retaining the original protocol-level results. Predetermined gene maps include poorly reconstructed and nearly constant fields (Supplementary Figs. S13,S14,S26 and S27).




Figure 7: Matched estimator and placement controls in previously inspected Wu patients. (a) Both operators share each historical mask and scored set; V2 values average the three geometries. (b) Region-exclusion minus distributed correlation on the identical H, with equal total anchor count and ten random seeds. The three individual patients are shown; masks and genes are not biological replicates. All points use 20% coverage and the frozen 50-gene panel.




Figure 8: Complete matched-information baseline grid in 22 previously inspected Bassiouni patients. Every declared estimator and budget is shown. Points are medians of paired patient effects relative to fixed IDW k16/p2 after averaging ten original mask trajectories. Correlation increases and error decreases improve. No estimator was selected from these outcomes.

2.5 Graph representations did not add a consistent external benefit

The GNN comparisons included local multilayer perceptrons, fixed mean/GCN/GraphSAGE representations, bounded nonlocal models and fixed-distance or latent-coordinate message passing. Their architectural and optimization regimes are distinguished in Methods. In the 30-head minibatch development analysis, image-only seed-mean correlations for local, GCN and GraphSAGE predictors were 0.682, 0.654 and 0.678, with MSEs 0.9114, 0.8999 and 0.8859. Residual-anchor correlations were 0.821, 0.810 and 0.825, with MSEs 0.2567, 0.2510 and 0.2338; the identical-mask IDW reference was 0.792 and 0.2684. These one-patient results did not establish a topology-specific advantage (Fig. 9; Supplementary Table S13). The full seed comparisons, including shuffled controls, remain available.




Figure 9: Minibatch neural comparisons in the development patient. Three training seeds for each local, mean, GCN and GraphSAGE configuration, including self-only and node-shuffled controls; all evaluated on TENX99, a single previously used development patient. Points average three fixed masks at 20% coverage. Dashed lines give identical-mask IDW reference values. Top: image-only prediction; bottom: the protocol-specified exploratory residual-anchor baseline. Color identifies training seed, not an independent patient. Panels use different y-ranges. IDW receives measured expression unavailable to the image-only branch; residual versus IDW tests incremental morphology. These fits were not tuned on Wu or Bassiouni outcomes.

The expanded matrix compared all 48 full-section-trained fits with the 30 frozen minibatch fits in TENX99, Wu 3 and Bassiouni 22, using identical masks and information routes (Fig. 10). In the primary 20% distributed Bassiouni comparison, the residual-anchor GraphGPS-inspired model changed paired patient correlation versus IDW by −0.0504 (95% interval −0.0629 to −0.0389), relative MSE by +13.4% (11.3–18.4%) and MAE by +0.0244 (0.0204–0.0295). The latent-coordinate EGNN gave −0.0480 (−0.0698 to −0.0368), +13.0% (9.1–18.3%) and +0.0252 (0.0188–0.0310), respectively. Neither primary model improved any of these endpoints in any of the 22 patients. Under identical-H region exclusion, their median relative MSE increases were 10.2% and 9.2% (Fig. 11).

Component comparisons also failed to establish the proposed gains: the full nonlocal model worsened correlation and errors relative to its local counterpart, and latent-coordinate updates did not improve the fixed-distance model (Table 4). Same-information conventional neural comparisons, all routes, individual Wu effects, training seeds and distance-bin support are retained in Supplementary Figs. S15–S25,S28 and Tables S14–S17. Conditional geometric and reach checks confirm the implemented properties; they do not establish biological long-range association or full-image equivariance. These negative findings concern the tested training budget, frozen features and low-resolution cross-cohort inputs.




Figure 10: Graph predictors and information boundaries. The image network receives frozen image embeddings and coordinate-defined structure. The GraphGPS-inspired branch combines local gated aggregation with bounded attention to 32 coordinate-only landmarks. The alternative EGNN changes private latent coordinates through relative displacements; the distance-only model fixes them. Original physical coordinates always define masks, anchor interpolation and physical-distance diagnostics. No hidden expression is an image-network input.




Figure 11: All 16 new architectures/controls versus molecular IDW in 22 Bassiouni patients. Exploratory residual route at 20% coverage; colors distinguish historical distributed V1 and identical-H region exclusion. Bars are descriptive 95% patient percentile intervals for median paired effects after mask/training-seed averages. Positive correlation and negative relative MSE improve. The two designs have different scored support and are not subtracted here. All models are shown regardless of direction; nominally equal parameter counts do not imply equal effective function classes.

Table 4: Predefined architecture contrasts in recovered Bassiouni patients. At20% historical V1 coverage, the exploratory residual route uses identical measured anchors for both models. Effects average all three training seeds and three mask seeds within each patient before pairing; relative MSE is normalized by the reference patient mean MSE after these averages. Intervals resample22 whole patients, not genes, masks or training seeds. They are descriptive, unadjusted intervals, not simultaneous significance claims. Other routes, controls, budgets and individual Wu effects are fully exported.

Candidate vs control	Effect	Median [95% interval]
GPS full vs GPS local	Δr	-0.0050 [-0.0074, -0.0044]
GPS full vs GPS local	MSE%	+2.7228 [+2.0494, +3.6169]
GPS full vs GPS local	Δ MAE	+0.0039 [+0.0030, +0.0054]
GPS full vs GPS no distances	Δr	-0.0121 [-0.0135, -0.0088]
GPS full vs GPS no distances	MSE%	+4.9138 [+3.6647, +5.7686]
GPS full vs GPS no distances	Δ MAE	+0.0064 [+0.0056, +0.0078]
EGNN latent vs Distance fixed	Δr	-0.0048 [-0.0074, -0.0008]
EGNN latent vs Distance fixed	MSE%	+1.6673 [+0.7107, +2.8664]
EGNN latent vs Distance fixed	Δ MAE	+0.0026 [+0.0014, +0.0046]
EGNN latent vs Distance disabled	Δr	-0.0065 [-0.0096, -0.0024]
EGNN latent vs Distance disabled	MSE%	+1.9736 [+1.0100, +3.3448]
EGNN latent vs Distance disabled	Δ MAE	+0.0028 [+0.0020, +0.0053]

2.6 Regional completion, distance and calibration defined failure regimes

The original contiguous-completion protocol was substantially harder than distributed reconstruction. At 20% coverage, Bassiouni correlation fell from 0.416 to 0.171, MSE increased from 0.0923 to 0.1496 and MAE from 0.2033 to 0.2624. Every patient deteriorated; median paired correlation change was -0.267 and relative MSE increase 52.8%. Wu correlation fell from 0.359 to 0.0815, with lower correlation and higher errors in every patient (Supplementary Table S9). Because the interpolation rule and scored tissue also change, these effects must not be attributed solely to geometry. A withheld region may still lie inside the anchors' convex hull.

Distance-resolved errors were heterogeneous (Fig. 12). At 20% Bassiouni coverage, farthest-bin MSE exceeded nearest-bin MSE in 15/22 patients under distributed interpolation and 20/22 under regional completion; strictly increasing curves occurred in only 6 and 13 patients. Wu distributed curves increased across occupied bins, while regional curves were nonmonotonic. Bin support and tissue composition differ, precluding a universal reliability cutoff (Supplementary Table S7).




Figure 12: Spatial diagnostics reveal heterogeneous error beyond an overall prediction score. Development results use TENX99 and the frozen 50-gene panel. **a,** Mean gene MSE versus nearest-anchor distance for molecular interpolation at 20% coverage under distributed and development block masks; only occupied distance bins are shown. **b,** Development comparison of random and spatial farthest-point sampling, shown as MSE at matched measured-location budgets. These are descriptive single-patient sampling diagnostics, not a demonstration of universally optimal acquisition. **c,d,** Absolute-error maps for the preselected gene *FOXA1*, fixed fusion, 20% coverage and replicate 0 under random and block masks. Both maps use the same error scale, capped at the 98th percentile of the pooled preselected map-source values, which also include the archived capture-window condition; no locations are removed. Missing-region shape, available anchors and scored tissue differ between the maps. The figure illustrates why distance-bin support and spatial errors should accompany global correlation and why development geometry findings require independent evaluation.

Fixed image fusion also failed to improve external reconstruction consistently. Bassiouni correlation, MSE and MAE at 20% distributed coverage were 0.407, 0.1022 and 0.2288, versus 0.416, 0.0923 and 0.2033 for interpolation. Only 3/22 patients improved in correlation and none in either error. The median paired correlation effect was −0.0108 (95% interval −0.0187 to −0.0081), with relative MSE increase 10.7% (6.6–14.5%). Wu showed correlation improvement in 1/3 patients and no error improvement. The original anchor-versus-image paired correlation gain was 0.329 (0.304–0.438), a contrast with different information access (Table 5). HER2ST calibration reduced scale-related error but did not establish robust joint correlation/error gains (Supplementary Figs. S1,S9 and S10). An affine transform followed by normalized interpolation and its inverse left molecular predictions unchanged, as expected algebraically (Supplementary Methods).

Table 5: Patient-level uncertainty reanalysis of original Bassiouni exports. Median paired effects at 20% V1 coverage, with 10,000 patient-resampling percentile 95% intervals (seed 20261101). Negative error changes improve; positive correlation changes improve. These are descriptive exploratory intervals, not simultaneous bands or p-values. Anchors versus image is an unequal-information contrast; fusion versus anchors tests incremental image contribution. All original point estimates and source bytes are unchanged.

Contrast	Effect	Median	95% interval	Improved
Anchors minus image	Δr	0.3293	[0.3044, 0.4382]	22/22
Anchors minus image	Relative MSE (%)	-95.4534	[-96.2444, -94.0201]	22/22
Anchors minus image	Δ MAE	-1.0074	[-1.1341, -0.9108]	22/22
Fusion minus anchors	Δr	-0.0108	[-0.0187, -0.0081]	3/22
Fusion minus anchors	Relative MSE (%)	10.7078	[6.6170, 14.5048]	0/22
Fusion minus anchors	Δ MAE	0.0238	[0.0171, 0.0278]	0/22

Anchor-vector and anchor-position permutations degraded molecular reconstruction in predefined controls; image permutation supplied no comparable evidence for a useful image contribution (Supplementary Figs. S11 and S12; Table S4). These related controls and exact hidden-label invariance checks support computational traceability without supplying additional biological replicates.

3 Discussion

Measurement placement was a determinant of reconstruction quality even when the amount of observed expression was held fixed. The 22-patient comparison shows that equal budgets can produce different performance when anchors are excluded from the region containing the scored targets. Its strength is the fixed hidden set and estimator, which separate placement from changes in tissue support or reconstruction rule. The result applies to the specified central-region design; it does not establish a universally optimal sampling policy or biological correlation length.

Distributed molecular anchors nevertheless supported useful reconstruction of the predefined panel across two breast-cancer studies. The coverage response suggests that nearby same-section measurements carry information that a transported image predictor may not recover. This is a task-specific advantage of measured expression, not an equal-information demonstration that interpolation outperforms all image-based systems. Reliability should be assessed

on the intended acquisition design: favorable distributed masking does not ensure accurate completion of a large unmeasured region[24].

The neural negative results sharpen this distinction. With the same anchors supplied to residual predictors and molecular baselines, the tested graph representations did not produce a consistent external gain. Adding nonlocal access or updating latent coordinates did not establish the hypothesized advantages. Functional attention reach and conditional equivariance are software properties, rather than evidence of learned molecular relationships. These findings argue for retaining strong molecular comparators and explicit component controls, while avoiding a general rejection of geometry-aware learning on better-supported data.

Cross-cohort histology remains a substantial limitation. Public images were heavily downsampled, encoders were frozen, and fitting used only two source patients. Differences in capture depth, support and composition remain after mathematical normalization. Calibration can change image predictions, but cannot create a new interpolation benefit when a per-gene affine transform is inverted around the same normalized weighted average. Native-resolution representations, broader source training and genuinely independent evaluation would be needed to establish whether morphology adds reproducible incremental information.

The scope is also limited by retrospective masking, a development-selected 50-gene panel (40 genes in HER2ST), small Wu patient numbers and restriction to breast tissue. The exploratory controls reused evaluated cohorts. Genes, sites and repeated masks do not increase biological sample size, and descriptive patient intervals do not establish clinical generalizability. Distance alone did not define a reliable error threshold; occupied-bin support and patient-specific failures should remain visible before reconstructed expression is used downstream. Approaches such as TISSUE motivate explicit uncertainty assessment[25], but calibrated predictive uncertainty was not evaluated here.

Coverage is a count of measured locations, not a measured financial saving. If a full assay has fixed cost C_0 and per-location cost c for N locations, the conditional ratio is

$$\frac{C_{\text{full}}}{C_{20\%}} = \frac{C_0 + cN}{C_0 + 0.2cN} \leq 5. \tag{1}$$

Only a purely location-dependent component could fall fivefold at 20% coverage. Selective capture, laboratory expenditure, sequencing requirements and downstream accuracy were not prospectively tested. Future acquisition studies should evaluate placement and error requirements together with realized costs.

GraphPathology therefore supports an information-explicit account of sparse reconstruction: measured molecular support can be useful, its placement matters under controlled conditions, and added image or graph complexity requires a demonstrated incremental benefit. Prospective sampling and independent native-resolution cohorts are the next tests of practical value.

4 Methods

4.1 Data, target panel and preprocessing

All experiments used previously released paired spatial expression and histology data. The four development samples were obtained through HEST[5]: NCBI783 and NCBI785 were the

training patients, TENX95 was the validation patient and TENX99 was the primary development evaluation patient. The cohort comprised 7,015 training locations, 7,760 validation locations and 20,761 primary evaluation locations. Earlier conditional leave-one-patient-out analyses reused frozen gene panels and development-selected configurations and were interpreted as sensitivity analyses, not unbiased external cross-validation.

HER2ST[2] contributed one patient-specific section for each of A1–H1. Initial evaluation was followed by patient-disjoint calibration development using the same inspected cohort; these two analytical roles are distinguished throughout. In the Wu local Visium cohort[3], patients 1142243F, CID4465 and CID4535 supplied calibration statistics, and patients 1160920F, CID4290 and CID44971 remained held out. The Bassiouni evaluation cohort was obtained from GSE210616[4]. The source study includes 43 sections from 22 patients. We selected the first numbered section for every published patient from metadata before target evaluation, yielding BASS_P01–P09, P11–P15 and P17–P24. Gaps in identifiers reflect source numbering rather than outcome-dependent exclusion. The source study's own reference and validation groups were not used as calibration groups in this reconstruction study.

Metadata, repository identifiers, prior-use records and available asset hashes were checked to exclude reused or reprocessed development samples. Such checks support cohort independence but cannot conclusively establish patient identity across unrelated anonymized repositories. Cohort roles and sample counts are summarized in Table 1; identifiers and exact panels are supplied in Supplementary Tables S1 and S2.

The primary panel contained the 50 genes ranked using development-training expression variance; ranking did not use evaluation expression. The development assay provided 280 harmonized biological genes, and larger 250- and 280-gene settings were used in earlier development analyses. The original external studies retained all 50 genes. HER2ST retained the frozen 40-gene intersection without replacing missing genes. The optional 278/280 intersection in the Bassiouni metadata was not used to claim a secondary whole-transcriptome experiment.

For the second replication, the source filtered feature-barcode H5 files, tissue positions, scalefactor JSON files and tissue images were acquired for each selected section. Barcode joins and published in-tissue flags defined included locations. No new expression-derived quality threshold, tissue-class annotation or deconvolution output was used to select evaluation locations. Counts were checked for finite, nonnegative integer values. For count c_{ig} at location i and gene g, the evaluation target was

$$y_{ig} = \frac{\log(1 + c_{ig}) - \mu_g^{\mathrm{dev}}}{\sigma_g^{\mathrm{dev}}}, \tag{2}$$

where the centering and scaling constants were the pooled location mean and population standard deviation frozen from development-training patients; a standard deviation below 10^{-8} was replaced by 1. No destination library-size or capture-area correction was fitted. Identical target units were used for all methods within each comparison. These common mathematical units do not imply cross-platform calibration of absolute molecular abundance. Supplementary Fig. S2 summarizes cross-cohort patterns without pooling panels; the gene-level results are provided in Supplementary Table S3.

4.2 Registration and frozen image features

Visium array coordinates were mapped to image coordinates by a least-squares fit to the source hexagonal lattice with 100-μm spacing. Coordinate residuals, image scale and anisotropy were checked against the frozen qualification settings, and overlays were inspected before outcome evaluation. These tests concern location consistency and do not establish cell-level registration. Images were not super-resolved.

The Wu and Bassiouni images contained only 15/25 and 24–29 native pixels, respectively, across the nominal 112-μm patch width. Resizing does not recover morphology absent from these public inputs. Image quality was therefore recorded as a limitation of the image comparator, not as a restriction on spatial interpolation that uses only coordinates and observed expression.

At each external location, an RGB patch covered a nominal 112-μm square, rounded to the available native pixels. The patch was resized with LANCZOS to 224×224 pixels. The torchvision `IMAGENET1K_V1` preprocessing then resized the shorter side to 256 pixels using bilinear interpolation, took a central 224×224 crop, rescaled intensities to $[0, 1]$, and normalized RGB channels using means $(0.485, 0.456, 0.406)$ and standard deviations $(0.229, 0.224, 0.225)$. The second resize and central crop were retained explicitly; 112 μm denotes the nominal extracted patch, rather than the final uncropped field seen by the network. Development embeddings used the supplied 224×224 HEST patches with the same torchvision transform. No stain normalization, random image augmentation or encoder fine-tuning was applied.

The frozen ResNet18[26] comprised a 7×7 convolutional stem with stride 2, batch normalization, a rectified-linear activation and 3×3 max pooling with stride 2, followed by four residual stages. The stages contained two basic blocks each, with channel widths 64, 128, 256 and 512. Each block used two 3×3 convolutions with batch normalization and a rectified-linear activation after the first convolution and after shortcut addition. Identity or projected shortcuts matched the feature dimensions. Global average pooling produced a 512-dimensional vector; the original ImageNet classifier was replaced by the identity operation. All encoder parameters and batch-normalization statistics remained fixed during feature extraction.

The final central crop retains approximately 0.875 of the extracted patch linear field.

4.3 Coordinate-defined graph representations

Each retained location is a node with physical position $q_i \in \mathbb{R}^2$ in micrometers and a frozen image embedding. Following message-passing principles[27], graphs were constructed independently within each section from physical coordinates, without expression labels. The candidates were symmetrized six-nearest-neighbor adjacency capped at 250 μm, radius graphs at 150 or 250 μm, and Delaunay adjacency capped at 250 μm. Symmetrization takes the union of directed nearest-neighbor edges, so final degree need not equal six. The binary adjacency A had no self edges. With $\widetilde{A} = A + I$, degree matrix $\widetilde{D}$ of $\widetilde{A}$, and degree matrix D of A, the three fixed operators were

$$P = \widetilde{D}^{-1}\widetilde{A}, \qquad Q = \widetilde{D}^{-1/2}\widetilde{A}\widetilde{D}^{-1/2}, \qquad R = D^{-1}A.$$

For R, an isolated node has a zero neighbor aggregate. The input to the MLP was PX for mean pooling, QX for GCN and $[X, RX]$ for mean-GraphSAGE, where X contains standardized frozen image embeddings and brackets denote concatenation. Each predictor therefore applies one fixed graph aggregation followed by the one-hidden-layer MLP defined above. These single-step predictors used neither learned adjacency nor attention; the additional message-passing models are defined below.

Graph choice was selected at seed 0 by the highest validation median gene Pearson correlation, then lower MSE and deterministic graph-name order. All three 50-gene predictors selected capped Delaunay adjacency. Mean pooling and GCN used width 128; GraphSAGE used width 67 to approximately match the initial local head's parameter count after doubling the input dimension. With ResNet18, these models contain 72,114 and 72,075 trainable head parameters, respectively; all use dropout 0.1. Supplementary Table S8 summarizes the selected dimensions.

4.4 Neural predictors and distinct training regimes

All neural fitting used NCBI783/NCBI785 and validation-only stopping on TENX95. The following regimes are distinct; parameter similarity does not imply identical update schedules or effective capacity.

Original primary image ensemble. Each expression head was a one-hidden-layer MLP with the exact sequence

$$512 \xrightarrow{\text{linear}} 256 \xrightarrow{\text{GELU}} 256 \xrightarrow{\text{dropout (0.1)}} 256 \xrightarrow{\text{linear}} 50, \tag{3}$$

where GELU denotes the Gaussian error linear unit. Both linear layers included biases and used the default PyTorch initialization, uniform on $[-1/\sqrt{d_{\text{in}}}, 1/\sqrt{d_{\text{in}}}]$. The output was linear, with no nonnegativity constraint. Each head contained 144,178 trainable parameters. Feature-wise centering and scaling, S_x, were fitted to the pooled 7,015 training locations from NCBI783 and NCBI785. A separate per-gene scaler, S_y, was fitted to their already normalized expression targets. Both used population standard deviations, replacing scales below 10^{-8} with 1. If f_{θ_s} denotes the head trained with seed $s \in \{0, 1, 2\}$ and ϕ_{R18} the frozen encoder, the image prediction in the common target units was

$$\widehat{\boldsymbol{y}}_i^{\text{image}} = \frac{1}{3} \sum_{s=0}^{2} S_y^{-1}[f_{\theta_s}(S_x(\phi_{\text{R18}}(x_i)))] \,. \tag{4}$$

Dropout was disabled at inference. The three inverse-scaled predictions were averaged without fitting ensemble weights.

Heads minimized mean squared error in the additionally standardized target space using AdamW, learning rate 10^{-3}, weight decay 10^{-4}, $(\beta_1, \beta_2) = (0.9, 0.999)$ and $\epsilon = 10^{-8}$. Weight decay applied to all head parameters, including biases. Each epoch shuffled the training locations into minibatches of 512, retaining the final partial batch, for at most 80 epochs. Training used CPU float32 computation and no learning-rate scheduler. Early stopping used the TENX95 validation patient: an improvement required validation MSE to decrease by more than 10^{-4}, and training stopped after 10 consecutive epochs without such an improvement.

The best qualifying parameters were restored. Hidden widths $\{128, 256\}$ and dropout probabilities $\{0, 0.1\}$ were compared at seed 0; the configuration with the highest validation median gene Pearson correlation was selected, with lower validation MSE breaking ties. This configuration-selection criterion was distinct from the within-fit stopping criterion. The selected width-256, dropout-0.1 heads were fitted with seeds 0, 1 and 2; their retained epochs were 12, 11 and 12, respectively. No external patient was used to select or refit these heads.

Additional minibatch heads. Thirty additional local, mean, GCN and GraphSAGE heads (three seeds per specified configuration/control) used the following settings. They were fitted separately from the original primary ensemble and were later frozen as references. A single learned linear/GELU/dropout/readout layer followed mean or symmetric GCN aggregation; GraphSAGE concatenated self and neighbor mean. Widths were 128 for local/mean/GCN and 67 for SAGE (72,114 versus 72,075 parameters). Spatial, self-only and degree-preserving node-shuffled controls shared each family's nominal parameters, labels and training settings. SAGE self-only concatenated X with X and has a smaller effective function class. AdamW used lr 10^{-3}, decay 10^{-4}, batch size 512, dropout 0.1, maximum 80 epochs, patience 10 and minimum validation improvement 10^{-4}; TENX95 alone selected stopping. Seeds 0/1/2 were all retained. Deterministic float32 CPU execution used four torch/BLAS threads; Local, mean-self and GCN-self identity controls used the same inputs. Their float32 weight/prediction comparisons allowed rtol=atol=$32\epsilon_{\text{float32}} = 3.8147 \times 10^{-6}$; this tolerance did not relax float64 molecular replay or exact hidden-label invariance. Further identity diagnostics are in Supplementary Methods.

Full-section comparison. The 48 additional fits crossed 16 configurations with seeds 0/1/2. AdamW used learning rate 10^{-3}, weight decay 10^{-4}, maximum 80 epochs, patience 10 and minimum validation-MSE improvement 10^{-4}. Each epoch made one full-section update per source training patient in seeded random order. Local/mean/GCN widths were 128 and concatenated GraphSAGE width 67. These baselines match the new models' update budget, whereas the 30 frozen minibatch heads do not. Both additional regimes trained directly in the common development target units, without the primary ensemble's extra head-target scaler. Model selection retained every seed; TENX99 was used only for evaluation. Parameter counts, selected epochs, histories and computational costs are in Supplementary Tables S15 and S16. Initial development searches, Phikon representation comparisons[28] and conditional leave-one-patient-out fits have separate specifications in Supplementary Methods S6.

4.5 Bounded nonlocal and distance-aware message passing

This implementation draws on GraphGPS principles rather than reproducing the reference implementation. Its blocks use a 512-to-64 projection, two 64-wide blocks, four attention heads, dropout 0.1, pre-LayerNorm, local linear messages gated by eight relative-distance radial features, parallel bounded attention, residual additions and a two-layer GELU feed-forward path. The local graph is Delaunay with a 250 μm cap. Global keys/values are at most 32 deterministic coordinate-only farthest-point landmarks, giving $O(NL)$ attention storage with $L \leq 32$; there is no dense N-by-N allocation. The nonlocal-disabled model attends self plus up to seven nearest graph neighbors, padding unused slots with self. Thus full/local changes attention

range and key count as well as connectivity, despite identical 121,018 nominal parameters. Local radial features use log(1+d/250) with eight centers 0–1 and width 0.15; attention relative bias uses eight centers 0–4 and width 0.5. The distance-disabled version removes explicit gates/biases but retains coordinate-defined graph/landmark selection and is not coordinate-free.

The invariant fixed-coordinate and EGNN alternatives use two 64-wide message blocks. With centered normalized $p_i = (q_i - \bar{q})/250$, where q_i is the physical position in micrometers, $m_{ij} = \phi_e(h_i, h_j, \mathrm{RBF}(\|p_i - p_j\|^2))$, node updates aggregate the mean message through ϕ_h, using SiLU, LayerNorm and dropout 0.1. Eight radial centers span 0–4 with width 0.5. EGNN additionally updates private coordinates as

$$p_i^{l+1} = p_i^l + \frac{1}{\max(1, |\mathcal{N}_i|)} \sum_{j \in \mathcal{N}_i} 0.1 \tanh(\phi_x(m_{ij}))(p_i^l - p_j^l).$$

Each ϕ_x is a bias-free 64-to-1 map initialized at standard deviation 0.001. Only the first coordinate update affects the second layer's scalar readout; the last update has no downstream scalar loss. EGNN has 87,282 nominal parameters versus 87,154 for fixed distance (128 additional coordinate coefficients, of which the last 64 do not affect prediction). Both are E(2)-compatible conditional on frozen features, so their accuracy difference cannot identify equivariance itself. Latent positions never change original masks, anchors or IDW distances.

Self-only and coordinate/graph-shuffled variants retain nominal parameter tensors but have different effective function classes; inactive attention or edge parameters do not make them capacity-equivalent. Conjugating the graph and assigned coordinates together preserves edge-length geometry while breaking image-to-location assignment. Degree/length rewiring preserves each node's degree and global 50 μm edge-length bins below the same 250 μm cap, targeting $E/4$ accepted double swaps with at most $20E$ attempts. Actual successes are reported, including infeasible or weak rewiring. No hidden expression enters any graph, landmark or distance encoding.

4.6 Observation masks and controlled placement

For a section with N retained locations, the observed set $\mathcal{O}$ contained $\lfloor Nb + 0.5 \rfloor$ anchors at budget $b \in \{0.10, 0.20, 0.40\}$. Every hidden location lacked its entire expression vector; coordinates and image features remained available. Primary random trajectories used seeds 0–9 with nested prefixes, so higher budgets added anchors without discarding earlier observations. All trajectories were frozen without using expression values.

In distributed interpolation (V1), anchors were eligible throughout the tissue and all unobserved locations were scored. In contiguous regional completion (V2), a forbidden region consisted of the nearest 40% of locations to a center determined from the 25th, 50th or 75th percentile of the first coordinate principal-component rank. The three centers defined three fixed geometries. Anchors were sampled outside the forbidden region, and only the forbidden region was scored. All other unobserved labels were also removed from model inputs. Thus the same nominal budget refers to the full retained section even when evaluation is restricted to a region.

A deterministic spatial farthest-point policy was retained as a secondary sampling analysis. It

started from the eligible location farthest from the eligible-coordinate centroid and selected locations without expression labels. It was not optimized on external outcomes. Earlier development block and capture-window masks differ from the final regional-completion definition and are labeled separately in Fig. 3 and the source tables.

The controlled identical-target placement experiment (B2) used only the predeclared central PCA-coordinate region R (40% of locations). H was a fixed random 20% subset of R (about 8% of all locations), selected using SHA 256("H|patient") first eight bytes, little-endian. Distributed anchors were drawn from all non-H sites; region-exclusion anchors from outside R. Seeds 0–9 gave nested prefixes with equal total counts at every budget. R, H, barcodes and anchor trajectories were persisted before scoring. Both estimators scored exactly H; no expression-based geometry or exclusion was used. Seed means preceded geometry means and patient pairing. Both primary error and correlation effects and nearest-anchor distances were exported, with occupied-bin errors and support.

The additional neural matrix used three original random masks per budget for TENX99, and original V1/central-V2 geometry 1 masks with seeds 0/1/2 externally. All 78 fits also used B2 seeds 0/1/2 at 10/20/40% budgets. This subset is distinct from the ten-seed, three-geometry molecular baseline analysis.

4.7 Molecular interpolation, calibration and residual prediction

For every unmeasured location $i \in \mathcal{U}$, we define directed support edges $i \to j$ to at most 16 measured locations $j \in \mathcal{O}$, identified by the smallest coordinate distance. The resulting observation-dependent edge-weight matrix $W_{\mathcal{U}\mathcal{O}}$ has row-normalized positive weights and zero entries for unavailable anchors. The molecular reconstruction operator is

$$\widehat{Y}^{\mathrm{ST}}_{\mathcal{U}} = W_{\mathcal{U}\mathcal{O}} Y_{\mathcal{O}}. \tag{5}$$

The operator has no explicit maximum-distance cutoff, permitting cross-gap transport of observed expression. It is not a learned attention matrix, and absence of a cutoff is not evidence of reliable long-range inference. The distinct V1 and V2 weights and scoring supports are defined next.

The molecular predictor used the up to 16 nearest observed anchors, $\mathcal{N}_{16}(i;\mathcal{O})$. For distinct observed and hidden coordinates, distributed interpolation used normalized inverse-distance-squared weights, a local form of inverse-distance interpolation[29]:

$$\widehat{y}^{\mathrm{ST}}_{ig} = \sum_{j \in \mathcal{N}_{16}(i;\mathcal{O})} w_{ij} y_{jg}, \qquad w_{ij} = \frac{\|q_i - q_j\|^{-2}}{\sum_{k \in \mathcal{N}_{16}(i;\mathcal{O})} \|q_i - q_k\|^{-2}}. \tag{6}$$

For the frozen V2 protocol, the same neighborhood size was used with equal weights $w_{ij} = 1/|\mathcal{N}_{16}|$. These rules were fixed during the Wu evaluation and were not selected on Bassiouni outcomes. Fewer than 16 anchors, if available, were handled by using all available anchors. Euclidean distances were rounded to six decimal places in micrometers and floored at 10^{-6} μm before inverse-square weighting; the normalized weights therefore remained finite at coincident coordinates.

The image-only prediction was denoted $\widehat{y}^{\mathrm{image}}$. The final fusion was

$$\widehat{y}_{ig}^{\mathrm{fusion}} = 0.8\widehat{y}_{ig}^{\mathrm{ST}} + 0.2\widetilde{y}_{ig}^{\mathrm{image}}, \tag{7}$$

where the image calibration was

$$\widetilde{y}_{ig}^{\mathrm{image}} = a_g^{\mathrm{W}} + b_g^{\mathrm{W}} \frac{\widehat{y}_{ig}^{\mathrm{image}} - a_g^{\mathrm{S}}}{b_g^{\mathrm{S}}}. \tag{8}$$

Here S denotes the two development-training patients and W the three Wu calibration patients. Moments were computed from all retained targets of the designated source and calibration patients, rather than their sparse anchor subsets or image predictions. Each patient contributed equal total weight: for a domain with P patients and n_p locations in patient p,

$$a_g = \frac{1}{P}\sum_{p=1}^{P}\frac{1}{n_p}\sum_{i=1}^{n_p} y_{pig}, \qquad b_g = \sqrt{\frac{1}{P}\sum_{p=1}^{P}\frac{1}{n_p}\sum_{i=1}^{n_p}(y_{pig} - a_g)^2}. \tag{9}$$

A scale below 10^{-8} was replaced by 1. This scale includes variation about the shared domain center; it is not the mean of patient-specific standard deviations. All moments used the common target units in Eq. 2. Bassiouni evaluation reused these source and Wu constants without fitting new moments or fusion weights. The calibrated image term used in fusion therefore differs from the raw image-only comparator reported in Table 2.

Per-gene affine transformation, normalized interpolation and inverse transformation are algebraically equivalent to interpolation in the original units (Supplementary Note S1). Calibration therefore changes the image term, not the underlying normalized molecular operator.

For additional graph comparisons, g_{ig} denotes the gene-g image prediction at site i in the common target space, calibrated by Eq. 8 for both external cohorts. Alongside image-only g and fixed fusion, we evaluated observed-residual interpolation,

$$\widehat{y}_{ig} = g_{ig} + \sum_{j\in\mathcal{O}} w_{ij}\,[y_{jg} - g_{jg}]. \tag{10}$$

Only observed residual vectors enter this predictor; no regression coefficients are learned from the evaluation section. It is distinct from the development ridge residual correction, which fits gene-specific coefficients to image and molecular graph summaries using training patients (Supplementary Methods S6). Neither should be interchanged with fixed image fusion. HER2ST patient-disjoint calibration and fusion-family selection are also specified in Supplementary Methods S6; the final external image weight and Wu-derived moments remained fixed.

The matched-estimator grid used all ten original random V1 trajectories at each of 10/20/40% coverage for nearest anchor; uniform k=4,8,16,32; IDW at those k with p=1,2,3; and Gaussian k=8,16,32. We retained cKDTree ordering, six-decimal distances and a 10^{-6} μm inverse-weight floor. Gaussian bandwidth was the median observed-anchor distance to its kth other observed neighbor, with k limited by available anchors; it never used hidden expression. Every declared estimator is reported. The operator-by-protocol comparison (B1) held each original mask and scored set fixed while changing IDW k16/p2 versus uniform k16, for both V1 and all three V2 geometries.

4.8 Metrics, patient effects and uncertainty

All expression metrics were computed on hidden locations only; the full observed expression vector was available at each anchor and the full target vector was withheld at every other location. Observed rows were selected before molecular prediction, so hidden expression could not influence anchor values or interpolation weights. For each gene, Pearson and Spearman correlations summarized spatial association, while MSE and MAE summarized prediction error. Panel Pearson values were medians across genes with defined correlations. Constant-target or constant-prediction correlations were left undefined rather than assigned zero; such genes did not pass positive-correlation threshold counts. Panel MSE and MAE were means of per-gene errors. In formulae, for the scored hidden set $\mathcal{H}$,

$$\mathrm{MSE} = \frac{1}{G|\mathcal{H}|}\sum_{g=1}^{G}\sum_{i\in\mathcal{H}}(\widehat{y}_{ig}-y_{ig})^2, \qquad \mathrm{MAE} = \frac{1}{G|\mathcal{H}|}\sum_{g=1}^{G}\sum_{i\in\mathcal{H}}|\widehat{y}_{ig}-y_{ig}|. \tag{11}$$

Within each patient and geometry, metrics were averaged over mask seeds; geometry means were then averaged equally. Headline cohort values were medians of these patient summaries. Tables label an MSE value as a cohort median of patient mean MSEs, not a pooled mean over all spots. Paired effects were calculated within patients before taking their median. This distinction is necessary because a median paired effect need not equal the difference or ratio of marginal medians.

The patient was the biological unit. Seeds described computational or mask variability, not additional patients. The original external-evaluation support criterion required a median gene Pearson gain of at least 0.02 over image-only prediction, relative mean MSE reduction of at least 2%, and non-worsening MAE. The cohort rule required at least $\lceil 2N/3 \rceil$ patients meeting the joint criterion at at least two budgets, positive median benefits at every budget, and the same support threshold for correlation/MSE improvement from 10% to 40%. For 22 patients this threshold was 15. The limited native image resolution restricts interpretation of the image comparisons even when the molecular support criterion is satisfied.

For the additional neural comparisons, mask means were formed within geometry, geometry means within patient and training seeds averaged within patient before paired effects. Relative MSE is the candidate-minus-reference patient mean MSE divided by the reference mean MSE after these averages. Bassiouni intervals resampled 22 paired patients 10,000 times (seed 20261101) and took percentile 95% limits for the median effect. They are descriptive, unadjusted intervals, not p-values or simultaneous bands. Cohorts were never pooled; Wu effects are individual, and no population interval was inferred from TENX99. Raw Spearman was secondary because near-tied ranks are sensitive to floating-point differences. Full defined-correlation counts accompany the panel summaries.

For each hidden location, the nearest-anchor distance was $d_i = \min_{j\in\mathcal{O}} \|q_i - q_j\|$. Distances were grouped into the frozen bins $[0, 150)$, $[150, 300)$, $[300, 600)$, $[600, 1200)$, $[1200, 2400)$ and $[2400, \infty)$ μm. Error summaries were reported for occupied bins; the Bassiouni binwise correlations required at least 20 scored locations, while the Wu distance summaries contained errors only. Empty bins were omitted, not filled with zero. Counts, geometry and run identities were retained to distinguish sparse support from low error. Nearest-versus-farthest contrasts and monotonicity summaries were descriptive properties of the occupied-bin curves, not estimates of a universal biological length scale.

4.9 Controls and reproducibility

In the second replication, control patient BASS_P01, 20% coverage, random-mask seed 0, V1 geometry 0 and V2 geometry 1 were designated before evaluation. Three control seeds permuted observed expression-vector assignments, anchor-position assignments or image predictions. The first two disrupt related molecular–location associations; they were not interpreted as independent biological evidence. Every saved prediction was also repeated after all unobserved expression was replaced with NaNs and with independent random values. Observed/hidden barcodes, mask identities, configurations and hashes accompanied the predictions.

Analyses used Python, NumPy[30], SciPy[31], pandas, PyTorch and related scientific libraries. Reproduction instructions and environment specifications are maintained with the code. Model training and original external evaluation used recorded configurations and checksums; source-data tables provide the numerical outputs underlying the published figures.

Graphs, anchor trajectories, feature scalers and landmark selection were label-free or fitted on the designated source patients only. Replacing every unobserved label, including unscored labels, with NaNs and independent random values had to leave fixed/residual predictions exactly unchanged. Image-only B2 metrics were also identical across placements and budgets because H was fixed. Input identities and barcode/coordinate joins were checked against original records; independently derived coordinates agreed with preserved label-free coordinates within $10^{-8}\,\mu$m, which governed exact neighbor ties. Numerical source tables, environment records and fitted-state hashes are retained in the repository; the detailed audit counts are summarized in the supplementary source guide.

The original external results were frozen before the additional controls. The latter followed timestamped local protocols, retained all specified configurations/seeds and reused previously inspected patients; they were exploratory, not formally preregistered. No evaluation outcome selected a new architecture, seed, calibration map or fusion coefficient. Original primary heads and newly fitted states have separate identities and availability records.

Conditional geometric tests applied rotations, reflections, translations and node permutations at float64/float32, recording scalar and physical latent errors separately. Float 64 graph-model tolerances were 10^{-8}; float32 and preaggregated baseline tolerances were 10^{-4}. Physical latent tolerances included the 250-μm normalization. Recomputed-graph tests used nondegenerate coordinates without distance ties. These tests condition on frozen image embeddings and documented graph/landmark handling, not invariance of the complete image pipeline. Attention diagnostics recorded physical/assigned connection fractions and actual mass beyond $250/600/1200\,\mu$m per head/layer/seed/patient. A coordinate-selected remote landmark feature was zeroed to test functional dependency, not biological correlation. The degree/length control accepted no swaps in 64/66 recovered-cohort patient–seed graphs, with at most one swap in the remaining graphs; those cases supply little independent comparison.

AI-assisted computational tools. Truong-Son Hy established the scientific ideas, objectives, research direction and experimental instructions and provided intellectual oversight. Under the author's direction, OpenAI Codex assisted with coding, data processing, model training/execution and computational tests; ChatGPT assisted with drafting, editing and La-

TeX preparation. The author is responsible for evaluating the analyses, their interpretation and the conclusions. AI tools are not authors.

5 Ethics

This work reanalyzed publicly released, anonymized datasets without new recruitment, tissue collection or experimental intervention. The original studies describe their ethics and consent procedures[2–4].

6 Data availability

Development data were drawn from HEST[5]. HER2ST data are available at https://zenodo.org/records/4751624, Wu data at https://zenodo.org/records/4739739, and Bassiouni data under GEO accession GSE210616, https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE210616. The selected Bassiouni inputs have been recovered and verified against original checksums. Source tables for all manuscript figures and tables accompany the journal package. The complete additional per-gene exports (approximately 523 MB) and newly trained model-state bundle (approximately 102 MB) are currently author-local, with file/checksum manifests in the repository; no public deposit or DOI is claimed. Raw tissue images and complete prediction caches are not redistributed with the manuscript.

7 Code availability

Code, protocols and reproduction instructions are available at https://github.com/HySonLab/GraphPathology. Publication figures and documents can be regenerated from the recorded exports. Full replay of the original image predictions remains limited by three missing historical primary image-head binaries. The additional 30 and 48 fitted states are distinct from those missing binaries; their release-ready manifests specify author-local access and required deposition actions.

8 Acknowledgements

We acknowledge the investigators who generated and released the source datasets. Figures were drawn from numerical exports using Matplotlib 3.10.8 and 3.11.2 (https://matplotlib.org/). The original conceptual spatial schematic is synthetic; gene maps show measured data, anchors, predictions and errors without inferred tissue annotations.

9 Funding

No funding to disclose.

10 Author contributions

Truong-Son Hy was responsible for study conception and design, methodology, intellectual oversight, evaluation of analyses and validation, interpretation, visualization, and manuscript writing and revision. Computational and editorial tool assistance is disclosed in Methods.

11 Competing interests

Truong-Son Hy is an Editorial Board Member of *Scientific Reports*. The author declares no other competing interests.

References

[1] Ståhl, P. L. *et al.* Visualization and analysis of gene expression in tissue sections by spatial transcriptomics. *Science* **353**, 78–82 (2016). https://doi.org/10.1126/science.aaf2403.

[2] Andersson, A. *et al.* Spatial deconvolution of HER2-positive breast cancer delineates tumor-associated cell type interactions. *Nat. Commun.* **12**, 6012 (2021). https://doi.org/10.1038/s41467-021-26271-2.

[3] Wu, S. Z. *et al.* A single-cell and spatially resolved atlas of human breast cancers. *Nat. Genet.* **53**, 1334–1347 (2021). https://doi.org/10.1038/s41588-021-00911-1.

[4] Bassiouni, R. *et al.* Spatial transcriptomic analysis of a diverse patient cohort reveals a conserved architecture in triple-negative breast cancer. *Cancer Res.* **83**, 34–48 (2023). https://doi.org/10.1158/0008-5472.CAN-22-2682.

[5] Jaume, G. *et al.* HEST-1k: A dataset for spatial transcriptomics and histology image analysis. In *Advances in Neural Information Processing Systems*, vol. 37 (2024). https://doi.org/10.52202/079017-1704.

[6] He, B. *et al.* Integrating spatial gene expression and breast tumour morphology via deep learning. *Nat. Biomed. Eng.* **4**, 827–834 (2020). https://doi.org/10.1038/s41551-020-0578-x.

[7] Monjo, T., Koido, M., Nagasawa, S., Suzuki, Y. & Kamatani, Y. Efficient prediction of a spatial transcriptomics profile better characterizes breast cancer tissue sections without costly experimentation. *Sci. Rep.* **12**, 4133 (2022). https://doi.org/10.1038/s41598-022-07685-4.

[8] Zeng, Y., Wei, Z., Yu, W. *et al.* Spatial transcriptomics prediction from histology jointly through transformer and graph neural networks. *Briefings in Bioinformatics* (2022).

[9] Jia, Y., Liu, J., Chen, L., Zhao, T. & Wang, Y. Thitogene: a deep learning method for predicting spatial transcriptomics from histological images. *Briefings in Bioinformatics* **25**, bbad464 (2024).

[10] Wang, Y. *et al.* Fmh2st: foundation model-based spatial transcriptomics generation from histological images. *Nucleic Acids Research* **53**, gkaf865 (2025).

[11] Liu, J., Eckstein, M., Wang, Z., Feuerhake, F. & Merhof, D. Spatial transcriptomics expression prediction from histopathology based on cross-modal mask reconstruction and contrastive learning. *Med. Image Anal.* **108**, 103889 (2026). https://doi.org/10.1016/j.media.2025.103889.

[12] Wang, C. *et al.* Benchmarking the translational potential of spatial gene expression prediction from histology. *Nat. Commun.* **16**, 1544 (2025). https://doi.org/10.1038/s41467-025-56618-y.

[13] Kipf, T. N. & Welling, M. Semi-supervised classification with graph convolutional networks. In *International Conference on Learning Representations* (2017). https://arxiv.org/abs/1609.02907.

[14] Hamilton, W., Ying, Z. & Leskovec, J. Inductive representation learning on large graphs. In *Advances in Neural Information Processing Systems*, vol. 30 (2017). https://proceedings.neurips.cc/paper/2017/hash/5dd9db5e033da9c6fb5ba83c7a7ebea9-Abstract.html.

[15] Rampasek, L. *et al.* Recipe for a general, powerful, scalable graph transformer. In *Advances in Neural Information Processing Systems*, vol. 35, 14501–14515 (2022).

[16] Satorras, V. G., Hoogeboom, E. & Welling, M. E(n) equivariant graph neural networks. In *Proceedings of the 38th International Conference on Machine Learning*, vol. 139 of *Proceedings of Machine Learning Research*, 9323–9332 (2021).

[17] Bergenstråhle, L. *et al.* Super-resolved spatial transcriptomics by deep data fusion. *Nat. Biotechnol.* **40**, 476–479 (2022). https://doi.org/10.1038/s41587-021-01075-3.

[18] Hu, J. *et al.* Deciphering tumor ecosystems at super resolution from spatial transcriptomics with tesla. *Cell Systems* **14**, 404–417.e4 (2023).

[19] Zhang, D., Schroeder, A., Yan, H. *et al.* Inferring super-resolution tissue architecture by integrating spatial transcriptomics with histology. *Nature Biotechnology* **42**, 1372–1377 (2024).

[20] Fang, Y., Qian, J., Wang, X., Cooper, L. A. & Zhou, B. Sparser2sparse: Single-shot sparser-to-sparse learning for spatial transcriptomics imputation with natural image co-learning. *IEEE Journal of Biomedical and Health Informatics* (2026).

[21] Zhu, J., Deng, R., Guo, J. *et al.* Scr^2-st: Combine single cell with spatial transcriptomics for efficient active sampling via reinforcement learning. In *Proceedings of the 9th International Conference on Medical Imaging with Deep Learning*, vol. 315, 488–509 (PMLR, 2026). https://proceedings.mlr.press/v315/zhu26a.html.

[22] He, H.-F., Peng, P., Yang, S.-T. *et al.* Unlocking single-cell level and continuous whole-slide insights in spatial transcriptomics with panospace. *Nature Computational Science* **6**, 358–371 (2026).

[23] Xu, H., Wang, S., Fang, M. *et al.* Spacel: deep learning-based characterization of spatial transcriptome architectures. *Nature Communications* **14**, 7603 (2023).

[24] Roberts, D. R. *et al.* Cross-validation strategies for data with temporal, spatial, hierarchical, or phylogenetic structure. *Ecography* **40**, 913–929 (2017). https://doi.org/10.1111/ecog.02881.

[25] Sun, E. D., Ma, R., Navarro Negredo, P., Brunet, A. & Zou, J. TISSUE: uncertainty-calibrated prediction of single-cell spatial transcriptomics improves downstream analyses. *Nat. Methods* **21**, 444–454 (2024). https://doi.org/10.1038/s41592-024-02184-y.

[26] He, K., Zhang, X., Ren, S. & Sun, J. Deep residual learning for image recognition. In *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition*, 770–778 (2016). https://doi.org/10.1109/CVPR.2016.90.

[27] Gilmer, J., Schoenholz, S. S., Riley, P. F., Vinyals, O. & Dahl, G. E. Neural message passing for quantum chemistry. In *Proceedings of the 34th International Conference on Machine Learning*, vol. 70, 1263–1272 (PMLR, 2017). https://proceedings.mlr.press/v70/gilmer17a.html.

[28] Filiot, A. *et al.* Scaling self-supervised learning for histopathology with masked image modeling. *medRxiv* (2023). https://doi.org/10.1101/2023.07.21.23292757. Preprint.

[29] Shepard, D. A two-dimensional interpolation function for irregularly-spaced data. In *Proceedings of the 1968 23rd ACM National Conference*, 517–524 (ACM, 1968). https://doi.org/10.1145/800186.810616.

[30] Harris, C. R. *et al.* Array programming with NumPy. *Nature* **585**, 357–362 (2020). https://doi.org/10.1038/s41586-020-2649-2.

[31] Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat. Methods* **17**, 261–272 (2020). https://doi.org/10.1038/s41592-019-0686-2.

Supplementary Information

GraphPathology enables geometry-aware reconstruction of spatial transcriptomics from sparse molecular measurements

Truong-Son Hy[1,*]

[1]Department of Computer Science, University of Alabama at Birmingham, Birmingham, Alabama, United States

[*]Correspondence: thy@uab.edu

Contents and source-data guide

This document provides supporting figures and tables, cohort-resolved graph comparisons, computational diagnostics and detailed development methods. Source tables retain all tested estimators, masks, training seeds and defined-correlation counts. Patients are biological replicates; masks and training seeds describe computational variation. The main Methods distinguish the original evaluation, retrospective controls and the two additional neural training regimes.

S1 Supplementary notes

S1.1 Affine interpolation identity

For a gene with transform $T(y) = (y - \mu)/\sigma$, $\sigma > 0$, and interpolation weights satisfying $\sum_j w_{ij} = 1$, we have

$$T^{-1}\left(\sum_j w_{ij} T(y_j)\right) = \sigma \sum_j w_{ij} \frac{y_j - \mu}{\sigma} + \mu$$
$$= \sum_j w_{ij} y_j.$$

Consequently, applying and inverting a fixed per-gene affine transform does not create a distinct molecular interpolation method when weights are unchanged. In the Wu evaluation, the raw and transformed predictions differed by at most 2.66×10^{-15}, consistent with roundoff. Tiny numerical changes can break tied ranks and alter raw Spearman calculations; such changes were logged as numerical rank sensitivity, not counted as harmonization benefits. Calibration can still alter an image branch or a learned fusion model; that is a different comparison. The manuscript therefore omits duplicate molecular interpolation curves and makes no claim of an affine-harmonization advantage for molecular interpolation itself.

S1.2 Dataset roles, masking and interpretation

The four-patient HEST cohort was used for development. HER2ST was first evaluated under a frozen protocol and subsequently used for calibration development; those two views reuse the same eight patients. The Wu cohort supplied three calibration and three evaluation patients. The 22 Bassiouni patients were evaluated without new cohort-specific fitting. Counts of independent evaluation patients must not add repeated HER2ST analyses or mask replicates as new patients.

Distributed evaluation reveals molecular vectors at sampled anchors and scores all remaining locations. The external regional-completion protocol instead withholds a contiguous 40% region, samples anchors outside it, and scores that region only. V1 uses inverse-distance-squared interpolation and V2 uses an equal-weight nearest-anchor mean. Their contrast is therefore a protocol-level difference, not an isolated geometry intervention. The earlier development block/capture masks are also not identical to the external forbidden-region design.

Public external H&E images are downsampled, and sparse molecular interpolation receives same-section expression unavailable to the image-only baseline. Image-resolution limitations and unequal available information must accompany comparisons with histology. An H&E-fusion failure under these frozen inputs does not establish that native-resolution histology is generally uninformative.

S1.3 Aggregation and metric interpretation

The primary external summaries compute gene-level metrics on hidden locations, summarize genes within each run, average random seeds within geometry, average geometries within pa-

tient, and finally summarize patients. Unless a figure legend states otherwise, external cohort markers are medians across patients. Several development and transportability plots instead show arithmetic means; their legends state this explicitly. A difference between two cohort medians is not a median paired patient effect.

Mean squared error and mean absolute error use the unchanged development-standardized log1p-count target. Shared numerical units do not remove capture-depth or assay differences. Correlations are undefined when the target or prediction range is at most 10^{-12}; they are excluded from correlation averages but not counted as successes in threshold fractions. Spearman correlations use average ranks for ties. Mask and model seeds describe computational variation, not independent biological replication. Descriptive patient-bootstrap intervals are reported for the 22-patient comparisons (main Tables 3–5); they are not p-values. Wu patients are reported individually, with no primary three-patient bootstrap claim.

S2 Supplementary figures




Figure S1: HER2ST transportability and target calibration. Eight HER2ST patients and the frozen 40-gene intersection are shown at 20% random measured-location coverage. **a,c,** Distributed interpolation (V1). **b,d,** Contiguous regional completion (V2). Panels a/b show median gene Pearson correlation and c/d show mean gene MSE in common target units on a logarithmic axis. Dots are patient summaries after averaging seeds and geometries; connected markers are medians across patients. T0 denotes the uncalibrated zero-shot analysis, whereas T1 denotes the calibrated target mapping with fixed fusion. Image weights were 0.25 for V1 and 0.5 for V2; these development weights differ from the final external weight of 0.2. These are the same patients, not independent replication groups. The T1 fixed-fusion analysis differs from the later fold-selected fusion results in Supplementary Fig. S10. Affine transformation and inverse transformation leave molecular interpolation unchanged; apparent image/fusion calibration effects are not a new interpolation advantage.




Figure S2: Cross-cohort synthesis without pooling different panels. Rows correspond to the development primary patient ($n = 1$), HER2ST zero-shot ($n = 8$), HER2ST calibrated ($n = 8$, the same patients), original Wu evaluation ($n = 3$) and original Bassiouni evaluation ($n = 22$). The development and original evaluation cohorts use 50 genes; HER2ST uses 40. **a–e,** Left-column paired patient correlation effects at 20% distributed coverage for molecular interpolation minus image-only prediction and fusion minus molecular interpolation; bars are median paired effects. **f–j,** Right-column individual patient molecular correlation curves at 10%, 20% and 40% distributed coverage. The panels preserve differences in analytical role and gene panel and do not define a pooled cross-cohort score. HER2ST calibration is development, not an additional independent validation cohort.




Figure S3: Morphology-only development baselines and controls. The morphology-only development analyses use the single primary development patient TENX99 and 50 genes. **a,** Median gene Pearson correlation for ridge regression (B1) and the small multilayer perceptron (B2). The training-mean baseline has constant predictions and undefined correlations and is not drawn as zero. **b,** MSE under training-target-row permutation (NC1) and training-embedding-row permutation (NC2). Dots represent saved model-seed runs and bars are arithmetic means of displayed runs, not independent patient estimates. The controls disrupt image–expression pairing while preserving validation and evaluation data. This initial image-only head differs from the frozen local MLP selected in the representation comparison and used for external replication.




Figure S4: Single-scale graph context did not by itself establish spatial specificity. The single-scale graph comparison uses TENX99 and the 50-gene panel. **a,** Median gene correlation for the frozen local MLP, neighborhood mean pooling plus MLP, GCN and GraphSAGE. **b,** Mean gene MSE for the archived shuffled-neighborhood runs. Dots are training-seed runs and bars are their arithmetic means. A graph model outperforming a local predictor does not establish that the correct spatial adjacency caused the gain; generic aggregation or model parameterization can also contribute. The results use the original fitted models without additional graph selection or local-predictor refitting.




Figure S5: Matched graph controls limited claims about exact topology. The matched graph controls use one development patient, TENX99, and 50 genes. **a,** Median gene Pearson correlation. **b,** Mean gene MSE. Local, self-only, mean-pooling and true-spatial GraphSAGE results accompany vertex-permuted, degree-rewired and degree-plus-edge-length-rewired controls. Training seeds are averaged within each graph randomization seed; the randomized controls have five graph seeds. Dots are these averages and bars are arithmetic means. Random graph seeds are not biological replicates. The experiment did not establish a reproducible advantage of the true spatial topology. This figure records a negative/mixed development result, not evidence that topology is irrelevant in every representation or tissue.




Figure S6: Pathology representation effects in the development patient. The local MLP representation comparison evaluates frozen ImageNet ResNet18 and Phikon representations in TENX99 for 50 genes. **a,** Median gene Pearson correlation. **b,** Mean gene MSE. Each dot is one of three training seeds; bars are arithmetic means. The panel-specific improvement did not establish general superiority on larger target panels or reproducible topology-specific gains. Phikon is not introduced into the final Bassiouni replication; that analysis retains the frozen ResNet18 image branch. These are development comparisons, not external encoder validation.




Figure S7: Sparse-reconstruction development controls. The sparse-reconstruction controls use the ResNet18 branch, TENX99, 50 genes and 20% measured-location coverage. In this figure, B1 denotes image-only prediction, B3 image–molecular fusion and B4 learned propagation. Controls include anchor-expression permutation (NC1), anchor-position permutation (NC2) and image-location permutation (NC3). Model labels are specific to this experiment. **a,** Median gene Pearson correlation. **b,** Mean gene MSE. All controls use the same random observation mask (replicate 0); dots represent three permutation seeds and bars are arithmetic means. These distributions are control diagnostics rather than estimates of a geometry effect. Separate integrity checks establish invariance to corruption of hidden labels. Related anchor-assignment disruptions must not be counted as independent biological mechanisms.




Figure S8: Sampling effects depended on the reconstruction regime. The sampling comparison uses interpolation-only results for TENX99, the ResNet18 reference branch, 50 genes and 20% coverage. **a,** Open-tissue interpolation. **b,** Contiguous missing-region evaluation. A0 is random sampling with ten seeds; A1 is spatial farthest-point sampling, A2 morphology-space farthest-point sampling, and A3 a locked joint policy. A1–A3 are deterministic given their stored initialization. Dots are policy runs and bars are arithmetic means. This limited baseline diagnostic is not a complete summary of all learned/fusion families and does not demonstrate universally optimal sampling.




Figure S9: HER2ST patient distributions under the frozen zero-shot protocol. The eight HER2ST patients, 40 shared primary genes and 20% random-anchor coverage are summarized separately for **a,** distributed interpolation and **b,** contiguous regional completion. E0 denotes H&E-only prediction, E1 molecular interpolation and E2 frozen fusion, with image weights 0.25 for V1 and 0.5 for V2. Dots are patient means across seeds/geometries. Bars are arithmetic patient means, unlike the median summaries in Supplementary Fig. S1 and the primary tables. These zero-shot results and the subsequent calibration analyses use the same patients; HER2ST therefore does not supply an untouched final validation cohort.




Figure S10: Patient-disjoint calibration did not establish robust fusion superiority. The calibration-development analysis includes eight HER2ST patients, the frozen 40-gene intersection and 20% coverage. **a,** Selected calibrated fusion minus molecular interpolation in Pearson correlation. **b,** Relative MSE reduction compared with molecular interpolation; positive values indicate lower fusion error. Dots are patient/fold results grouped by V1/V2 and bars are arithmetic means. The selected fusion rule was determined within patient-disjoint training/calibration folds, not from the displayed outer-patient outcomes. Joint correlation and error improvements were not consistent across patients. These selected-fusion results should not be interchanged with the T1 fixed-fusion target study shown in Supplementary Fig. S1.




Figure S11: Matched negative controls in original Wu evaluation. The predeclared held-out patient 1160920F, 50 genes, 20% coverage and random mask seed 0 were used, with V1 geometry 0 and V2 geometry 1. Three permutation seeds were evaluated for NC1 anchor-vector reassignment, NC2 anchor-position reassignment and NC3 image-prediction permutation. **a,** Control minus matched-clean median gene Pearson correlation. **b,** Matched-clean minus control MSE, so negative values indicate increased control error. Molecular interpolation (M1) and fixed image fusion (M3) are shown. Dots are matched control settings and bars are means; they are not independent patients. Molecular controls disrupt the correct spatial anchor assignments, whereas image permutation provides no robust supporting evidence for the frozen image contribution.




Figure S12: Replication controls and native image resolution. The independent-replication controls use the predeclared Bassiouni patient BASS_P01, 50 genes, 20% coverage, random mask seed 0, V1 geometry 0 and V2 geometry 1. **a,b,** Matched Pearson and MSE effects for M1/M3 under three control seeds, with definitions and signs as in Supplementary Fig. S11. Dots are matched control settings and bars are arithmetic means. **c,** Native pixels per nominal 112-μm patch for all 22 patients in source order. The supplied images provide only 24–29 native pixels before resizing for the encoder. This is image-input quality control, not expression-prediction performance. It explains why the image comparisons cannot establish the limits of native-resolution histopathology. Coordinate overlays for all patients are recorded in the archived qualification audit.

S3 Supplementary tables

The tables below provide compact views of the accompanying machine-readable data and definitions needed to interpret the supplementary figures. Table S2 is rounded for presentation; the digital files retain unrounded exported values. Large gene-level and distance-resolved tables are supplied as separate files rather than reproduced in full in this PDF. Mask, model and permutation seeds are computational replicates and do not increase the number of independent patients.

Table S1: Primary gene panel. The 50 genes are listed in the order fixed by variance ranking in the development-training patients. The original Wu and Bassiouni evaluation retain all 50 genes. HER2ST analyses use the 40-gene intersection with this panel; unavailable genes are not replaced. The full data file also includes the larger development panels and cohort-specific availability records: `source_data/SuppTable1_gene_panels.csv`.

Order	Gene	Order	Gene
1	FOXA1	26	LUM
2	SCD	27	MYO5B
3	KRT7	28	SFRP4
4	ERBB2	29	KRT14
5	POSTN	30	TOP2A
6	CEACAM6	31	IL7R
7	SERHL2	32	AR
8	ANKRD30A	33	KRT5
9	EPCAM	34	PCLAF
10	ABCC11	35	KLF5
11	TACSTD2	36	SFRP1
12	GATA3	37	TRAF4
13	FASN	38	CXCR4
14	MLPH	39	MYH11
15	S100A14	40	TENT5C
16	CDH1	41	TCIM
17	CD9	42	NARS
18	KRT8	43	ACTG2
19	CCND1	44	PDK4
20	DSP	45	CD3E
21	SERPINA3	46	TRAC
22	LYPD3	47	JUP
23	TFAP2A	48	LYZ
24	TPD52	49	CENPF
25	PTGDS	50	CCDC80

Table S2: Patient-resolved reconstruction metrics. Compact view of the 25 original external-evaluation patients at 20% distributed coverage, using 50 genes. Values average mask seeds within patient. r is the median gene Pearson correlation; MSE is mean gene squared error in the common target units. I denotes image-only, ST molecular interpolation, and F fixed image fusion. Complete exported patient summaries for the available cohorts, regimes, budgets and methods are supplied in `source_data/SuppTable2_per_patient.csv`.

Patient	r_{I}	r_{ST}	r_{F}	$\mathrm{MSE}_{\mathrm{I}}$	$\mathrm{MSE}_{\mathrm{ST}}$	$\mathrm{MSE}_{\mathrm{F}}$
Wu						
1160920F	-0.045	0.359	0.335	0.846	0.082	0.095
CID4290	0.161	0.325	0.333	2.915	0.099	0.100
CID44971	0.059	0.502	0.496	2.949	0.115	0.118
Bassiouni						
BASS_P01	0.077	0.382	0.370	1.304	0.087	0.106
BASS_P02	0.009	0.459	0.431	1.067	0.093	0.127
BASS_P03	0.030	0.545	0.537	2.585	0.081	0.089
BASS_P04	-0.039	0.334	0.312	1.882	0.113	0.125
BASS_P05	0.096	0.636	0.628	1.581	0.133	0.155
BASS_P06	-0.005	0.312	0.297	1.567	0.069	0.072
BASS_P07	0.013	0.191	0.180	2.684	0.092	0.097
BASS_P08	0.095	0.403	0.393	2.120	0.088	0.095
BASS_P09	0.075	0.391	0.380	1.763	0.120	0.139
BASS_P11	-0.081	0.474	0.464	2.052	0.104	0.109
BASS_P12	0.182	0.435	0.436	1.748	0.113	0.120
BASS_P13	0.151	0.612	0.615	2.228	0.104	0.119
BASS_P14	0.041	0.540	0.536	1.458	0.126	0.154
BASS_P15	0.093	0.386	0.389	1.777	0.101	0.115
BASS_P17	0.095	0.428	0.421	1.773	0.088	0.096
BASS_P18	0.090	0.516	0.496	2.391	0.083	0.086
BASS_P19	0.068	0.537	0.501	2.431	0.068	0.078
BASS_P20	0.067	0.326	0.319	2.407	0.096	0.098
BASS_P21	0.075	0.401	0.340	2.607	0.092	0.108
BASS_P22	-0.029	0.322	0.300	2.149	0.087	0.090
BASS_P23	0.267	0.541	0.525	2.166	0.054	0.061
BASS_P24	0.106	0.350	0.302	1.626	0.048	0.052

Table S3: Per-gene results and metric definitions. The accompanying digital table, `source_data/SuppTable3_per_gene.csv.gz`, contains the exported Wu and Bassiouni gene-level results, with run, patient, panel, mask and method identifiers. Compression changes storage only. Every metric is calculated using the hidden locations scored by the corresponding protocol. The definitions below explain the gene-level statistics and their summaries; genes and repeated runs are not independent patient observations.

Quantity	Definition and interpretation
Pearson correlation	Correlation between predicted and measured values across scored locations, calculated separately for each gene. Constant targets or predictions yield undefined correlations.
Spearman correlation	Correlation between the ranks of predicted and measured values across scored locations, calculated separately for each gene. Numerical changes in tied ranks do not establish a biological improvement.
Mean squared error (MSE)	Mean squared prediction error across scored locations for each gene, evaluated in the common development-standardized log1p-count units.
Mean absolute error (MAE)	Mean absolute prediction error across scored locations for each gene, evaluated in the same target units.
Panel correlation	Median of defined gene correlations. Undefined correlations are excluded from this median and do not count as positive-correlation successes.
Panel error	Arithmetic mean of the gene-level MSEs or MAEs. All genes are retained in error summaries.
Patient and cohort summaries	Mask seeds are averaged within geometry and geometries within patient. Headline cohort values are medians of patient summaries; paired effects are calculated within patients before summarizing them.

Table S4: Negative-control families. Control definitions are experiment-specific and are given in the corresponding figure legends. Training-row controls differ from inference-time controls that permute permitted sparse inputs. Permuting anchor-expression assignments and anchor-position assignments disrupts related associations and does not constitute independent biological evidence. Exported control results are supplied in `source_data/SuppTable4_negative_controls.csv`.

Control family	Interpretation
Training-target or training-embedding row permutation	Disrupts local morphology–target pairing during development while preserving the validation and evaluation data.
Anchor-expression assignment permutation	Reassigns observed expression vectors among anchors, disrupting their correspondence with spatial position.
Anchor-position assignment permutation	Disrupts the spatial assignment of observed anchors while retaining the permitted molecular information.
Image-location or image-prediction permutation	Tests dependence on the image input while retaining the allowed molecular measurements.

Table S5: Recorded integrity and replay checks. Counts are taken from the original Wu and Bassiouni technical audits; these audits were not rerun during manuscript preparation. Method predictions and metric arrays are distinct audit units. For each prediction, hidden labels were independently replaced by NaNs and random values to test that unobserved expression could not influence model inputs. Digital records: `source_data/SuppTable5_integrity.csv`.

Audit quantity	Wu	Bassiouni
Mask trajectories	132	968
Clean run bundles	396	2,904
Control run bundles	18	18
Method prediction replays	1,656	8,766
Metric array replays	6,624	35,064
Hidden-label corruption checks	3,312	17,532

Table S6: Observation masks and sampling protocols. Coverage is the proportion of retained locations with observed complete expression vectors, not the proportion of tissue area assayed. All unobserved labels are excluded from model inputs. The three regional geometries and ten random-mask seeds are computational replicates. The accompanying `source_data/SuppTable6_masks.csv` records the exported mask identities, patients, regimes, geometries and sampling trajectories.

Protocol component	Definition
Measurement budgets	For N retained locations and $b \in \{0.10, 0.20, 0.40\}$, the observed count is $\lfloor bN + 0.5 \rfloor$. Random anchor sets use nested prefixes of the same trajectory as coverage increases.
Distributed interpolation (V1)	Anchors may be sampled throughout the retained tissue; every remaining location is scored. Predictions use up to 16 nearest observed anchors with normalized inverse-distance-squared weights.
Contiguous regional completion (V2)	A prespecified contiguous region contains the 40% of locations nearest to its center. Anchors are sampled outside that region, and only locations within it are scored. Predictions use the mean of up to 16 nearest observed anchors.
Regional geometry	Three centers are selected at the 25th, 50th and 75th percentile of first coordinate principal-component rank. The three geometries are averaged equally within patient.
Secondary spatial sampling	Label-free farthest-point sampling starts from the eligible location farthest from the eligible-coordinate centroid. It is not optimized using external expression outcomes.
Development masks	Earlier block and capture-window masks differ from the external regional-completion protocol and are interpreted separately.

Table S7: Distance-resolved evaluation and patient support. Distance is measured from each scored hidden location to its nearest observed anchor. Empty bins are omitted. Bassiouni binwise Pearson correlations require at least 20 scored locations in the corresponding run; Wu distance bins report MSE and MAE only. Error summaries may include smaller occupied bins. The patient counts below summarize the Bassiouni cohort at 20% coverage and do not define a universal distance cutoff. Exported bin-level metrics and support counts are supplied in `source_data/SuppTable7_distance_bins.csv`.

Evaluation component	Definition
Distance bins (μm)	$[0, 150)$, $[150, 300)$, $[300, 600)$, $[600, 1200)$, $[1200, 2400)$ and $[2400, \infty)$.
Bin support	Number of scored hidden locations contributing to the patient, method, mask and distance-bin summary; occupied-bin support may differ across patients and protocols.
Nearest-to-farthest comparison	Compares MSE in the nearest and farthest occupied bins within each patient.
Monotonicity	Assesses whether MSE strictly increases across the occupied distance bins.

Bassiouni patient summary	V1	V2
Higher MSE in farthest than nearest occupied bin	15/22	20/22
Strictly increasing occupied-bin MSE	6/22	13/22

Table S8: Selected supporting predictor architectures for the 50-gene panel. Dimensions denote input → hidden → output, and every listed MLP has GELU, dropout 0.1 and a linear output layer. Encoder parameters remain frozen; counts include trainable head biases. All MLPs use the shared optimizer and stopping rule above. The initial local predictor and the representation-comparison predictor are distinct fitted heads.

Predictor	Selected dimensions	Fitting settings or fixed aggregation
Initial ResNet18 ridge	$512 \to 50$	Ridge penalty 100; intercept; 25,650 coefficients.
Initial ResNet18 MLP	$512 \to 128 \to 50$	Lowest validation MSE; 72,114 head parameters.
ResNet18 mean pooling / GCN	$512 \to 128 \to 50$	PX or QX on capped Delaunay adjacency; 72,114 head parameters.
ResNet18 mean-GraphSAGE	$1024 \to 67 \to 50$	$[X, RX]$ on capped Delaunay adjacency; 72,075 head parameters.
ResNet18 representation-comparison MLP	$512 \to 256 \to 50$	Highest validation median gene Pearson; 144,178 head parameters.
Phikon local MLP	$768 \to 256 \to 50$	Same candidate grid and selection criterion; 209,714 head parameters.
Phikon spatial MLPs	$768 \to 128 \to 50$; $1536 \to 67 \to 50$	Mean pooling and mean-GraphSAGE, respectively; inherited graph and head widths.
Sparse residual correction	34 covariates plus intercept per gene	Gene-specific ridge correction of fixed fusion; penalty 10,000; 1,750 coefficients for 50 genes.

Table S9: Coverage response and contiguous-region completion in the two original evaluation studies. All rows use the same frozen 50-gene panel and spatial interpolation. Values are medians of patient summaries, with $n = 3$ for Wu and $n = 22$ for Bassiouni. V1 is distributed interpolation with inverse-distance-squared weighting; V2 is completion of a predefined contiguous region using the mean of nearby anchors. Both the scoring support and interpolation rule change between V1 and V2. The comparison therefore describes protocol-level deterioration rather than an isolated causal geometry effect. In both studies, every evaluation patient improved in Pearson correlation and MSE from 10% to 40% V1 coverage; every patient deteriorated in Pearson, MSE and MAE under V2 relative to V1 at 20%.

Cohort	Coverage	V1 Pearson r	V1 MSE	V2 Pearson r	V2 MSE
Wu, original evaluation	10%	0.313	0.1080	0.111	0.1356
	20%	0.359	0.0987	0.081	0.1366
	40%	0.393	0.0899	0.088	0.1385
Bassiouni, original evaluation	10%	0.380	0.1001	0.121	0.1520
	20%	0.416	0.0923	0.171	0.1496
	40%	0.465	0.0852	0.181	0.1518

S4 Additional controls and source-data guide

The matched molecular and neural comparisons use previously evaluated cohorts. Complete source tables are under `source_data/journal/` and `source_data/extension/`; protocols, input identities, fitted-state records and numerical audits are documented in the repository. Exact Bassiouni inputs are available and verified. The complete additional per-gene exports and model states remain author-local with release-ready manifests, not publicly deposited assets. Publication regeneration from exported tables is distinct from replay of the three missing historical primary image heads.

Table S10: All 20 matched-information estimators in Wu at 20% V1 coverage. Each patient value averages ten original random trajectories. P1=1160920F, P2=CID4290, P3=CID44971. Relative MSE is candidate minus fixed IDW k16/p2, divided by that reference; negative values improve. No external winner was selected. All 10/20/40% results, MAE and individual paired effects accompany the CSV exports. Gaussian bandwidth uses observed-anchor geometry only.

Estimator	r P1	r P2	r P3	MSE% P1	MSE% P2	MSE% P3
nearest	0.223	0.228	0.368	+68.3	+61.9	+63.7
uniform_k4	0.328	0.293	0.474	+8.0	+8.2	+7.5
uniform_k8	0.354	0.294	0.475	+0.7	+4.6	+4.0
uniform_k16	0.359	0.265	0.442	-0.8	+7.5	+6.8
uniform_k32	0.348	0.220	0.396	+0.8	+13.7	+11.7
idw_k4_p1	0.327	0.305	0.483	+8.5	+6.6	+6.4
idw_k4_p2	0.313	0.299	0.475	+13.1	+9.7	+10.2
idw_k4_p3	0.297	0.289	0.460	+19.8	+15.3	+16.3
idw_k8_p1	0.360	0.321	0.499	-0.0	+0.8	+0.3
idw_k8_p2	0.341	0.317	0.495	+4.4	+2.8	+3.0
idw_k8_p3	0.315	0.301	0.474	+12.7	+9.4	+10.4
idw_k16_p1	0.378	0.318	0.497	-3.1	+0.7	+0.0
idw_k16_p2	0.359	0.325	0.502	+0.0	+0.0	+0.0
idw_k16_p3	0.324	0.305	0.479	+9.1	+6.6	+7.6
idw_k32_p1	0.387	0.307	0.480	-3.3	+3.8	+2.1
idw_k32_p2	0.371	0.328	0.503	-2.2	-0.8	-1.2
idw_k32_p3	0.329	0.308	0.481	+7.2	+5.2	+6.3
gaussian_k8	0.360	0.308	0.489	-0.1	+2.6	+2.0
gaussian_k16	0.374	0.286	0.461	-1.9	+4.8	+4.4
gaussian_k32	0.358	0.249	0.417	-0.5	+10.3	+9.5

Table S11: Operator by historical protocol at 20% coverage. IDW and uniform operators share each original mask and scored set. V2 averages geometries equally after averaging seeds. The V1/V2 contrast still changes spatial support and is not a causal geometry estimate. All budgets are supplied in source data.

Patient	Protocol	IDW r	IDW MSE	Uniform r	Uniform MSE
1160920F	V1	0.359	0.0817	0.359	0.0811
1160920F	V2	0.179	0.1029	0.176	0.1028
CID4290	V1	0.325	0.0987	0.265	0.1061
CID4290	V2	0.050	0.1367	0.036	0.1366
CID44971	V1	0.502	0.1150	0.442	0.1228
CID44971	V2	0.113	0.1781	0.081	0.1767

Table S12: Identical-target anchor-placement effects at 20% coverage in Wu. Region-exclusion minus distributed anchors, with the same H and total anchor count. Each row averages ten seeds. The holdout H is fixed across conditions and budgets. All three patients deteriorate in Pearson, MSE and MAE with both estimators. This central-region result is exploratory robustness evidence in previously inspected patients, not new blind validation.

Patient	Operator	Δr	Relative MSE%	Δ MAE
1160920F	IDW	-0.1056	+20.1	+0.0214
1160920F	Uniform	-0.0928	+21.7	+0.0196
CID4290	IDW	-0.2452	+45.4	+0.0464
CID4290	Uniform	-0.1875	+36.0	+0.0361
CID44971	IDW	-0.1274	+37.3	+0.0386
CID44971	Uniform	-0.1605	+38.9	+0.0387

Table S13: All new controlled neural fits at 20% TENX99 coverage. Each value averages three identical original masks. Models fit NCBI783/NCBI785 and use TENX95 alone for stopping. Seeds are training variability, not patients. Residual is the exploratory image plus interpolated observed-residual predictor. All budgets, fixed 20% fusion and anchor-only values are in CSVs. No architecture or seed was selected using TENX99. Nominal parameter counts are 72,114 (local/mean/GCN) and 72,075 (SAGE); SAGE self-only duplicates self inputs and has a smaller effective function class.

Configuration	Seed	Image r	Image MSE	Residual r	Residual MSE
local local	0	0.666	0.9142	0.818	0.2576
local local	1	0.686	0.9008	0.824	0.2569
local local	2	0.695	0.9191	0.821	0.2555
MEAN spatial	0	0.651	0.8825	0.812	0.2458
MEAN spatial	1	0.647	0.9095	0.813	0.2497
MEAN spatial	2	0.658	0.9206	0.813	0.2446
MEAN self	0	0.666	0.9142	0.818	0.2576
MEAN self	1	0.686	0.9008	0.824	0.2569
MEAN self	2	0.695	0.9191	0.821	0.2555
MEAN shuffled	0	0.117	1.2311	0.787	0.2747
MEAN shuffled	1	0.227	1.1425	0.771	0.2865
MEAN shuffled	2	0.213	1.1819	0.778	0.2889
GCN spatial	0	0.652	0.8871	0.809	0.2508
GCN spatial	1	0.659	0.8885	0.812	0.2518
GCN spatial	2	0.653	0.9240	0.809	0.2503
GCN self	0	0.666	0.9142	0.818	0.2576
GCN self	1	0.686	0.9008	0.824	0.2569
GCN self	2	0.695	0.9191	0.821	0.2555
GCN shuffled	0	0.124	1.2297	0.788	0.2743
GCN shuffled	1	0.230	1.1264	0.767	0.2909
GCN shuffled	2	0.228	1.1830	0.777	0.2908
SAGE spatial	0	0.667	0.8611	0.823	0.2340
SAGE spatial	1	0.688	0.8792	0.831	0.2346
SAGE spatial	2	0.679	0.9173	0.822	0.2329
SAGE self	0	0.667	0.9364	0.815	0.2574
SAGE self	1	0.656	0.9010	0.812	0.2590
SAGE self	2	0.672	0.9599	0.810	0.2582
SAGE shuffled	0	0.686	0.8615	0.812	0.2511
SAGE shuffled	1	0.695	0.8655	0.816	0.2564
SAGE shuffled	2	0.684	0.9124	0.818	0.2403




Figure S13: Predetermined real Wu V1 gene maps. Patient 1160920F, 20% random seed0, original mask, fixed IDW k16/p2. Genes ERBB2, EPCAM, FOXA1 and POSTN were designated before comparisons. Truth/prediction use shared per-gene scales over scored sites; errors have separate scales. Anchors are crosses and gray sites are other registered locations. Standardized log1p counts are not absolute transcript abundance. Constant scored fields are labelled rather than omitted or given misleading contrast. Source: Wu et al., Zenodo 4739739, CC-BY4.0; no tumor/cell annotations are inferred.




Figure S14: Predetermined real Wu V2 gene maps. Same patient, genes, 20% budget and seed0, central original V2 geometry 1, historical uniform k16 operator. Colored truth/prediction/error points are the original forbidden scoring region; anchor crosses lie outside. Scales and source terms match the preceding figure. FOXA1 has a constant scored field; numerical residuals do not establish biological recovery. These maps retain unsuccessful/constant examples and are not independent validation. The corresponding Bassiouni maps are provided in Supplementary Figs. S26 and S27.

S5 Graph comparisons and computational diagnostics

S5.1 Complete cohort and prediction-route matrices

All matrices use 20% distributed coverage and the fixed 50-gene panel. Sixteen full-section configurations and ten frozen minibatch configuration families are shown, each retaining three training seeds, alongside fixed IDW and uniform molecular estimators. Metrics are averaged over the same three mask seeds and then training seeds within patient. Dots are patient summaries, and black ticks are cohort medians; their spread is not a confidence interval. The image-only route uses frozen image features and permitted coordinates; fixed fusion and observed-residual interpolation additionally use the same measured anchors as the molecular references. Thus image-only versus molecular rows have different information access. Architecture names identify configurations, not outcome-selected winners.




Figure S15: HEST development: image-only at 20% distributed coverage. Results for one previously evaluated patient, TENX99. Pearson, MSE and MAE are shown after averaging three mask and three training seeds within patient. Neural rows use image features and permitted geometry; the two molecular rows receive measured anchors. The dot represents one patient; no population interval is inferred.




Figure S16: HEST development: fixed fusion at 20% distributed coverage. Results for one previously evaluated patient, TENX99. Pearson, MSE and MAE are shown after averaging three mask and three training seeds within patient. Neural rows combine 0.2 calibrated image prediction with 0.8 IDW; molecular rows use the same anchor vectors without images. The dot represents one patient; no population interval is inferred.




Figure S17: HEST development: observed-residual interpolation at 20% distributed coverage. Results for one previously evaluated patient, TENX99. Pearson, MSE and MAE are shown after averaging three mask and three training seeds within patient. Neural rows add interpolation of observed residuals to image prediction; molecular rows use the same anchors without images. The dot represents one patient; no population interval is inferred.




Figure S18: Wu: image-only at 20% distributed coverage. Results for three previously evaluated patients. Pearson, MSE and MAE are shown after averaging three mask and three training seeds within patient. Neural rows use image features and permitted geometry; the two molecular rows receive measured anchors. Colors identify 1160920F, CID4290 and CID44971 individually.




Figure S19: Wu: fixed fusion at 20% distributed coverage. Results for three previously evaluated patients. Pearson, MSE and MAE are shown after averaging three mask and three training seeds within patient. Neural rows combine 0.2 calibrated image prediction with 0.8 IDW; molecular rows use the same anchor vectors without images. Colors identify 1160920F, CID4290 and CID44971 individually.




Figure S20: Wu: observed-residual interpolation at 20% distributed coverage. Results for three previously evaluated patients. Pearson, MSE and MAE are shown after averaging three mask and three training seeds within patient. Neural rows add interpolation of observed residuals to image prediction; molecular rows use the same anchors without images. Colors identify 1160920F, CID4290 and CID44971 individually.




Figure S21: Bassiouni: image-only at 20% distributed coverage. Results for 22 previously evaluated patients. Pearson, MSE and MAE are shown after averaging three mask and three training seeds within patient. Neural rows use image features and permitted geometry; the two molecular rows receive measured anchors. Gray dots identify individual patients; black ticks are medians of their summaries.




Figure S22: Bassiouni: fixed fusion at 20% distributed coverage. Results for 22 previously evaluated patients. Pearson, MSE and MAE are shown after averaging three mask and three training seeds within patient. Neural rows combine 0.2 calibrated image prediction with 0.8 IDW; molecular rows use the same anchor vectors without images. Gray dots identify individual patients; black ticks are medians of their summaries.




Figure S23: Bassiouni: observed-residual interpolation at 20% distributed coverage. Results for 22 previously evaluated patients. Pearson, MSE and MAE are shown after averaging three mask and three training seeds within patient. Neural rows add interpolation of observed residuals to image prediction; molecular rows use the same anchors without images. Gray dots identify individual patients; black ticks are medians of their summaries.

Table S14: All 384 saved-checkpoint conditional geometric checks. Maxima across all three seeds and recorded transforms. Physical latent errors are in micrometers. Tests are conditional on fixed image features, not the complete CNN pipeline. Baseline preaggregation remains float 32 even when the head is tested in float 64.

Model	Precision	Max scalar error	Max latent error	Checks
Distance disabled	float 64	0	1.14×10^{-13}	12
Distance disabled	float 32	7.75×10^{-7}	2.40×10^{-5}	12
Distance fixed	float 64	1.11×10^{-15}	1.14×10^{-13}	12
Distance fixed	float 32	8.42×10^{-7}	2.40×10^{-5}	12
EGNN rewired	float 64	8.88×10^{-16}	1.14×10^{-13}	12
EGNN rewired	float 32	7.15×10^{-7}	3.13×10^{-5}	12
EGNN latent	float 64	1.33×10^{-15}	1.14×10^{-13}	12
EGNN latent	float 32	7.45×10^{-7}	2.86×10^{-5}	12
EGNN self	float 64	0	1.14×10^{-13}	12
EGNN self	float 32	7.15×10^{-7}	2.40×10^{-5}	12
EGNN shuffled	float 64	1.33×10^{-15}	1.14×10^{-13}	12
EGNN shuffled	float 32	7.45×10^{-7}	2.70×10^{-5}	12
GPS rewired	float 64	1.33×10^{-15}	1.14×10^{-13}	12
GPS rewired	float 32	9.54×10^{-7}	2.40×10^{-5}	12
GPS full	float 64	1.33×10^{-15}	1.14×10^{-13}	12
GPS full	float 32	9.54×10^{-7}	2.40×10^{-5}	12
GPS local	float 64	1.11×10^{-15}	1.14×10^{-13}	12
GPS local	float 32	7.75×10^{-7}	2.40×10^{-5}	12
GPS no distances	float 64	0	1.14×10^{-13}	12
GPS no distances	float 32	8.34×10^{-7}	2.40×10^{-5}	12
GPS self	float 64	0	1.14×10^{-13}	12
GPS self	float 32	7.15×10^{-7}	2.40×10^{-5}	12
GPS shuffled	float 64	1.55×10^{-15}	1.14×10^{-13}	12
GPS shuffled	float 32	9.54×10^{-7}	2.40×10^{-5}	12
GCN	float 64	0	1.14×10^{-13}	12
GCN	float 32	8.94×10^{-8}	2.40×10^{-5}	12
Local MLP	float 64	0	1.14×10^{-13}	12
Local MLP	float 32	3.58×10^{-7}	2.40×10^{-5}	12
Mean	float 64	3.49×10^{-9}	1.14×10^{-13}	12
Mean	float 32	1.19×10^{-7}	2.40×10^{-5}	12
GraphSAGE	float 64	0	1.14×10^{-13}	12
GraphSAGE	float 32	1.19×10^{-7}	2.40×10^{-5}	12

Table S15: Resource accounting for every new configuration. All three seeds are included. Training wall times were measured under the documented concurrent workload and are not a controlled throughput comparison. Standardized inference uses the same 7,760-node TENX95 section, four CPU threads, two warmups and median of five forwards, including model-specific feature aggregation/Tensor conversion but excluding frozen CNN and graph construction. Peak process RSS includes interpreter and data, not only model activations. No GPU or paid external service was used. Full per-seed values, graph construction times and environments are exported.

Model	Parameters	Best epoch range	Fit s	Forward ms	RSS MiB
GPS full	121018	5–60	5.67	78.4	546
GPS local	121018	5–56	7.04	102.3	568
GPS no distances	121018	5–42	5.45	69.6	568
GPS self	121018	47–53	15.76	34.6	568
GPS shuffled	121018	6–76	22.15	76.5	568
GPS rewired	121018	5–60	3.78	81.5	546
Distance fixed	87154	7–53	12.47	56.5	533
EGNN latent	87282	7–57	13.95	54.2	538
Distance disabled	87154	7–74	12.10	54.1	537
EGNN self	87282	8–35	1.91	11.7	534
EGNN shuffled	87282	7–40	14.76	54.6	545
EGNN rewired	87282	7–57	13.95	62.2	544
Local MLP	72114	8–77	5.30	9.9	520
Mean	72114	8–9	1.24	36.2	550
GCN	72114	8–9	1.26	75.8	581
GraphSAGE	72075	39–60	3.77	58.4	550

Table S16: Every new fit, without seed selection. Histories, optimizer states, input hashes and complete checkpoint SHA 256 values are available in the fit JSON files and inventory CSV; weights are distributed separately from Git.

Model	Seed	Best epoch	Ran	Seconds	RSS MiB
Distance disabled	0	74	80	35.82	632
Distance disabled	1	30	40	12.10	643
Distance disabled	2	7	17	5.66	650
Distance fixed	0	53	63	28.79	613
Distance fixed	1	30	40	12.47	638
Distance fixed	2	7	17	4.35	632
EGNN rewired	0	23	33	13.95	642
EGNN rewired	1	57	67	20.21	651
EGNN rewired	2	7	17	5.02	643
EGNN latent	0	23	33	13.95	640
EGNN latent	1	57	67	19.52	639
EGNN latent	2	7	17	5.50	648
EGNN self	0	35	45	4.64	506
EGNN self	1	8	18	1.88	504
EGNN self	2	8	18	1.91	502
EGNN shuffled	0	24	34	18.71	634

Model	Seed	Best epoch	Ran	Seconds	RSS MiB
EGNN shuffled	1	40	50	14.76	635
EGNN shuffled	2	7	17	4.77	643
GPS rewired	0	60	70	35.00	631
GPS rewired	1	5	15	3.70	633
GPS rewired	2	6	16	3.78	624
GPS full	0	60	70	15.56	626
GPS full	1	5	15	5.67	629
GPS full	2	6	16	3.82	629
GPS local	0	56	66	30.11	692
GPS local	1	5	15	5.24	679
GPS local	2	5	15	7.04	674
GPS no distances	0	42	52	31.17	600
GPS no distances	1	5	15	3.67	599
GPS no distances	2	6	16	5.45	596
GPS self	0	51	61	20.15	543
GPS self	1	47	57	15.76	546
GPS self	2	53	63	13.54	546
GPS shuffled	0	76	80	47.42	631
GPS shuffled	1	46	56	22.15	630
GPS shuffled	2	6	16	6.00	634
GCN	0	8	18	1.26	524
GCN	1	8	18	0.98	522
GCN	2	9	19	1.36	522
Local MLP	0	8	18	2.20	497
Local MLP	1	77	80	5.30	497
Local MLP	2	74	80	6.21	497
Mean	0	8	18	1.27	521
Mean	1	8	18	1.01	522
Mean	2	9	19	1.24	522
GraphSAGE	0	48	58	6.51	570
GraphSAGE	1	60	70	3.77	570
GraphSAGE	2	39	49	2.81	570

Diagnostic interpretation. Undefined correlations remain undefined, with valid-gene counts in the exports. Raw Spearman is secondary because near-tied ranks are floating-point sensitive. Distance curves retain occupied-bin support; different bins need not contain the same locations or patients. Conditional geometric checks and remote-feature sensitivity do not establish biological association. Degree/length controls with zero accepted swaps do not

provide an independent rewired comparison. Optional anchor-conditioned learning and full-image equivariance tests were not performed.




Figure S24: Conditional attention reach. Dots are individual patient summaries after equal seed/head/layer means; large dots are cohort medians. Colors identify250/600/1200micrometer thresholds. This is a software/connectivity diagnostic, not evidence of molecular correlation. All GPS variants, seeds and patients are exported.




Figure S25: All48 source-only training histories. Every one of16 configurations and all three seeds are retained. Solid curves show training MSE, dashed curves validation MSE. Blue, orange and green identify seeds0,1 and2, respectively. Only TENX95 validation MSE selected epochs; evaluation outcomes did not select architecture or seeds.




Figure S26: Predetermined Bassiouni distributed map. BASS P01 at 20percent seed0, the four prespecified genes, original IDW k16/p2. Truth and prediction share each gene scale; anchor colorbar extensions mark clipped anchor values. Gray sites are unscored context. Constant fields are retained and explicitly marked; no tissue annotations are inferred.




Figure S27: Predetermined Bassiouni regional map. BASS P01 central V2 at 20percent seed0, uniform k16 and the same four predetermined genes. Historical V2 has different scored support and estimator from V1; this map comparison does not isolate geometry. Identical-H effects are evaluated separately. Truth/prediction scales are shared within each gene, errors use a separate nonnegative scale.




Figure S28: Distance-resolved Bassiouni reconstruction. All16 new models and both molecular baselines,20percent region-exclusion B2. Curves are medians of per-patient conditional-bin mean errors after equal mask and training-seed averages. Occupied bins only; empty bins are omitted. Source tables retain number of patients and min/median/max mean locations per mask in each bin. Different bins need not contain the same patients or locations and do not identify a biological correlation length.

Table S17: All 48 full-section development-test fits, with every training seed retained. TENX99 at 20% coverage, averaging the same three original masks within each seed. Left metrics are image-only; right metrics use exploratory residual anchors. This is one previously inspected patient, not 48 biological replicates. No confidence interval or seed selection is inferred. Full precision, fixed-fusion, all budgets, external patient metrics and older frozen heads accompany the CSV exports.

Model	Seed	Image r	MSE	MAE	Residual r	MSE	MAE
GPS full	0	0.714	0.9751	0.7814	0.834	0.2349	0.3694
GPS full	1	0.529	1.1646	0.8594	0.801	0.2693	0.3967
GPS full	2	0.569	1.2240	0.8969	0.813	0.2634	0.3909
GPS local	0	0.708	0.9202	0.7560	0.826	0.2662	0.3921
GPS local	1	0.573	1.1372	0.8470	0.804	0.2694	0.3958
GPS local	2	0.508	1.2035	0.8928	0.811	0.2617	0.3885
GPS no distances	0	0.711	1.0426	0.8070	0.835	0.2242	0.3589
GPS no distances	1	0.556	1.1672	0.8560	0.809	0.2596	0.3878
GPS no distances	2	0.577	1.2357	0.8978	0.819	0.2545	0.3827
GPS self	0	0.664	1.0495	0.8149	0.803	0.3344	0.4438
GPS self	1	0.669	1.0613	0.8177	0.795	0.3441	0.4500
GPS self	2	0.671	1.0259	0.8074	0.808	0.3001	0.4204
GPS shuffled	0	0.717	0.9751	0.7806	0.822	0.2865	0.4111
GPS shuffled	1	0.710	1.2113	0.8818	0.829	0.2456	0.3793
GPS shuffled	2	0.552	1.2417	0.9061	0.812	0.2733	0.4001
GPS rewired	0	0.714	0.9751	0.7814	0.834	0.2349	0.3694
GPS rewired	1	0.529	1.1646	0.8594	0.801	0.2693	0.3967
GPS rewired	2	0.569	1.2240	0.8969	0.813	0.2634	0.3909
Distance fixed	0	0.680	0.9648	0.7772	0.820	0.2945	0.4164
Distance fixed	1	0.637	0.9801	0.7883	0.793	0.3136	0.4327
Distance fixed	2	0.471	1.1048	0.8520	0.793	0.2867	0.4118
EGNN latent	0	0.628	1.0248	0.8072	0.799	0.3033	0.4259
EGNN latent	1	0.711	0.9576	0.7706	0.825	0.2646	0.3915
EGNN latent	2	0.536	1.0697	0.8344	0.792	0.3030	0.4239
Distance disabled	0	0.706	1.0438	0.8028	0.831	0.2609	0.3887
Distance disabled	1	0.640	0.9790	0.7877	0.794	0.3128	0.4320
Distance disabled	2	0.471	1.1047	0.8519	0.793	0.2866	0.4118
EGNN self	0	0.595	0.9884	0.7966	0.778	0.3307	0.4476
EGNN self	1	0.546	1.1053	0.8471	0.791	0.3008	0.4227
EGNN self	2	0.545	1.0548	0.8283	0.793	0.3046	0.4253
EGNN shuffled	0	0.627	1.0708	0.8276	0.794	0.3082	0.4303
EGNN shuffled	1	0.665	1.0025	0.7999	0.785	0.3441	0.4550
EGNN shuffled	2	0.532	1.0732	0.8359	0.791	0.3038	0.4245
EGNN rewired	0	0.628	1.0248	0.8072	0.799	0.3033	0.4259

Model	Seed	Image r	MSE	MAE	Residual r	MSE	MAE
EGNN rewired	1	0.711	0.9576	0.7706	0.825	0.2646	0.3915
EGNN rewired	2	0.536	1.0697	0.8344	0.792	0.3030	0.4239
Local MLP	0	0.604	0.9930	0.8144	0.829	0.2433	0.3775
Local MLP	1	0.669	0.9033	0.7628	0.820	0.2591	0.3910
Local MLP	2	0.667	0.9185	0.7714	0.811	0.2571	0.3897
Mean	0	0.624	1.0157	0.8231	0.827	0.2361	0.3667
Mean	1	0.593	1.0522	0.8471	0.819	0.2387	0.3688
Mean	2	0.627	1.0723	0.8525	0.824	0.2366	0.3674
GCN	0	0.620	1.0175	0.8241	0.825	0.2386	0.3692
GCN	1	0.591	1.0536	0.8478	0.818	0.2406	0.3707
GCN	2	0.623	1.0727	0.8529	0.821	0.2385	0.3694
GraphSAGE	0	0.671	0.8701	0.7488	0.827	0.2333	0.3675
GraphSAGE	1	0.688	0.8818	0.7530	0.829	0.2337	0.3683
GraphSAGE	2	0.667	0.9245	0.7785	0.821	0.2307	0.3653

S6 Detailed development methods

These specifications distinguish the original development comparators from the additional minibatch and full-section comparisons. They retain the model-selection and calibration details needed to reproduce the corresponding figures.

S6.1 Development head searches and conditional patient holdouts

The original development MLPs used one hidden linear layer, GELU, dropout and a linear gene-output layer. Features were standardized on fitting patients before graph aggregation; a separate fitting-only head-target scaler was inverted before scoring. Population scales below 10^{-8} were replaced by 1. AdamW used learning rate 10^{-3}, weight decay 10^{-4}, minibatches of 512 with shuffled rows and retained final partial batches, at most 80 epochs, patience 10 and minimum validation-MSE improvement 10^{-4}. Best qualifying parameters were restored, with dropout disabled at prediction. Three training seeds 0/1/2 were retained; hyperparameter candidates used seed 0. Computation used deterministic CPU float32 tensors and four threads, without a learning-rate schedule or auxiliary loss. These settings describe the original development comparisons below, not the later full-section regime.

The initial local comparison in Supplementary Fig. S3 selected hidden width from $\{128, 256\}$ and dropout from $\{0, 0.1\}$ by the lowest validation MSE. For 50 genes, the selected architecture was $512 \to 128 \to 50$ with dropout 0.1. The ridge baseline used an intercept and a Cholesky solver, with penalty selected from $\{10^{-4}, 10^{-3}, 10^{-2}, 10^{-1}, 1, 10, 100\}$ by the highest validation median gene Pearson correlation, breaking ties by lower MSE and then smaller penalty; the selected 50-gene penalty was 100. Training-target and training-embedding permutations disrupted pairing within the fitting data; validation and evaluation data were unchanged.

The representation comparison in Supplementary Fig. S6 used the same four width/dropout candidates but selected the highest validation median gene Pearson correlation, with lower MSE as the first tie-breaker. Both ResNet18 and Phikon selected width 256 and dropout 0.1 for 50 genes. Thus the initial local predictor in Supplementary Fig. S3 and the ResNet18 predictor used in the representation comparison and final sparse pipeline have different selected widths. The latter predictor is described in the main Methods. The representation comparison also evaluated ridge penalties $\{0.1, 1, 10, 100, 1000\}$; both encoders selected 1000 for 50 genes. Equal width and search budget did not imply equal parameter count because the input dimensions differed.

Conditional leave-one-patient-out sensitivity fits used the remaining three development patients, refitted feature and head-target scalers on those patients, retained the primary-selected hyperparameters and used fixed epoch counts without early stopping on the held-out patient. For the 50-gene heads, these counts were 14 for the initial MLP, 14 for mean pooling and GCN, 7 for GraphSAGE, 12 for the representation-comparison ResNet18 head and 1 for the Phikon head. These were rounded medians of the corresponding primary fits' best epochs. Since the configurations and underlying target definition had already been selected during development, these fits are conditional sensitivity analyses.

S6.2 Original graph controls and pathology representation

The matched controls in Supplementary Fig. S5 retained the GraphSAGE head and training settings. The self-only control used $[X, X]$. Vertex permutation preserved the graph's degree distribution, while double-edge rewiring preserved each node's degree. The additional distance-constrained rewiring preserved the numbers of physical edge lengths in $[0, 175)$, $[175, 250)$ and $[250, \infty)$ μm. Rewiring targeted five accepted swaps per edge, with limits of 100 attempts per edge for degree-only rewiring and 1,000 for distance-constrained rewiring; the latter used candidate endpoints within 275 μm. Five graph randomization seeds were crossed with the three model seeds. The Phikon spatial sensitivity comparison reused the selected graph, hidden widths and dropout: mean pooling used $768 \to 128 \to 50$, and GraphSAGE used $1536 \to 67 \to 50$, with true and degree-plus-length-matched graphs. These inherited widths do not constitute exact parameter matching between encoders.

Phikon used the public `owkin/phikon` ViT-B/16 model, pinned to revision `057cc0295895`. Its 12-layer transformer produced the 768-dimensional final-layer-normalized classification-token representation, without a pooler. The official `ViTImageProcessor` processed the same RGB patches as the ResNet18 analysis: 224×224 pixels, rescaling by $1/255$ and ImageNet channel means and standard deviations. There was no recropping, stain normalization or augmentation. Extraction used the model in evaluation mode with frozen weights, float32 precision, batches of 16 and eight CPU threads (Transformers 4.57.3, PyTorch scaled dot-product attention). The head and validation selection were as specified above; the encoder was not fine-tuned.

S6.3 Development ridge residual correction

The learned-propagation model in Supplementary Fig. S7 is a gene-specific ridge residual correction using fixed graph summaries. Its image prediction H is the equal average of the three frozen local MLP predictions from the representation comparison. Let o be the observed-location indicator and Z the expression matrix containing permitted observed vectors and zeros elsewhere. A self-inclusive row-normalized operator P was constructed from the symmetrized six-nearest-neighbor graph capped at 250 μm. The observed masses and molecular messages were

$$m^{(1)} = Po, \quad m^{(2)} = P^2 o, \qquad V_{ig}^{(1)} = \frac{(PZ)_{ig}}{\max(m_i^{(1)}, 10^{-6})}, \quad V_{ig}^{(2)} = \frac{(P^2 Z)_{ig}}{\max(m_i^{(2)}, 10^{-6})}.$$

Image features were standardized on fitting patients and reduced to eight whitened principal components with randomized singular-value decomposition and random seed 0. Denote these components by X_{PCA}. The 29 shared covariates at each location were X_{PCA}, PX_{PCA}, $P^2 X_{\mathrm{PCA}}$, o, $m^{(1)}$, $m^{(2)}$, $\log(1 + d_1/100)$, and the fraction of the nearest 16 observed anchors lying within 300 μm, where d_1 is nearest-anchor distance in μm. Five further covariates for gene g were the image prediction H_g, the fixed fusion prediction F_g, Z_g, $V_g^{(1)}$ and $V_g^{(2)}$. The resulting design has 34 covariates and an intercept for each gene.

Predictions were $\widehat{y}_g = F_g + \Phi_g \theta_g$, where the last column of Φ_g is one. Coefficients were fitted by

$$\underset{\theta_g}{\operatorname{argmin}} \ \|(y_g - F_g) - \Phi_g \theta_g\|_2^2 + \alpha \, \|\theta_{g,1:34}\|_2^2 .$$

The intercept was unpenalized. Fitting sampled up to 256 hidden locations without replacement per training mask using that mask's seed; their expression entered only the supervised response. Masks covered 5, 10, 20, 40, 60 and 80% of locations under random, block and capture-window sampling, with three replicates per condition. The penalty grid was $\{1, 100, 10000\}$, selected by mean validation MSE across masks with equal mask weights; both encoders selected $\alpha = 10000$ for 50 genes. No additional expression scaler or auxiliary loss was introduced.

The underlying development fusion was $F = \lambda H + (1 - \lambda)M$. Nearest-anchor mean interpolation selected k from $\{1, 4, 8, 16\}$; inverse-distance interpolation used up to 16 anchors and selected power from $\{1, 2, 3\}$. The stronger family and $\lambda \in \{0, 0.25, 0.5, 0.75, 1\}$ were chosen by mean validation MSE separately by panel and sampling geometry. For 50 genes, random sampling selected inverse-distance-squared interpolation and $\lambda = 0.25$; block sampling selected the 16-anchor mean and $\lambda = 0.5$; capture-window sampling selected inverse-distance-cubed interpolation and $\lambda = 0.5$. These development settings differ from the final external fusion settings. Supplementary Fig. S7 evaluates three permutation seeds on one random 20% observation mask, with the fitted models unchanged. The plotted B1, B3 and B4 labels denote the frozen image predictor, fixed fusion and ridge residual correction, respectively.

S6.4 Patient-disjoint HER2ST calibration and fusion selection

Supplementary Figs. S1 and S10 use the same eight HER2ST patients and 40 shared genes. Four outer folds held out the pairs A1/B1, C1/D1, E1/F1 and G1/H1, respectively. Within each remaining six-patient set, target-mapping candidates were evaluated by leave-one-patient-out records: the mapping for an inner patient's predictions was estimated from the other five patients. The fixed source moments came from NCBI783 and NCBI785. The candidates were the unchanged target, per-gene mean/standard-deviation mapping and per-gene median/interquartile-range mapping. Each patient had equal total weight in moment or quantile estimates. An interquartile range below 10^{-8} fell back to the corresponding standard deviation, and a standard deviation below 10^{-8} was replaced by one. Source and destination moments were estimated separately. Image predictions were expressed in source-standardized coordinates, anchors in destination-standardized coordinates, and final errors were computed after mapping predictions back to common target units.

The target-mapping comparison retained fixed image weights 0.25 for V1 and 0.5 for V2. Selection used inner random-mask seeds 0–2 and four equally weighted conditions: V1 coverage 10, 20 and 40%, and V2 coverage 20%. Seeds and geometries were first averaged within patient and condition, then conditions within patient, then the six patients. Candidates within 0.005 of the highest mean patient median gene Pearson correlation were eligible; among these, the candidate with lowest common-unit MSE was selected. The mean/standard-deviation transform was selected in every fold; the corresponding outer mapping used moments estimated from all six outer-training patients.

The selected-fusion analysis then used the corresponding inner records to fit and select among four families: molecular interpolation alone; one global image weight from $\{0, 0.1, \ldots, 1\}$; 35 nondecreasing three-bin weight rules with weights from $\{0, 0.1, 0.25, 0.5, 1\}$ and nearest-anchor boundaries 300 and 600 μm; and a five-parameter sigmoid gate. All weights were shared

across genes, regimes and budgets within an outer fold. The sigmoid gate was

$$w_i = \mathrm{sigmoid}(\beta_0 + \widetilde{q}_i^{\top}\beta), \qquad \widehat{y}_i = M_i + w_i(H_i - M_i),$$

where the four features before fitting-data standardization were $\log(1+d_1/200)$, $\log(1+\overline{d}_{16}/200)$, $\log(1+n_{600})$ and $\log(1+v)$: nearest-anchor distance, mean distance to up to 16 nearest anchors, the number of observed anchors within 600 μm, and mean across-gene variance among the neighboring observed vectors. The gate minimized common-unit squared error with equal patient/condition/run/location contributions, divided by each run's interpolation MSE floored at 10^{-8}, plus $0.01\|\beta\|_2^2 + 10^{-6}\beta_0^2$. The same weights defined feature means and standard deviations. L-BFGS-B used initial coefficients $(-2, 0, 0, 0, 0)$, bounds $[-12, 12]$, maximum 300 iterations, function tolerance 10^{-12} and gradient tolerance 10^{-7}. No outer-patient outcomes entered fitting or candidate selection.

The same correlation-then-MSE rule selected the fusion family. The first three patient-pair folds selected global image weight 0.2; the G1/H1 fold selected the sigmoid gate. Outer evaluation used mask seeds 0–9. Supplementary Fig. S10 reports these fold-selected rules, whereas Supplementary Fig. S1 shows the fixed-fusion target-mapping comparison. Both remain calibration-development analyses of previously examined HER2ST patients.

The full-precision primary image feature scaler, target normalization and source-to-Wu calibration constants are supplied in `methods_data/`. The primary head settings and recorded software versions are provided there in machine-readable form. These constants specify the published transformations without refitting them.

S6.5 Identity controls and numerical audit scope

Local, mean-self and GCN-self inputs in the 30-head comparison were bitwise identical. Seeds 1/2 predictions also agreed bitwise; seed 0 weights differed by at most 2.98×10^{-8} and predictions by 9.54×10^{-7}, within the stated float32 guard. These are computational identity checks, not independent biological support. At 20% coverage, matched-seed GCN spatial-minus-self correlation changes ranged from -0.0425 to -0.0149; GraphSAGE changes ranged from $+0.0002$ to $+0.0318$. Shuffled GraphSAGE seed-mean image prediction exceeded the spatial version in TENX99. These development findings do not establish a general topology advantage.

The original Bassiouni evaluation produced 2,904 clean and 18 control bundles (8,766 method predictions); all 17,532 hidden-label substitutions were invariant. The recorded audit reproduced 8,766 predictions and 35,064 per-gene metric arrays and reconstructed 968 label-free mask trajectories. Additional Wu molecular comparisons comprised 2,700 runs and 5,400 exact substitutions; all 360 original random references and 54,000 primary gene values passed 10^{-10} tolerances. Additional Bassiouni comparisons comprised 19,800 runs and 39,600 exact substitutions, with 2,640 original random references and 396,000 primary per-gene values passing the same tolerances. The neural matrix retained 2,028 model–patient bundles and 289,224 exact hidden-label checks; all 384 conditional geometric checks passed. Raw Spearman is retained but excluded from exact-replay claims because of near ties. Hashes, protocols, complete results and environment records provide the detailed provenance.