
MYCRO-TEC ®

SP

AVAILABLE IN



UNLOCK · RELEASE · FEED

ALL PROGRAMS

Most soils already hold the nutrition. SP is the biology that gets it back.

01 UNLOCK

Organic acids break the bonds holding P to Ca · Al · Fe



02 RELEASE

Phosphorus and tied minerals return to solution beside feeder roots



03 FEED

The crop takes up nutrition already in the soil



HORTICULTURE

IN-FURROW	FERTIGATION
5–10 L/ha	5–10 L/ha

PLANTING TO ACTIVE GROWTH

Vegetable and tree crops.

VITICULTURE

IN-FURROW	FERTIGATION
5–10 L/ha	5–10 L/ha

BUDBURST TO ACTIVE GROWTH

Established and developing vines.

BROADACRE & PASTURE

IN-FURROW	COATING
1–2 L/ha	2 L/t

AT SOWING

Cereals, canola and pasture.

INSIDE THIS BROCHURE

PAGE 2 Where SP fits

PAGE 3 Where SP sits


PAGE 4 Composition & function

PAGE 5 Mode of action

PAGE 6 Results in practice

HYBRID AG
PRECISION NUTRITION

AVAILABLE IN



Where SP fits.

The living consortium that turns tied reserves back into crop nutrition.

OPTIMISING INPUTS · WHY SP FITS

The nutrient bank already in your soil.

Only 20–30% of applied phosphorus is taken up. The rest fixes — to calcium at high pH, to aluminium and iron at low pH. SP's four screened organisms break those bonds — returning phosphorus, calcium, iron and tied trace elements to the crop.

ROLE	APPLICATION	WINDOW
mineral-unlock layer	*inject · fertigation · coating*	*planting to active growth*



YOU PAY FOR ALL OF IT

The phosphorus goes on in full. Fixation decides how much stays available to the crop.

WHAT SP CHANGES

SP works on the access side — biology that supports release of reserves already in the soil.

THE BANK IS ALREADY THERE

Most soils already hold years of nutrition. The constraint is access, not supply.

WHAT SP CHANGES

Placed where roots meet fixed reserves, it supports solubilisation from establishment.

NUTRIENT FOLLOWS PHOSPHORUS

When phosphate returns to solution, nitrogen and micronutrient uptake follow it.

WHAT SP CHANGES

A mineral-access layer alongside the program — phosphorus first, the wider bank follows.

WHAT HAS TO BE TRUE

SP works alongside a nutrition program, in soils where testing shows nutrient present but constrained — a nutritional limit, not a structural one.

MAXIMISING OUTCOMES · WHAT GROWERS SEE

Most soils already hold the nutrition. The constraint is access, not supply.

UNLOCKS PHOSPHORUS

At high pH, phosphorus commonly fixes with calcium; at low pH, with aluminium and iron. Organic acids help release it.

PROGRAM SIGNAL · P AVAILABILITY



BREAKS THE P BONDS

Gluconic, oxalic and citric acids chelate the calcium, aluminium and iron holding phosphorus, releasing it near active roots.

PROGRAM SIGNAL · MINERAL MOBILITY



STRONGER ROOT SYSTEMS

More available nutrition supports feeder-root density and crop establishment.

PROGRAM SIGNAL · ROOT RESPONSE



NITROGEN EFFICIENCY

As phosphate returns to solution, nitrogen and micronutrient uptake follow it.

PROGRAM SIGNAL · UPTAKE EFFICIENCY




PRECISION NUTRITION
Physical · Mineral · Biological

OPTIMISING INPUTS
Addressed · Optimised · Enabled

MAXIMISING OUTCOMES
Measured · Proven · Delivered

HYBRID AG
PRECISION NUTRITION

AVAILABLE IN



Where SP sits in your program.

The same four lanes your program already runs.

THE SEASON · FIVE WINDOWS, ONE PROGRAM

Simplified, illustrative program to show you when it goes on — actual programs vary with crop, variety, soil, season, test data, goals.



ESTABLISHMENT

GROUND SPREAD
Prescription dry blend

SOIL PRIMER
—

+ SP · planting pass

FOLIAR
—

FERTIGATION
—



EARLY GROWTH

GROUND SPREAD
—

SOIL PRIMER
Biological primer

FOLIAR
Trace elements

FERTIGATION
Base nutrition

+ SP · fertigation



FLOWERING & SET

GROUND SPREAD
—

SOIL PRIMER
—

FOLIAR
Calcium · boron

FERTIGATION
Base nutrition



FRUIT FILL

GROUND SPREAD
—

SOIL PRIMER
—

FOLIAR
Calcium · potassium

FERTIGATION
Base nutrition



FINISH


GROUND SPREAD
—

SOIL PRIMER
—

FOLIAR
Potassium

FERTIGATION
—


SP rides the planting pass and early fertigation — rates by segment on the front cover.



YOUR PROGRAM
NO EXTRA PASS
Goes in with planting and the watering you already run — programmed, not a rescue.



NEXT · PAGE 4
COMPOSITION & FUNCTION
What SP actually is — the organisms inside it and the job each one does.



THEN · PAGE 5
MODE OF ACTION
The stage-by-stage breakdown for agronomists and technical growers.

ALL OUR PROGRAMS CONTAIN

NUTRIENT LOADING
Every product's N, P, K, S, Ca, Mg and trace contribution, totalled per hectare.


COST PER HECTARE
Each pass costed per hectare — judged on what it costs to grow the crop.

AGAINST THE TARGET
Supply weighed against target yield and removal, so the program balances.

Build the biology in. Carry the response through the season.

HYBRID AG
PRECISION NUTRITION


AVAILABLE IN



Composition & function.

Four screened organisms, three fungi and one dominant bacterium — built to mobilise tied nutrition.

INSIDE SP · WHAT EACH GROUP DOES



SOLUBILISING BACTERIA

WHAT IT IS

Enterobacter cloacae — the high-load bacterial engine.

WHAT IT DOES

Produces organic acids and drives rapid mineral mobilisation through the active root zone.

CONTAINS

E. cloacae >2×10⁹ CFU/mL

RAPID NUTRIENT RELEASE



BENEFICIAL FUNGI

WHAT IT IS

Trichoderma, Aspergillus and Penicillium — three complementary fungi.

WHAT IT DOES

Colonise soil and mineral surfaces, extend the working zone and support ongoing nutrient cycling.

CONTAINS

T. harzianum · A. niger · P. radicum

MULTI-GENUS CONSORTIUM



ORGANIC-ACID ENGINE

WHAT IT IS

Gluconic, oxalic and citric acids produced by the consortium.

WHAT IT DOES

Chelate metals and break the bonds tying phosphorus and other nutrients to calcium, aluminium and iron.

CONTAINS

Gluconic · oxalic · citric acid

BOND BREAKING + MOBILITY

TYPICAL ANALYSIS · VIABLE ORGANISMS

Enterobacter cloacae	Trichoderma harzianum	Aspergillus niger	Penicillium radicum
E. cloacae	*T. harzianum*	*A. niger*	*P. radicum*
$>2\times10^{9}$ CFU/mL	$>2\times10^{8}$ CFU/mL	$>2\times10^{7}$ CFU/mL	$>1\times10^{7}$ CFU/mL

Concentrations at production. Formula per current SP label; use the current label and SDS for final directions.

MIXING · STORAGE · SAFETY

MIXING & USAGE

Shake well. Dilute 1:100 in cool, clean, non-chlorinated water. Apply promptly; never hold diluted product overnight. Avoid crop-protection mixes unless confirmed.

STORAGE & HANDLING

Store at 5–30 °C, out of direct sunlight. Do not freeze. Keep tightly closed. 12-month shelf life from manufacture when stored correctly.

SAFETY

Keep out of reach of children. May irritate skin or eyes. Follow the current label and SDS.

HYBRID AG
PRECISION NUTRITION

Mode of action.

SP works in five linked steps — from placement to crop uptake.

AVAILABLE IN



THE FIVE-STAGE CASCADE

Designed for the agronomist who needs to see how tied reserves become available.

1 APPLICATION & PLACEMENT

Applied by liquid injection, fertigation, seed coating or fertiliser coating — positioned close to the active root zone.

KEY · TARGETED PLACEMENT



2 RHIZOSPHERE COLONISATION

The four screened organisms establish along roots, mineral surfaces and the surrounding rhizosphere.

KEY · LIVING BIOLOGY ESTABLISHED



3 ORGANIC-ACID SECRETION

While active, the living consortium produces gluconic, oxalic and citric acids around tied mineral reserves.

KEY · GLUCONIC + OXALIC + CITRIC



4 BOND BREAKING & RELEASE

Acids chelate metals and disrupt the bonds tying phosphorus and other nutrients to calcium, aluminium and iron.

KEY · MINERALS MOBILISED



5 UPTAKE & GROWTH

Released nutrition moves toward feeder roots, supporting establishment, root density and crop response.

KEY · PLANT-AVAILABLE NUTRITION



PRECISION NUTRITION
Physical · Mineral · Biological

OPTIMISING INPUTS
Addressed · Optimised · Enabled


MAXIMISING OUTCOMES
Measured · Proven · Delivered

HYBRID AG
PRECISION NUTRITION

Results in practice.

Gerogery NSW canola · Nov 2025 · published GSB field result · granular formulation · site-specific.

AVAILABLE IN



BETTER TOGETHER · THREE COMPLEMENTARY JOBS



DIGESTER · DIGESTS RESIDUE

ROLE


Cellulose- and lignin-degrading microbes and their enzymes break down last season's stubble.

PAYOFF

Returns the nutrition tied up in crop residue to the soil, and improves structure and water-holding for the next crop.

DIGEST
Breaks down residue

MYCRO-TEC DIGESTER



SP · UNLOCKS NUTRIENT

ROLE


Four screened microbes secrete gluconic, oxalic and citric acids that break the bonds locking up soil nutrient.

PAYOFF

Helps release phosphorus and other nutrients fixed with calcium, aluminium and iron.

UNLOCK
Releases fixed P

MYCRO-TEC SP



LOCKOUT · COLONISES THE ZONE

ROLE

Purpose-selected living biology colonises the rhizosphere and competes for space and food.

PAYOFF

Supports a more balanced root-zone microbiome and a productive living zone.

COLONISE
Colonises the living zone

MYCRO-TEC LOCKOUT

RESULTS IN PRACTICE · BROADACRE TRIAL

Phosphorus locked by aluminium.

The granular formulation was applied in a treated strip through a 52.7 ha canola paddock; pre-harvest tests and the yield map showed a clear response.

SOURCE · GSB CASE STUDY 2195 · GRANULAR FORMULATION



THE PROOF IS IN THE PADDOCK

The published case study reports a measured yield and soil-nutrient response across the treated strip.

SCAN QR CODE TO VIEW THE FULL CANOLA TRIAL FROM GREAT SOUTHERN BIOLOGY

52.7 ha	**3.40 → 3.98**	**+17%**	**+0.58 t/ha**
canola paddock	*t/ha average yield*	*yield improvement*	*above paddock average*

+35.7%	**+17.9%**	**26 → 38**
Colwell P · 42 → 57 mg/kg	*Colwell K · 184 → 217 mg/kg*	*nitrate mg/kg, treated strip*

PRECISION NUTRITION
Physical · Mineral · Biological

OPTIMISING INPUTS
Addressed · Optimised · Enabled

MAXIMISING OUTCOMES
Measured · Proven · Delivered