
R=0.405 | ALL pre-PID | log




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old unaccepted); no fit / no PID changes

Nuclear window mode 64.25, N 8005120; window mean/RMS are NOT Gaussian fit

R=0.405 | TOF ^{3}He window | log




1.78285 < $(m/q)^2$ < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old unaccepted); no fit / no PID changes

Nuclear window mode 76.75, N 376; window mean/RMS are NOT Gaussian fit

R=0.405 | TOF D / ^{4}He-compatible | log




3.26877 < $(m/q)^2$ < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old unaccepted); no fit / no PID changes

Nuclear window mode 71.75, N 131; window mean/RMS are NOT Gaussian fit

R=0.405 | ALL pre-PID | linear




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old unaccepted); no fit / no PID changes

Nuclear window mode 64.25, N 8005120; window mean/RMS are NOT Gaussian fit

R=0.405 | TOF ^{3}He window | linear




1.78285 < $(m/q)^2$ < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old unaccepted); no fit / no PID changes

Nuclear window mode 76.75, N 376; window mean/RMS are NOT Gaussian fit

R=0.405 | TOF D / ^{4}He-compatible | linear




3.26877 < $(m/q)^2$ < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old unaccepted); no fit / no PID changes

Nuclear window mode 71.75, N 131; window mean/RMS are NOT Gaussian fit

R=0.415 | ALL pre-PID | log




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 63.25, N 8787753; window mean/RMS are NOT Gaussian fit

R=0.415 | TOF ^{3}He window | log




1.78285 < (m/q)^2 < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 72.75, N 597; window mean/RMS are NOT Gaussian fit

R=0.415 | TOF D / ^{4}He-compatible | log




3.26877 < (m/q)^2 < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 69.75, N 177; window mean/RMS are NOT Gaussian fit

R=0.415 | ALL pre-PID | linear




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 63.25, N 8787753; window mean/RMS are NOT Gaussian fit

R=0.415 | TOF ^{3}He window | linear




1.78285 < (m/q)^2 < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 72.75, N 597; window mean/RMS are NOT Gaussian fit

R=0.415 | TOF D / ^{4}He-compatible | linear




3.26877 < (m/q)^2 < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 69.75, N 177; window mean/RMS are NOT Gaussian fit

R=0.425 | ALL pre-PID | log




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 62.25, N 9348458; window mean/RMS are NOT Gaussian fit

R=0.425 | TOF ^{3}He window | log




1.78285 < $(m/q)^2$ < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 69.25, N 769; window mean/RMS are NOT Gaussian fit

R=0.425 | TOF D / ^{4}He-compatible | log




3.26877 < $(m/q)^2$ < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 62.75, N 169; window mean/RMS are NOT Gaussian fit

R=0.425 | ALL pre-PID | linear




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 62.25, N 9348458; window mean/RMS are NOT Gaussian fit

R=0.425 | TOF ^{3}He window | linear




1.78285 < $(m/q)^2$ < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 69.25, N 769; window mean/RMS are NOT Gaussian fit

R=0.425 | TOF D / ^{4}He-compatible | linear




3.26877 < $(m/q)^2$ < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 62.75, N 169; window mean/RMS are NOT Gaussian fit

R=0.435 | ALL pre-PID | log




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 60.75, N 9467393; window mean/RMS are NOT Gaussian fit

R=0.435 | TOF ^{3}He window | log




$1.78285 < (m/q)^2 < 2.06485$; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 71.25, N 1066; window mean/RMS are NOT Gaussian fit

R=0.435 | TOF D / ^{4}He-compatible | log




$3.26877 < (m/q)^2 < 3.69315$; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 62.25, N 242; window mean/RMS are NOT Gaussian fit

R=0.435 | ALL pre-PID | linear




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 60.75, N 9467393; window mean/RMS are NOT Gaussian fit

R=0.435 | TOF ^{3}He window | linear




$1.78285 < (m/q)^2 < 2.06485$; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 71.25, N 1066; window mean/RMS are NOT Gaussian fit

R=0.435 | TOF D / ^{4}He-compatible | linear




$3.26877 < (m/q)^2 < 3.69315$; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 62.25, N 242; window mean/RMS are NOT Gaussian fit

R=0.445 | ALL pre-PID | log




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 59.75, N 9747313; window mean/RMS are NOT Gaussian fit

R=0.445 | TOF ^{3}He window | log




1.78285 < $(m/q)^2$ < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 67.75, N 1627; window mean/RMS are NOT Gaussian fit

R=0.445 | TOF D / ^{4}He-compatible | log




3.26877 < $(m/q)^2$ < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 61.25, N 229; window mean/RMS are NOT Gaussian fit

R=0.445 | ALL pre-PID | linear




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 59.75, N 9747313; window mean/RMS are NOT Gaussian fit

R=0.445 | TOF ^{3}He window | linear




1.78285 < $(m/q)^2$ < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 67.75, N 1627; window mean/RMS are NOT Gaussian fit

R=0.445 | TOF D / ^{4}He-compatible | linear




3.26877 < $(m/q)^2$ < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 61.25, N 229; window mean/RMS are NOT Gaussian fit

R=0.455 | ALL pre-PID | log




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 58.75, N 9830174; window mean/RMS are NOT Gaussian fit

R=0.455 | TOF ^{3}He window | log




1.78285 < $(m/q)^2$ < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 67.25, N 2348; window mean/RMS are NOT Gaussian fit

R=0.455 | TOF D / ^{4}He-compatible | log




3.26877 < $(m/q)^2$ < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 60.25, N 235; window mean/RMS are NOT Gaussian fit

R=0.455 | ALL pre-PID | linear




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 58.75, N 9830174; window mean/RMS are NOT Gaussian fit

R=0.455 | TOF ^{3}He window | linear




1.78285 < $(m/q)^2$ < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 67.25, N 2348; window mean/RMS are NOT Gaussian fit

R=0.455 | TOF D / ^{4}He-compatible | linear




3.26877 < $(m/q)^2$ < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 60.25, N 235; window mean/RMS are NOT Gaussian fit

R=0.465 | ALL pre-PID | log




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 57.25, N 9283517; window mean/RMS are NOT Gaussian fit

R=0.465 | TOF ^{3}He window | log




1.78285 < $(m/q)^2$ < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 64.75, N 3291; window mean/RMS are NOT Gaussian fit

R=0.465 | TOF D / ^{4}He-compatible | log




3.26877 < $(m/q)^2$ < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 57.25, N 179; window mean/RMS are NOT Gaussian fit

R=0.465 | ALL pre-PID | linear




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 57.25, N 9283517; window mean/RMS are NOT Gaussian fit

R=0.465 | TOF ^{3}He window | linear




1.78285 < $(m/q)^2$ < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 64.75, N 3291; window mean/RMS are NOT Gaussian fit

R=0.465 | TOF D / ^{4}He-compatible | linear




3.26877 < $(m/q)^2$ < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 57.25, N 179; window mean/RMS are NOT Gaussian fit

R=0.475 | ALL pre-PID | log




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 56.25, N 8886824; window mean/RMS are NOT Gaussian fit

R=0.475 | TOF ^{3}He window | log




1.78285 < $(m/q)^2$ < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 62.25, N 4532; window mean/RMS are NOT Gaussian fit

R=0.475 | TOF D / ^{4}He-compatible | log




3.26877 < $(m/q)^2$ < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 57.75, N 236; window mean/RMS are NOT Gaussian fit

R=0.475 | ALL pre-PID | linear




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 56.25, N 8886824; window mean/RMS are NOT Gaussian fit

R=0.475 | TOF ^{3}He window | linear




1.78285 < $(m/q)^2$ < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 62.25, N 4532; window mean/RMS are NOT Gaussian fit

R=0.475 | TOF D / ^{4}He-compatible | linear




3.26877 < $(m/q)^2$ < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 57.75, N 236; window mean/RMS are NOT Gaussian fit

R=0.485 | ALL pre-PID | log




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 55.25, N 8288655; window mean/RMS are NOT Gaussian fit

R=0.485 | TOF ^{3}He window | log




1.78285 < $(m/q)^2$ < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 63.25, N 6771; window mean/RMS are NOT Gaussian fit

R=0.485 | TOF D / ^{4}He-compatible | log




3.26877 < $(m/q)^2$ < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 56.75, N 250; window mean/RMS are NOT Gaussian fit

R=0.485 | ALL pre-PID | linear




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 55.25, N 8288655; window mean/RMS are NOT Gaussian fit

R=0.485 | TOF ^{3}He window | linear




1.78285 < $(m/q)^2$ < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 63.25, N 6771; window mean/RMS are NOT Gaussian fit

R=0.485 | TOF D / ^{4}He-compatible | linear




3.26877 < $(m/q)^2$ < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 56.75, N 250; window mean/RMS are NOT Gaussian fit

R=0.495 | ALL pre-PID | log




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 55.25, N 8398890; window mean/RMS are NOT Gaussian fit

R=0.495 | TOF ^{3}He window | log




1.78285 < $(m/q)^2$ < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 61.25, N 8753; window mean/RMS are NOT Gaussian fit

R=0.495 | TOF D / ^{4}He-compatible | log




3.26877 < $(m/q)^2$ < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 60.25, N 297; window mean/RMS are NOT Gaussian fit

R=0.495 | ALL pre-PID | linear




Unselected ALL; same basic quality as TOF controls

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 55.25, N 8398890; window mean/RMS are NOT Gaussian fit

R=0.495 | TOF ^{3}He window | linear




1.78285 < $(m/q)^2$ < 2.06485; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 61.25, N 8753; window mean/RMS are NOT Gaussian fit

R=0.495 | TOF D / ^{4}He-compatible | linear



3.26877 < $(m/q)^2$ < 3.69315; pre-TPC subset, NOT pure isotope

from040 centers are references (old accepted); no fit / no PID changes

Nuclear window mode 60.25, N 297; window mean/RMS are NOT Gaussian fit