


d



d



d



Nominal primary: d

Gaussian: center = arithmetic mean; width = std deviation.

Log-normal: center = mode; width = local curvature scale.

Widths have different meanings: this is NOT variance-unfolding.

Native-R projected responses; neither width is intrinsic validation.

D/N uses the whole SAME fitted TAG window, not primary purity.

For d and 4He, D/N repeats the SAME TAG_D fit (not independent).


Gaussian/log-normal are non-nested; delta D is not significance.

No accepted isotope identities, production curves, or purity.


Four markers: filled/open circle = Gaussian pol1/exp;

filled/open square = log-normal pol1/exp. See TSV for flags.


t



t



t



Nominal primary: t

Gaussian: center = arithmetic mean; width = std deviation.

Log-normal: center = mode; width = local curvature scale.

Widths have different meanings: this is NOT variance-unfolding.

Native-R projected responses; neither width is intrinsic validation.

D/N uses the whole SAME fitted TAG window, not primary purity.


For d and 4He, D/N repeats the SAME TAG_D fit (not independent).


Gaussian/log-normal are non-nested; delta D is not significance.


No accepted isotope identities, production curves, or purity.

Four markers: filled/open circle = Gaussian pol1/exp;

filled/open square = log-normal pol1/exp. See TSV for flags.







Nominal primary: he3

Gaussian: center = arithmetic mean; width = std deviation.

Log-normal: center = mode; width = local curvature scale.

Widths have different meanings: this is NOT variance-unfolding.

Native-R projected responses; neither width is intrinsic validation.

D/N uses the whole SAME fitted TAG window, not primary purity.


For d and 4He, D/N repeats the SAME TAG_D fit (not independent).


Gaussian/log-normal are non-nested; delta D is not significance.


No accepted isotope identities, production curves, or purity.

Four markers: filled/open circle = Gaussian pol1/exp;

filled/open square = log-normal pol1/exp. See TSV for flags.







Nominal primary: he4

Gaussian: center = arithmetic mean; width = std deviation.

Log-normal: center = mode; width = local curvature scale.

Widths have different meanings: this is NOT variance-unfolding.

Native-R projected responses; neither width is intrinsic validation.

D/N uses the whole SAME fitted TAG window, not primary purity.

For d and 4He, D/N repeats the SAME TAG_D fit (not independent).

Gaussian/log-normal are non-nested; delta D is not significance.

No accepted isotope identities, production curves, or purity.

Four markers: filled/open circle = Gaussian pol1/exp;

filled/open square = log-normal pol1/exp. See TSV for flags.