



CS 228 : Logic in Computer Science

Krishna. S

Welcome

What is this course about? A mini-zoo of logics.

Welcome

What is this course about? A mini-zoo of logics. Here are some typical questions you will learn to answer:

Welcome

What is this course about? A mini-zoo of logics. Here are some typical questions you will learn to answer:

- ▶ Q1: Given a formula φ in a logic L , is φ satisfiable?

Welcome

What is this course about? A mini-zoo of logics. Here are some typical questions you will learn to answer:

- ▶ Q1: Given a formula φ in a logic L , is φ satisfiable?
- ▶ Q2: Given a formula φ in a logic L , is φ valid?

Welcome

What is this course about? A mini-zoo of logics. Here are some typical questions you will learn to answer:

- ▶ Q1: Given a formula φ in a logic L , is φ satisfiable?
- ▶ Q2: Given a formula φ in a logic L , is φ valid?
- ▶ Q3: How easy is to answer Q1 and Q2?

Welcome

What is this course about? A mini-zoo of logics. Here are some typical questions you will learn to answer:

- ▶ Q1: Given a formula φ in a logic L , is φ satisfiable?
- ▶ Q2: Given a formula φ in a logic L , is φ valid?
- ▶ Q3: How easy is to answer Q1 and Q2?
- ▶ Q4: Can you write an algorithm to answer Q1 and Q2?

Welcome

What is this course about? A mini-zoo of logics. Here are some typical questions you will learn to answer:

- ▶ Q1: Given a formula φ in a logic L , is φ satisfiable?
- ▶ Q2: Given a formula φ in a logic L , is φ valid?
- ▶ Q3: How easy is to answer Q1 and Q2?
- ▶ Q4: Can you write an algorithm to answer Q1 and Q2?
- ▶ Q5: Can you “prove” any factually correct statement using the chosen logic L ?

Welcome

What is this course about? A mini-zoo of logics. Here are some typical questions you will learn to answer:

- ▶ Q1: Given a formula φ in a logic L , is φ satisfiable?
- ▶ Q2: Given a formula φ in a logic L , is φ valid?
- ▶ Q3: How easy is to answer Q1 and Q2?
- ▶ Q4: Can you write an algorithm to answer Q1 and Q2?
- ▶ Q5: Can you “prove” any factually correct statement using the chosen logic L ?
- ▶ Q6: How is logic L used in computer science?

Welcome

What is this course about? A mini-zoo of logics. Here are some typical questions you will learn to answer:

- ▶ Q1: Given a formula φ in a logic L , is φ satisfiable?
- ▶ Q2: Given a formula φ in a logic L , is φ valid?
- ▶ Q3: How easy is to answer Q1 and Q2?
- ▶ Q4: Can you write an algorithm to answer Q1 and Q2?
- ▶ Q5: Can you “prove” any factually correct statement using the chosen logic L ?
- ▶ Q6: How is logic L used in computer science?
- ▶ Q7: What are the techniques needed to go about these questions?

Members of the mini-zoo

We will restrict ourselves to the following members:

Members of the mini-zoo

We will restrict ourselves to the following members:

- ▶ Propositional Logic

Members of the mini-zoo

We will restrict ourselves to the following members:

- ▶ Propositional Logic
- ▶ First Order Logic

Members of the mini-zoo

We will restrict ourselves to the following members:

- ▶ Propositional Logic
- ▶ First Order Logic
- ▶ Monadic Second Order Logic

Members of the mini-zoo

We will restrict ourselves to the following members:

- ▶ Propositional Logic
- ▶ First Order Logic
- ▶ Monadic Second Order Logic
- ▶ Linear Temporal Logic

Members of the mini-zoo

We will restrict ourselves to the following members:

- ▶ Propositional Logic
- ▶ First Order Logic
- ▶ Monadic Second Order Logic
- ▶ Linear Temporal Logic
- ▶ Their applications in CS

More if time permits!

References

- ▶ To start with, the text book of Huth and Ryan : Logic for CS.

References

- ▶ To start with, the text book of Huth and Ryan : Logic for CS.
- ▶ As we go ahead, lecture notes/monographs/other text books.

References

- ▶ To start with, the text book of Huth and Ryan : Logic for CS.
- ▶ As we go ahead, lecture notes/monographs/other text books.
- ▶ Classes : Slot 5. Tutorial: To discuss.
- ▶ Confirmed TAs: Anish Yogesh Kulkarni, Ameya Vikrama Singh, Om Swostik Mishra, Agnipratim Das, Nilabha Saha, Ashwin Abraham

Propositional Logic

Syntax

- ▶ Finite set of propositional variables p, q, \dots

Syntax

- ▶ Finite set of propositional variables p, q, \dots
- ▶ Each of these can be true/false

Syntax

- ▶ Finite set of propositional variables p, q, \dots
- ▶ Each of these can be true/false
- ▶ Combine propositions using $\neg, \vee, \wedge, \rightarrow$

Syntax

- ▶ Finite set of propositional variables p, q, \dots
- ▶ Each of these can be true/false
- ▶ Combine propositions using $\neg, \vee, \wedge, \rightarrow$
- ▶ Parantheses as required

Syntax

- ▶ Finite set of propositional variables p, q, \dots
- ▶ Each of these can be true/false
- ▶ Combine propositions using $\neg, \vee, \wedge, \rightarrow$
- ▶ Parantheses as required
- ▶ Example : $[p \wedge (q \vee r)] \rightarrow [\neg r \wedge p]$
- ▶ \neg binds tighter than \vee, \wedge , which bind tighter than \rightarrow . In the absence of parantheses, $p \rightarrow q \rightarrow r$ is read as $p \rightarrow (q \rightarrow r)$

Natural Deduction

- ▶ If it rains, Tia is outside and does not have any raingear with her, she will get wet. $\varphi = (R \wedge TiaOut \wedge \neg RG) \rightarrow TiaWet$

Natural Deduction

- ▶ If it rains, Tia is outside and does not have any raingear with her, she will get wet. $\varphi = (R \wedge TiaOut \wedge \neg RG) \rightarrow TiaWet$
- ▶ It is raining, and Tia is outside, and is not wet.
 $\psi = (R \wedge TiaOut \wedge \neg TiaWet)$

Natural Deduction

- ▶ If it rains, Tia is outside and does not have any raingear with her, she will get wet. $\varphi = (R \wedge TiaOut \wedge \neg RG) \rightarrow TiaWet$
- ▶ It is raining, and Tia is outside, and is not wet.
 $\psi = (R \wedge TiaOut \wedge \neg TiaWet)$
- ▶ So, Tia has her rain gear with her. RG
- ▶ Thus, $\chi = \varphi \wedge \psi \rightarrow RG$. You can deduce RG from $\varphi \wedge \psi$.
- ▶ Is χ valid? Is χ satisfiable?

Two Examples of Natural Deduction

Solve Sudoku

Consider the following kid's version of Sudoku.

	2	4	
1			3
4			2
	1	3	

Rules:

- ▶ Each row must contain all numbers 1-4
- ▶ Each column must contain all numbers 1-4
- ▶ Each 2×2 block must contain all numbers 1-4
- ▶ No cell contains 2 or more numbers

Encoding as Propositional Satisfiability

- ▶ Proposition $P(i, j, n)$ is true when cell (i, j) has number n

Encoding as Propositional Satisfiability

- ▶ Proposition $P(i, j, n)$ is true when cell (i, j) has number n
- ▶ $4 \times 4 \times 4$ propositions

Encoding as Propositional Satisfiability

- ▶ Proposition $P(i, j, n)$ is true when cell (i, j) has number n
- ▶ $4 \times 4 \times 4$ propositions
- ▶ Each row must contain all 4 numbers
 - ▶ Row 1: $[P(1, 1, 1) \vee P(1, 2, 1) \vee P(1, 3, 1) \vee P(1, 4, 1)] \wedge$
 $[P(1, 1, 2) \vee P(1, 2, 2) \vee P(1, 3, 2) \vee P(1, 4, 2)] \wedge$
 $[P(1, 1, 3) \vee P(1, 2, 3) \vee P(1, 3, 3) \vee P(1, 4, 3)] \wedge$
 $[P(1, 1, 4) \vee P(1, 2, 4) \vee P(1, 3, 4) \vee P(1, 4, 4)]$

Encoding as Propositional Satisfiability

- ▶ Proposition $P(i, j, n)$ is true when cell (i, j) has number n
- ▶ $4 \times 4 \times 4$ propositions
- ▶ Each row must contain all 4 numbers
 - ▶ Row 1: $[P(1, 1, 1) \vee P(1, 2, 1) \vee P(1, 3, 1) \vee P(1, 4, 1)] \wedge$
 $[P(1, 1, 2) \vee P(1, 2, 2) \vee P(1, 3, 2) \vee P(1, 4, 2)] \wedge$
 $[P(1, 1, 3) \vee P(1, 2, 3) \vee P(1, 3, 3) \vee P(1, 4, 3)] \wedge$
 $[P(1, 1, 4) \vee P(1, 2, 4) \vee P(1, 3, 4) \vee P(1, 4, 4)]$
 - ▶ Row 2: $[P(2, 1, 1) \vee \dots$
 - ▶ Row 3: $[P(3, 1, 1) \vee \dots$
 - ▶ Row 4: $[P(4, 1, 1) \vee \dots$

Encoding as Propositional Satisfiability

Each column must contain all numbers 1-4

Encoding as Propositional Satisfiability

Each column must contain all numbers 1-4

- ▶ Column 1: $[P(1, 1, 1) \vee P(2, 1, 1) \vee P(3, 1, 1) \vee P(4, 1, 1)] \wedge$
 $[P(1, 1, 2) \vee P(2, 1, 2) \vee P(3, 1, 2) \vee P(4, 1, 2)] \wedge$
 $[P(1, 1, 3) \vee P(2, 1, 3) \vee P(3, 1, 3) \vee P(4, 1, 3)] \wedge$
 $[P(1, 1, 4) \vee P(2, 1, 4) \vee P(3, 1, 4) \vee P(4, 1, 4)]$

Encoding as Propositional Satisfiability

Each column must contain all numbers 1-4

- ▶ Column 1: $[P(1, 1, 1) \vee P(2, 1, 1) \vee P(3, 1, 1) \vee P(4, 1, 1)] \wedge$
 $[P(1, 1, 2) \vee P(2, 1, 2) \vee P(3, 1, 2) \vee P(4, 1, 2)] \wedge$
 $[P(1, 1, 3) \vee P(2, 1, 3) \vee P(3, 1, 3) \vee P(4, 1, 3)] \wedge$
 $[P(1, 1, 4) \vee P(2, 1, 4) \vee P(3, 1, 4) \vee P(4, 1, 4)]$
- ▶ Column 2: $[P(1, 2, 1) \vee \dots$
- ▶ Column 3: $[P(1, 3, 1) \vee \dots$
- ▶ Column 4: $[P(1, 4, 1) \vee \dots$

Encoding as Propositional Satisfiability

Each 2×2 block must contain all numbers 1-4

Encoding as Propositional Satisfiability

Each 2×2 block must contain all numbers 1-4

- ▶ Upper left block contains all numbers 1-4:

$$\begin{aligned} & [P(1, 1, 1) \vee P(1, 2, 1) \vee P(2, 1, 1) \vee P(2, 2, 1)] \wedge \\ & [P(1, 1, 2) \vee P(1, 2, 2) \vee P(2, 1, 2) \vee P(2, 2, 2)] \wedge \\ & [P(1, 1, 3) \vee P(1, 2, 3) \vee P(2, 1, 3) \vee P(2, 2, 3)] \wedge \\ & [P(1, 1, 4) \vee P(1, 2, 4) \vee P(2, 1, 4) \vee P(2, 2, 4)] \end{aligned}$$

Encoding as Propositional Satisfiability

Each 2×2 block must contain all numbers 1-4

- ▶ Upper left block contains all numbers 1-4:

$$\begin{aligned} & [P(1, 1, 1) \vee P(1, 2, 1) \vee P(2, 1, 1) \vee P(2, 2, 1)] \wedge \\ & [P(1, 1, 2) \vee P(1, 2, 2) \vee P(2, 1, 2) \vee P(2, 2, 2)] \wedge \\ & [P(1, 1, 3) \vee P(1, 2, 3) \vee P(2, 1, 3) \vee P(2, 2, 3)] \wedge \\ & [P(1, 1, 4) \vee P(1, 2, 4) \vee P(2, 1, 4) \vee P(2, 2, 4)] \end{aligned}$$

- ▶ Upper right block contains all numbers 1-4:

$$[P(1, 3, 1) \vee P(1, 4, 1) \vee P(2, 3, 1) \vee P(2, 4, 1)] \wedge \dots$$

- ▶ Lower left block contains all numbers 1-4:

$$[P(3, 1, 1) \vee P(3, 2, 1) \vee P(4, 1, 1) \vee P(4, 2, 1)] \wedge \dots$$

- ▶ Lower right block contains all numbers 1-4:

$$[P(3, 3, 1) \vee P(3, 4, 1) \vee P(4, 3, 1) \vee P(4, 4, 1)] \wedge \dots$$

Encoding as Propositional Satisfiability

No cell contains 2 or more numbers

- ▶ For cell(1,1):

$$P(1, 1, 1) \rightarrow [\neg P(1, 1, 2) \wedge \neg P(1, 1, 3) \wedge \neg P(1, 1, 4)] \wedge$$

$$P(1, 1, 2) \rightarrow [\neg P(1, 1, 1) \wedge \neg P(1, 1, 3) \wedge \neg P(1, 1, 4)] \wedge$$

$$P(1, 1, 3) \rightarrow [\neg P(1, 1, 1) \wedge \neg P(1, 1, 2) \wedge \neg P(1, 1, 4)] \wedge$$

$$P(1, 1, 4) \rightarrow [\neg P(1, 1, 1) \wedge \neg P(1, 1, 2) \wedge \neg P(1, 1, 3)] \wedge$$

- ▶ Similar for other cells

Encoding as Propositional Satisfiability

Encoding Initial Configuration:

$$P(1, 2, 2) \wedge P(1, 3, 4) \wedge P(2, 1, 1) \wedge P(2, 4, 3) \wedge \\ P(3, 1, 4) \wedge P(3, 4, 2) \wedge P(4, 2, 1) \wedge P(4, 3, 3)$$

Solving Sudoku

To solve the puzzle, just conjunct all the above formulae and find a satisfiable truth assignment!

Gold Rush

(**Box1**) *The gold is not here*

(**Box2**) *The gold is not here*

(**Box3**) *The gold is in Box 2*

Only one message is true; the other two are false. Which box has the gold?

Solve Gold Rush

- ▶ Propositions $M1, M2, M3$ representing messages in boxes 1,2,3
- ▶ Propositions $G1, G2, G3$ representing gold in boxes 1,2,3
- ▶ Formalize what is given to you

Solve Gold Rush

- ▶ Propositions $M1, M2, M3$ representing messages in boxes 1,2,3
- ▶ Propositions $G1, G2, G3$ representing gold in boxes 1,2,3
- ▶ Formalize what is given to you
 - ▶ $M1 \leftrightarrow \neg G1,$

Solve Gold Rush

- ▶ Propositions $M1, M2, M3$ representing messages in boxes 1,2,3
- ▶ Propositions $G1, G2, G3$ representing gold in boxes 1,2,3
- ▶ Formalize what is given to you
 - ▶ $M1 \leftrightarrow \neg G1, M2 \leftrightarrow \neg G2,$

Solve Gold Rush

- ▶ Propositions $M1, M2, M3$ representing messages in boxes 1,2,3
- ▶ Propositions $G1, G2, G3$ representing gold in boxes 1,2,3
- ▶ Formalize what is given to you
 - ▶ $M1 \leftrightarrow \neg G1, M2 \leftrightarrow \neg G2, M3 \leftrightarrow G2$

Solve Gold Rush

- ▶ Propositions $M1, M2, M3$ representing messages in boxes 1,2,3
- ▶ Propositions $G1, G2, G3$ representing gold in boxes 1,2,3
- ▶ Formalize what is given to you
 - ▶ $M1 \leftrightarrow \neg G1, M2 \leftrightarrow \neg G2, M3 \leftrightarrow G2$
 - ▶ $\neg(M1 \wedge M2 \wedge M3),$

Solve Gold Rush

- ▶ Propositions $M1, M2, M3$ representing messages in boxes 1,2,3
- ▶ Propositions $G1, G2, G3$ representing gold in boxes 1,2,3
- ▶ Formalize what is given to you
 - ▶ $M1 \leftrightarrow \neg G1, M2 \leftrightarrow \neg G2, M3 \leftrightarrow G2$
 - ▶ $\neg(M1 \wedge M2 \wedge M3), M1 \vee M2 \vee M3,$

Solve Gold Rush

- ▶ Propositions $M1, M2, M3$ representing messages in boxes 1,2,3
- ▶ Propositions $G1, G2, G3$ representing gold in boxes 1,2,3
- ▶ Formalize what is given to you
 - ▶ $M1 \leftrightarrow \neg G1, M2 \leftrightarrow \neg G2, M3 \leftrightarrow G2$
 - ▶ $\neg(M1 \wedge M2 \wedge M3), M1 \vee M2 \vee M3,$
 - ▶ $(\neg M1 \wedge \neg M2) \vee (\neg M1 \wedge \neg M3) \vee (\neg M2 \wedge \neg M3)$

Solve Gold Rush

- ▶ Propositions $M1, M2, M3$ representing messages in boxes 1,2,3
- ▶ Propositions $G1, G2, G3$ representing gold in boxes 1,2,3
- ▶ Formalize what is given to you
 - ▶ $M1 \leftrightarrow \neg G1, M2 \leftrightarrow \neg G2, M3 \leftrightarrow G2$
 - ▶ $\neg(M1 \wedge M2 \wedge M3), M1 \vee M2 \vee M3,$
 - ▶ $(\neg M1 \wedge \neg M2) \vee (\neg M1 \wedge \neg M3) \vee (\neg M2 \wedge \neg M3)$
 - ▶ Conjoin all these, and call the formula φ .

Solve Gold Rush

- ▶ Propositions $M1, M2, M3$ representing messages in boxes 1,2,3
- ▶ Propositions $G1, G2, G3$ representing gold in boxes 1,2,3
- ▶ Formalize what is given to you
 - ▶ $M1 \leftrightarrow \neg G1, M2 \leftrightarrow \neg G2, M3 \leftrightarrow G2$
 - ▶ $\neg(M1 \wedge M2 \wedge M3), M1 \vee M2 \vee M3,$
 - ▶ $(\neg M1 \wedge \neg M2) \vee (\neg M1 \wedge \neg M3) \vee (\neg M2 \wedge \neg M3)$
 - ▶ Conjoin all these, and call the formula φ .
 - ▶ Is there a unique satisfiable assignment for φ ?

Solve Gold Rush

- ▶ Propositions $M1, M2, M3$ representing messages in boxes 1,2,3
- ▶ Propositions $G1, G2, G3$ representing gold in boxes 1,2,3
- ▶ Formalize what is given to you
 - ▶ $M1 \leftrightarrow \neg G1, M2 \leftrightarrow \neg G2, M3 \leftrightarrow G2$
 - ▶ $\neg(M1 \wedge M2 \wedge M3), M1 \vee M2 \vee M3,$
 - ▶ $(\neg M1 \wedge \neg M2) \vee (\neg M1 \wedge \neg M3) \vee (\neg M2 \wedge \neg M3)$
 - ▶ Conjoin all these, and call the formula φ .
 - ▶ Is there a unique satisfiable assignment for φ ?
 - ▶ For example, is $M1 = \text{true}$ a part of the satisfiable assignment?