# Quantum Computing for Beginners: Building Qubits

Suzanne Gildert

Condensed Matter Physics Research (Quantum Devices Group)
University of Birmingham

28/03/2007

Overview of this presentation

What is a Qubit?

What are the 'basic' principles of the qubit system?

What are the design rules?

© Delft 2005

1 μm

What's so great about them?!

How are they fabricated?

How do researchers test them?

What are the applications?

UNIVERSITY OF BIRMINGHAM

What is a Qubit?

What is a Qubit?

- A qubit (or QUantum BIT) is similar in concept to a standard 'bit' - it is a memory element. It can hold not only the states '0' and '1' but a linear superposition of both states, $\alpha|0> + \beta|1>$

- In physicicts terms, we denote the states $|0>$ and $|1>$ respectively.

QUBIT vs. CLASSICAL BIT

UNIVERSITY OF BIRMINGHAM

What's so good about the Qubit?

- Qubits are useful for quantum computation

- Useful when lots of permutations need to be tried

- Suppose we want to calculate and store all possible outcomes of a particular calculation...

A classical register

| 1 |

| 1 |

| 0 0 |
| 0 1 |
| 1 0 |
| 1 1 |

Computation no. 4

A quantum register

$|0> + |1>$

$|0> + |1>$

$|00>$
+
$|01>$
+
$|10>$
+
$|11>$

SINGLE COMPUTATIONAL STEP

UNIVERSITY OF BIRMINGHAM

Choosing your Qubit

- Qubits can be realised using many different methods:
Photonic systems
Laser Ion traps / NMR systems
Quantum Dots / artificial atoms

- We are working towards Solid State Qubits, using *Superconductor* technology

There are 2 main families of Superconducting Qubits:

- Charge Qubit and Flux/Phase Qubit

It is widely believed that the only way to realise large scale integration of qubits will be with solid state technologies, in the same way that current computation can be scaled up by using transistors.

CHARGE QUBIT

1 μm
SET gate
SET
Readout circuit
Charge trap
Reservoir
Qubit
Control gates
Readout gates
© Riken 2006

FLUX QUBIT

3 μm
© Nippon 2004

UNIVERSITY OF BIRMINGHAM

What are the basic principles?

- The Josephson Junction is the basic building block of a superconducting qubit, and thus a quantum computer.
- Junction consists of 2 superconducting regions separated by a weak link (usually an insulator)

Electron wavefunction

Superconductor

Weak link

Energy diagram of a ring with a junction

$$U = \tfrac{1}{2} LI^2$$

Energy minimum

Energy

Energy maximum

0

Phase, or flux inside junction

- The qubit consists of a ring of 1 or more superconducting Josephson Junctions, with this same energy landscape.

3 μm

QUBIT

Measuring loop

UNIVERSITY OF BIRMINGHAM

Designing Qubits

- When the energy of the system is correctly chosen (by carefully choosing the capacitance and inductance of the device), quantum mechanics dominates the junction behaviour...

Energy Of Qubit Under Different Biasing Conditions.

$\frac{U}{E_J}$ Phase $\frac{\phi}{\phi_0}$ 0

$\frac{U}{E_J}$ $|\uparrow>$ $|\downarrow>$ 0.5 Phase $\frac{\phi}{\phi_0}$

© J Healey 2007

- Apply a *particular* magnetic field, and both energy states are equally favourable.

- What's more – they become quantum mechanically 'linked' through the barrier!

- The 'ground state' of the system - no currents flow. Apply a magnetic field, currents will start to flow around the loop.

- The qubit can support currents flowing in both directions at once!
- We can encode our data, |0> and |1>, into these two states.

State 0   State 1

State |0> + |1>

UNIVERSITY OF BIRMINGHAM

How do you make one?

Ramp Junction Technology

Insulating barrier

Superconductor

Gold Electrodes

- Use material deposition techniques to place a layer of superconductor

- Pattern a ramp into the superconductor

- Oxidise the surface of the superconductor (or deposit an insulating layer on top) to form the Josephson barrier

- Make a via (hole) into the insulating layer and attach electrodes to each side of the junction

- Different superconductors can be used for the two layers, to investigate interesting effects

UNIVERSITY OF BIRMINGHAM

How do you make lots?

- Single device vs. VLSI techniques

- Single device is OK for research...

- ...But VLSI needed for anything useful!

- Standard fabrication techniques in device physics/engineering to make Junctions

- Metal sputtering / Pulsed Laser deposition

- FIB etching - to produce a single device.... or an e-beam mask for multiple devices...

Quantum Computing for beginners: Building Qubits

UNIVERSITY OF BIRMINGHAM

Our fabrication plan

- We will employ a double-angle shadow technique - simple, quick and can reproduce junctions to within a small tolerance

E-beam direction x2

Shadow mask

Junctions

Superconductor regions become overlaid

Double angle shadow evaporation

- However, we will also be looking at other, more novel fabrication techniques in the meantime - there are lots!

Self-alignment
Single crystal whisker stacks?

Ramp Junctions?

UNIVERSITY OF BIRMINGHAM

Testing and measuring qubits

What temperatures do the qubits operate at?

- They need to be in the superconducting regime
- They need to be cold enough that thermal fluctuations are unimportant.
- So in the energy well diagram, the energy of the system is unlikely to be accidentally 'excited' over the barrier.

Thermal fluctuation

$|\uparrow>$ $|\downarrow>$

- The qubit sits in a copper box, which reaches the lowest temperatures – here approx. 30mK, (or 0.03 of a degree above absolute zero!)


QUBIT MYTH:
Even if you had Room Temperature Superconductors, qubits would still need to be refrigerated down to very low temperatures.

UNIVERSITY OF BIRMINGHAM

Why is it difficult to investigate qubits?

- Qubit research is difficult – you have to be really careful... 2 main problems.

  - Noise can affect your experiment – as explained earlier. Many steps taken to reduce the noise level...

    - Specialist amplifiers & comparators
    - Custom designed shielded electronics systems
    - Take many, many measurements to compile a good average
    - Great care is taken to calculate the effect of noise so that it can be eliminated

  - The measurement problem – reading a qubit without disturbing it....

  - Most people use either a SQUID loop or a resonant coil to infer the state of the qubit

UNIVERSITY OF BIRMINGHAM

A typical experimental set-up

What does it look like?

- The lower section of the apparatus is housed within a vacuum can

- Filter modules at various temperatures along the length of the cryostat

- Run apparatus from battery power and send signals through fibre optic (non electrical) links

- Fully shielded cables and fully shielded ground

0.3mm Attenuating coaxial cable

Sub 1K filter modules

Mixing chamber

Sample box

Junction

UNIVERSITY OF BIRMINGHAM

Experimental Setup (2)

Time Interval Counter

Preamplifier

IV characteristic display

Fridge Insert

Stanford Amplifier

Histogram display

Fridge dewar (He4)

Fridge control unit

UNIVERSITY OF BIRMINGHAM

What can you actually measure?

- Rabi oscillations & Ramsey fringes can illustrate the quantum coherence (showing that the two states, |0> and |1>, really are in co-existence

- Microwave spectroscopy can probe the energy levels of the qubit and help us to understand the potential of these devices


© Quantronics CEA

© University of Maryland

These experiments show very fundamental quantum properties which signify a working qubit, before trying to couple the devices together

UNIVERSITY OF BIRMINGHAM

# New experiments – our contribution

Cutting edge qubit technology!

High Tc experiments

Microwave cavity experiments

Resonator Cavity

Coaxial Cable

SMP connector

Sample - CPW

PCB

© J Healey 2007

E-Beam | Spot | Det | FWD | Tilt | 11/30/05 | Mag | 5 µm
5.00 kV | 3 | SED | 4.968 | 0.0° | 17:22:28 | 8.01 kX |

UNIVERSITY OF BIRMINGHAM

Applications
(...so can I factorise 15 with your qubits?)

- Unfotunately.....not (yet)

- May have heard of the IBM 'quantum computer' *factorising* the number 15 – the system uses an NMR technique – ensemble quantum computer

- This is currently beyond the reach of solid state quantum computers (you need at least 7 qubits)

- However solid state qubit technology is progressing very rapidly indeed, and hopefully there will be more and more results and demonstrations in the next few years

- D-wave systems 'The quantum Computing company' are in the process of producing a large scale qubit system using similar Josephson Junction technology

- Possible cryptography applications, and integrating with current optical quantum technology

IBM's 7 qubit 'molecule'

D-wave's 16 qubit device

UNIVERSITY OF BIRMINGHAM

Conclusions

- Qubits are interesting both from a physicist and an engineer's point of view, and are definitely worth investigating further.

- With enough qubits, it is possible to run quantum computations, which can outperform current technology in some specific cases, and may provide other means of replacing (or enhancing) semiconductor systems.

- The fabrication technology is improving all the time and soon qubits will be routinely implemented into superconducting circuits

- New materials are being discovered all the time which may improve the operation of the qubits further - in the same way that the semiconductor industry is progressing now.

- The applications are somewhat limited at the moment, but hopefully the field will continue to advance in the future.

UNIVERSITY OF BIRMINGHAM

Quantum computing for beginners: Building Qubits

Thank you for your attention - Any Questions?

Further Reading for interested audience members:

- *M. Tinkham 'Introduction to Superconductivity', Chapter 7*

- *D-wave systems: http://www.dwavesys.com/*

- *Some nice introductions to quantum computing:*
  ➔ *http://www.cm.ph.bham.ac.uk/scondintro/qubitsintro.html*
  ➔ *http://www.cs.caltech.edu/~westside/quantum-intro.html*
  ➔ *http://en.wikipedia.org/wiki/Qubits*
  ➔ *http://www.csr.umd.edu/csrpage/research/quantum/index.htm*

Suzanne Gildert
Condensed Matter Physics Research (Devices Group)
University of Birmingham