

exponentially in time for $K > K_c$ ($K < K_c$) (later shown rigorously and quantitatively in Ref. [16]). Crawford[22], using center manifold theory, obtains (Eq. (108) of Ref. [22]) an equation of the form $d\rho/dt = a(K - K_c)\rho + b\rho^3 + O(\rho^5)$ for K near K_c . Another work of interest is that of Watanabe and Strogatz[23] who consider the case where all oscillators have the same frequency for both finite and infinite N . By use of a nonlinear transformation of the phase variables $\theta_i(t)$, these authors show that the dynamics reduces to a solution of three coupled first order ordinary differential equations. Thus, while macroscopic behavior of order-parameter dynamics has been previously addressed for the standard Kuramoto problem, it has, until now, never been demonstrated fully (e.g., without the restriction of [22] to small amplitude, or the restriction of [23] to identical frequencies). Our paper does this and also demonstrates that our technique can be usefully applied to a host of other important related problems.

Our work also suggests other future lines of study. For example, can any rigorous results be obtained relevant to whether our macroscopic order-parameter attractors obtained by considering f in the manifold M have general validity[24]? Are there interesting qualitative differences between the behavior for Lorentzian $g(\omega)$ as compared to other monotonic symmetric oscillator distribution functions $g(\omega)$? What other systems, in addition to those discussed in Sec. IV, can our method be applied to?

We thank B. R. Hunt, M. Girvan, R. Faghih and J. Platig for very useful interactions. This work was supported by the ONR (N00014-07-1-0734) and by the NSF (PHY0456249).

-
- [1] A. Pikovsky, M. Rosenblum and J. Kürths, *Synchronization: A Universal Concept in Nonlinear Science* (Cambridge University Press, 2001); S. H. Strogatz, *Sync: The Emerging Science of Spontaneous Order* (Penguin Science Press, 2004); S. Boccaletti, J. Kürths, G. Osipov, D. L. Valladeres and C. S. Zhou, Phys. Reports **366**, 1 (2002).
 - [2] S. Yamaguchi, et al., Science **302**, 1408 (2002).
 - [3] I. Z. Kiss, Y. Zhai and J. L. Hudson, Science **296**, 1676 (2005).
 - [4] K. Wiesenfeld and J. W. Swift, Phys. Rev. E **51**, 1020 (1995).
 - [5] J. Pantaleone, Phys. Rev. D **58**, 073002 (1998).
 - [6] R. Häubler, R. Bartussek, and P. Hänggi, in *Applied Nonlinear Dynamics and Stochastic*

- Systems Near the Millennium*, edited by J. B. Kadtko and A. Bulsara, AIP Conf. Proc. No. 411 (AIP, New York, 1997) pp. 243–248.
- [7] Y. Kuramoto, in *International Symposium on Mathematical Problems in Theoretical Physics*, Lecture Notes in Physics, Vol. 39, edited by H. Araki (Springer-Verlag, Berlin, 1975); *Chemical Oscillations, Waves and Turbulence* (Springer, 1984).
 - [8] For reviews of the Kuramoto model see E. Ott, *Chaos in Dynamical Systems*, second edition, chapter 6 (Cambridge University Press, 2002); S. H. Strogatz, *Physica D* **143**, 1 (2000); and J. A. Acebron et al., *Rev. Mod. Phys.* **77**, 137 (2005).
 - [9] H. Sakaguchi, *Prog. Theor. Phys.* **79**, 39 (1988); H. Kori and A. S. Mikhailov, *Phys. Rev. E* **74**, 066115 (2006).
 - [10] T. M. Antonsen, R. Faghiih, M. Girvan, E. Ott and J. Platig, arXiv:0711.435; *Chaos* **18** (this issue) (2008).
 - [11] E. Barreto, B. Hunt, E. Ott and P. So, *Phys. Rev. E* **77**, 036107 (2008).
 - [12] E. Montbrió, J. K urths and B. Blasius, *Phys. Rev. E* **70**, 056125 (2004).
 - [13] See Theorem 8.4 of Chapter 1 of the book by E. A. Coddington and N. Levinson, *Theory of Ordinary Differential Equations* (McGraw-Hill, New York, 1955).
 - [14] This follows by noting that the Cauchy-Riemann conditions imply that the complex function $\alpha(\omega_r + i\omega_i, 0)$ satisfies Laplace’s equation in ω , $\nabla_\omega^2 \alpha(\omega_r + i\omega_i, 0) = 0$, where $\nabla_\omega^2 = \partial^2/\partial\omega_r^2 + \partial^2/\partial\omega_i^2$. Setting $\alpha = |\alpha|e^{-i\psi}$, this gives $\nabla_\omega^2 |\alpha| = |\alpha|[(\partial\psi/\partial\omega_r)^2 + (\partial\psi/\partial\omega_i)^2]$. Since $\nabla_\omega^2 |\alpha| \geq 0$ in $\omega_i < 0$, $|\alpha|$ cannot assume a maximum anywhere interior to the lower-half complex ω -plane. Hence the maximum value of $|\alpha|$ in $\omega_i \leq 0$ must occur on $\omega_i = 0$, i.e., on the real ω -axis.
 - [15] R. Mirollo and S. H. Strogatz, *J. Nonlinear Sci.* **17**, 309 (2007).
 - [16] S. H. Strogatz, R. E. Mirollo and P. C. Matthews, *Phys. Rev. Lett.* **68**, 2730 (1992).
 - [17] E. Niebur, H. G. Schuster and D. M. Kammen, *Phys. Rev. Lett.* **67**, 2753 (1991); S. Kim, S. H. Park and C. S. Ryu, *Phys. Rev. Lett.* **79**, 2911 (1997); H. Kori and Y. Kuramoto, *Phys. Rev. E* **63**, 046214 (2001); E. Montbri , D. Pazo and J. Schmidt, *Phys. Rev. E* **74**, 056201 (2006); and Ref.[20].
 - [18] M. K. S. Yeung and S. H. Strogatz, *Phys. Rev. Lett.* **82**, 648 (1999); to put (17) in the notation of this reference, make the replacement $s + i\omega_0 \rightarrow s$.
 - [19] M. Choi, H. J. Kim, D. Kim and H. Hong, *Phys. Rev. E* **61**, 371 (2000).
 - [20] B. Eckhardt, et al., *Phys. Rev. E* **75**, 021110 (2007).

- [21] Y. Kuramoto and I. Nishikawa, J. Stat. Phys. **49**, 569 (1986).
- [22] J. D. Crawford, J. Stat. Phys. **74**, 1047 (1994).
- [23] S. Watanabe and S. H. Strogatz, Phys. Rev. Lett. **70**, 2391 (1993); Physica D **74**, 197 (1994).
- [24] While the time-asymptotic behavior of the order-parameter obtained from the dynamics of the full system is seen to correspond to the attractors of the reduced systems (10) and (13) (where (13) is treated in [10]), we emphasize that M need not be attracting for the *microscopic* state $f(\omega, \theta, t)$. This can be simply seen in the case of the Kuramoto problem (4), for which $f(\omega, \theta, t) = [g(\omega)/2\pi]\{1 + \sum_m \beta_m(\omega) \exp[im(\theta - \omega t)]\}$ and $r(t) = 0$ for all t is a solution for $\beta_{\pm 1}(\omega) = 0$ and *any* choice of perturbations $\beta_m(\omega)$, $|m| \geq 2$.