
Original Paper:

(...) anatomical abnormalities included complex pelvicalyceal anatomy in 98, pelviureteric junction obstruction (pujo) in 35, ectopic, horseshoe, cross-fused ectopic kidneys in 40, and duplex system with nonfunctioning upper or lower moiety in 14 patients. (...) of the 1404 patients from rural area , 547 (39%) were managed by open surgery as compared to 362 (22%) of the patients from urban areas. (...) the mean size of the kidney stone was 5.05 5.88 cm, ureter 1.95 1.33 cm, and bladder 7.6 3.2 cm. (...) overall success rate was 90% in pcnl, 96% in urs, 100% in cystolithotripsy, and 81% in swl.

Summary by Human:*Results*

Results of 3969 surgeries, 2794 (70%) were minimally invasive surgery (mis) techniques to include swl 19%, pcnl 16 % , urs 18.9% , and pucl + pccl 16% and 1175 (30%) were open surgeries . the main factors necessitating open surgery were large stone burden 37%, anatomical abnormalities 16%, stones with renal failure 34%, gross hydronephrosis with thin cortex 58% , urinary tract infection (uti) 25%, and failed mis 18%. nearly 50% of the surgeries were necessitated by economic constraints and long distance from center where one-time treatment was preferred by the patient. tone-free rates by open surgeries were pyelolithotomy 91%, ureterolithotomy 100%, and cystolithotomy 100% with complication rate of up to 3%.

Model w/ full attn.:*Results*

results: the mean age of the patients was 6.5 years. the mean follow-up period was 12.5 months. mean stone size was 1.5cm. the majority of the stones were larger stones (n=454). the stone free rate was 100%. open surgery was performed in children with large bulk of stones, anatomical abnormalities, marked obstructive cortical atrophy and scarring, gross hydronephrosis, or urinary tract infection (uti).

Model w/ HEPOS enc-dec attn.:*Results*

the mean age of the patients was 6.35, 3.7 years with a male-to-female ratio of 2.8:1, and a range of 25 days (<1 month) to 15 years. anatomical abnormalities included complex pelvicalyceal anatomy in 98, pelviureteric junction obstruction (pujo) in 35, ectopic, horseshoe, cross-fused ectopic kidneys in 40, and duplex system with nonfunctioning upper or lower moiety in 14 patients. the mean size of the kidney stone was 5.05 5.88 cm3. of the 1404 patients from rural areas, 547 (39%) were managed by surgery as compared to 362 (22%) patients from urban areas. overall success rate was 90% in pcnl , 96% in urs , 100% in cystolithotripsy , and 81% in swl.

Figure 8: Sample summaries for a scientific paper. Model with truncated input generates fabricated facts. Our HEPOS encoder-decoder attention with LSH encoder attention are more faithful for the aspect of “results”.

Aspect	Example
Why GAO Did This Study	To protect data that are shared with state government agencies, federal agencies have established cybersecurity requirements and related compliance assessment programs. Specifically, they have numerous cybersecurity requirements for states to follow when accessing, storing, and transmitting federal data. GAO was asked to evaluate federal agencies' cybersecurity requirements and related assessment programs for state agencies. The objectives were to determine the extent to which (...)
What GAO Found	Although the Centers for Medicare and Medicaid Services (CMS), Federal Bureau of Investigation (FBI), Internal Revenue Service (IRS), and Social Security Administration (SSA) each established requirements to secure data that states receive, these requirements often had conflicting parameters. Such parameters involve agencies defining specific values like the number of consecutive unsuccessful logon attempts prior to locking out the user. Among the four federal agencies, the percentage of total requirements with conflicting parameters ranged from 49 percent to 79 percent. Regarding variance with National Institute of Standards and Technology guidance, GAO found that the extent to which the four agencies did not fully address guidance varied from 9 percent to 53 percent of total requirements. The variances were due in part to the federal agencies' insufficient coordination in establishing requirements. (...)
What GAO Recommends	GAO is making 12 recommendations to the four selected agencies and to OMB. Three agencies agreed with the recommendations and one agency (IRS) partially agreed or disagreed with them. OMB did not provide comments. GAO continues to believe all recommendations are warranted.

Table 10: Sample reference summary with aspects in a GAO report.

Aspect	Keywords	Example
Introduction and Literature	introduction, case, objectives, purposes, objective, purpose, background, literature, related work	background : the present study was carried out to assess the effects of community nutrition intervention based on advocacy approach on malnutrition status among school - aged children in shiraz , iran . introduction . low serum vitamin d levels are associated with increased postural sway . vitamin d varies seasonally . this study investigates whether postural sway varies seasonally and is associated with serum vitamin d and falls .
Methods	materials and methods, techniques, methodology, materials, research design, study design	materials and methods : this case - control nutritional intervention has been done between 2008 and 2009 on 2897 primary and secondary school boys and girls (7 - 13 years old) based on advocacy approach in shiraz , iran . the project provided nutritious snacks in public schools over a 2 - year period along with advocacy oriented actions in order to implement and promote nutritional intervention . for evaluation of effectiveness of the intervention growth monitoring indices of pre and post - intervention were statistically compared .
Results	results, experiments, observations	results : the frequency of subjects with body mass index lower than 5% decreased significantly after intervention among girls ($p = 0.02$) . however , there were no significant changes among boys or total population . (...)
Discussion and Conclusion	discussion, limitation, conclusions, concluding	conclusion : this study demonstrates the potential success and scalability of school feeding programs in iran . community nutrition intervention based on the advocacy process model is effective on reducing the prevalence of underweight specifically among female school aged children .

Table 11: Sample reference summary with aspects labeled in a PubMed article. Keywords are used to match different parts of the summaries to the four aspects.