

Example 17. A n -dimensional family of Kolmogorov complete integrable systems. Consider the polynomial functions $R = 1$,

$$\Phi(\mathbf{x}) = (x_1, x_1x_2, x_2x_3, \dots, x_{n-1}x_n)$$

and the C^r function

$$\mathbf{G}(\mathbf{u}) = (G_1(\mathbf{u}), 1 + G_2(\mathbf{u}), \dots, 1 + G_{n-1}(\mathbf{u}), 0),$$

with $\mathbf{u} \in (\mathbb{R}^n, \mathbf{0})$ and $G_i(\mathbf{0}) = 0$ for $i \geq 2$.

From the identity $\mathbf{A} \operatorname{adj}(\mathbf{A}) = \det(\mathbf{A})\mathbf{I}_n$ and by defining

$$P_{ij} := (-1)^{i+j} \prod_{\mu=1}^{j-2} x_\mu \prod_{\nu=j+1}^{n-1} x_\nu,$$

with $\prod_{\mu=1}^{j-2} x_\mu \equiv 1$ if $j \leq 2$ and $\prod_{\nu=j+1}^{n-1} x_\nu \equiv 1$ if $j \geq n-1$, it follows that

$$\operatorname{adj}(D\Phi(\mathbf{x})) = \begin{pmatrix} x_1 P_{11} & 0 & \cdots & 0 \\ x_2 P_{21} & x_2 P_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ x_n P_{n1} & x_n P_{n2} & \cdots & P_{nn} \end{pmatrix}.$$

Hence, system (3) becomes

$$\begin{aligned} \dot{x}_1 &= x_1 P_{11} G_1(\Phi(\mathbf{x})), \\ \dot{x}_2 &= x_2 (P_{21} G_1(\Phi(\mathbf{x})) + P_{22} (1 + G_2(\Phi(\mathbf{x})))), \\ &\vdots \\ \dot{x}_n &= x_n (P_{n1} G_1(\Phi(\mathbf{x})) + \sum_{j=2}^{n-1} P_{nj} (1 + G_j(\Phi(\mathbf{x})))), \end{aligned}$$

which is a C^r Kolmogorov system in $(\mathbb{R}^n, \mathbf{0})$. Since $\Phi(\mathbf{0}) = \mathbf{0}$ and $\mathbf{G}(\mathbf{0}) \neq \mathbf{0}$, this system is C^r completely integrable in $(\mathbb{R}^n, \mathbf{0})$ by Theorem 1.

ACKNOWLEDGEMENTS

The first author is supported by Dirección de Investigación of the UCSC through project DIREG 01/2024. The second author is supported by the Spanish State Research Agency, through the project PID2022-136613NB-I00 grant and the grant 2021-SGR-00113 from AGAUR, Generalitat de Catalunya. The third author is supported by European research project H2020-MSCA-RISE-2017-777911 and Universidad del Bío-Bío grant RE2320122.

REFERENCES

- [1] A. Algaba, C. García, J. Giné, *Analytic integrability around a nilpotent singularity*. J. Differential Equations **267** (2019), 443–467.
- [2] A. Algaba, C. García, M. Reyes, *Invariant curves and analytic integrability of a planar vector field*. J. Differential Equations **266** (2019), 1357–1376.
- [3] A. Algaba, C. García, M. Reyes, *A note on integrability of planar vector fields*. Eur. J. Appl. Math. (2012), 555–562.

- [4] A. Andreev, A. P. Sadovskii, V. A. Tskialyuk, *The center-focus problem for a system with homogeneous nonlinearities in the case zero eigenvalues of the linear part*. Diff. Equats. **39** (2003), 155–164.
- [5] J. Bastos, C. Buzzi, J. Torregrosa, *Orbitally symmetric systems with applications to planar centers*. Commun. Pure Appl. Anal. **20** (2021), 3319–3347.
- [6] L. Cairó, J. Llibre, *Darboux integrability for 3D Lotka–Volterra systems*. J. Phys. A: Math. Gen. **33** (2000), 2395–2406.
- [7] A. Castañeda, A. van den Essen, *A new class of nilpotent Jacobians in any dimension*. J. Algebra **566** (2021), 283–301.
- [8] F. Crespo, F. Molero, S. Ferrer, *Poisson and integrable systems through the Nambu bracket and its Jacobi multiplier*. J. Geom. Mech. **2** (2016), 169–178.
- [9] G. Darboux, *Mémoire sur les équations différentielles algébriques du premier ordre et du premier degré (Mélanges)*. Bull. Sci. Math. **2** (1878), 60–96; 123–144; 151–200.
- [10] G. Darboux, *Del’emploi des solutions particulières algébriques dans l’intégration des systèmes d’équations différentielles algébriques*. C.R. Math. Acad. Sci. Paris **86** (1878), 1012–1014.
- [11] I. A. García, H. Giacomini, M. Grau, *The inverse integrating factor and the Poincaré map*. Trans. Amer. Math. Soc. **362** (2010), 3591–3612.
- [12] A. Goriely, Integrability and Nonintegrability of Dynamical Systems, Advanced Series in Nonlinear Dynamics, vol. **19**, World Scientific Publishing Co., Inc., River Edge, NJ, 2001.
- [13] S. Labrunie, *On the polynomial first integrals of the (a, b, c) Lotka–Volterra system*. J. Math. Phys. **37** (1996), 5539–5550.
- [14] P. D. Lax, *Integrals of nonlinear equations of evolution and solitary waves*. Commun. Pure Appl. Math. **21** (1968), 467–490.
- [15] J. Llibre, R. Ramírez, V. Ramírez, *Integrability of a class of N -dimensional Lotka–Volterra and Kolmogorov systems*. J. Differential Equations **269** (2020), 2503–2531.
- [16] J. Llibre, X. Zhang, *Darboux theory of integrability in \mathbb{C}^n taking into account the multiplicity*. J. Differential Equations **246** (2009), 541–551.
- [17] J. Llibre, M. Messias, *Global dynamics of the Rikitake system*. Physica D **238** (2009), 241–252.
- [18] J. Llibre, C. Valls, *Global analytic first integrals for the real planar Lotka–Volterra system*. J. Math. Phys. **48** (2007), 033507–13.
- [19] J. Llibre, X. Zhang, *Darboux integrability for the Rössler system*. Int. J. Bifurcation Chaos Appl. Sci. Eng. **12** (2002), 421–428.
- [20] J. Llibre, X. Zhang, *Invariant algebraic surfaces of the Rikitake system*. J. Phys. A: Math. Gen. **33** (2000) 7613–7635.
- [21] P. J. Olver, Applications of Lie Groups to Differential Equations, second ed., Graduate Texts in Mathematics, vol. **107**, Springer-Verlag, New York, 1993.
- [22] T. Rikitake, *Oscillations of a system of disk dynamos*. Proc. Cambridge Philos. Soc. **54** (1958), 89–105.
- [23] O. Rössler, *An equation for continuous chaos*. Phys. Lett. A **57** (1976), 397–398.
- [24] W. Sarlet, F. Cantrijn, *Generalizations of Noether’s theorem in classical mechanics*. SIAM Rev. **23** (1981), 467–494.

DEPARTAMENTO DE MATEMÁTICA Y FÍSICA APLICADAS, UNIVERSIDAD CATÓLICA DE
LA SANTÍSIMA CONCEPCIÓN, CONCEPCIÓN, CHILE

Email address: `jgarcias@ucsc.cl`

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, BARCELONA,
SPAIN

Email address: `armengol.gasull@uab.cat`

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDAD DEL BÍO-BÍO, CONCEPCIÓN, CHILE

Email address: `srebollo@ubiobio.cl`