

Figure 16 | Performance of our ReST^{EM} optimized revision model as the sequential to parallel ratio is varied. We use majority voting to select the answer. We see that this optimized revision model demonstrates substantial performance degradations with additional sequential revisions.

L. Revision Model Example Outputs

In Figures 17, 18, 19, 20, 21, 22, and 23, we include select examples of our revision model's outputs.

M. PRM Beam Search Example Outputs

In Figures 24, 25, 26, 27, 28, and 29, we include select examples of PRM beam search. We include the PRM score, between 0 and 1, for each step in the examples.

If $f(x) = \frac{3x-2}{x-2}$, what is the value of $f(-2) +f(-1)+f(0)$? Express your answer as a common fraction.

Attempt 1:

```
Let's evaluate $f(-2)$.
We have $f(-2)=\frac{3(-2)-2}{-2-2}=-\frac{4}{-4}=\frac{1}{1}=1$.
So $f(-2)=1$.
Let's evaluate $f(-1)$.
We have $f(-1)=\frac{3(-1)-2}{-1-2}=-\frac{5}{-3}=\frac{5}{3}$.
So $f(-1)=\frac{5}{3}$.
Let's evaluate $f(0)$.
We have $f(0)=\frac{3(0)-2}{0-2}=\frac{-2}{-2}=1$.
So $f(0)=1$.
So we want to compute
$1+\frac{5}{3}+1=\frac{3}{3}+\frac{5}{3}=\frac{3+5}{3}=\frac{8}{3}$.
So the answer is $\boxed{\frac{8}{3}}$.
#### \frac{8}{3}
```

Attempt 2:

```
Let's first find $f(-2)$.
$f(-2) = \frac{3(-2)-2}{-2-2} = \frac{-6-2}{-4} = \frac{-8}{-4} = 2$.
Now let's find $f(-1)$.
$f(-1) = \frac{3(-1)-2}{-1-2} = \frac{-3-2}{-3} = \frac{-5}{-3} = \frac{5}{3}$.
Now let's find $f(0)$.
$f(0) = \frac{3(0)-2}{0-2} = \frac{-2}{-2} = 1$.
So we have $f(-2) +f(-1)+f(0) = 2+\frac{5}{3}+1 =
\frac{7}{3}+\frac{5}{3}+\frac{3}{3} = \frac{7+5+3}{3} = \frac{15}{3} = 5$.
So the answer is $\boxed{5}$.
#### 5
```

Attempt 3:

```
Let's plug in $-2$, $-1$, and $0$ for $x$ in $f(x)$.
We get $f(-2) = \frac{3(-2)-2}{-2-2} = \frac{-8}{-4} = \frac{2}{1} = 2, f(-1) =
\frac{3(-1)-2}{-1-2} = \frac{-5}{-3} = \frac{5}{3}, f(0) = \frac{3(0)-2}{0-2} =
\frac{-2}{-2} = 1$.
So we have $2+\frac{5}{3}+1=\frac{6}{3}+\frac{5}{3}+\frac{3}{3}=\frac{14}{3}$.
So the answer is $\boxed{\frac{14}{3}}$.
#### \frac{14}{3}
```

Figure 17 | Revision model example 1. The model calculates the sum at the end incorrectly on the first two attempts, but on the third attempt it succeeds and gets the answer correct.

In how many ways can 7 people sit around a round table if no two of the 3 people Pierre, Rosa, and Thomas can sit next to each other? (Seating arrangements which are rotations of each other are treated as the same.)

Attempt 1:

```
Let's start by figuring out how many ways there are to seat the three people who
can't sit next to each other.
So we need to seat them in three seats that are not right next to each other.
So there are 4 chairs that they can sit in.
So we have 4 slots to fill with 3 people.
We can do that in 4 choose 3 ways.
That is 4!/(3! * 1!) = 4 ways.
Now we need to figure out how many ways there are to seat the remaining 4 people.
So we can seat the first person in any of the seven chairs.
And then we can seat the second person in any of the remaining six chairs.
And so on.
So we can seat the four people in 7 * 6 * 5 * 4 ways.
So we need to multiply the number of ways to seat the three people who can't sit
next to each other, and the number of ways to seat the other four people.
That is 4 * 7 * 6 * 5 * 4 = 840.
So the answer is $\boxed{840}$.
#### 840
```

Attempt 2:

```
So we have to figure out how many ways there are to seat 7 people around a table
if no two of the 3 people can sit next to each other.
Let's start by seating the 3 people who can't sit next to each other.
We know that they can't sit next to each other, but they can sit on the ends of
the table.
So they can sit in 3 different ways.
Now let's seat the 4 people who can sit anywhere.
We can do that in 4! ways.
Now let's seat the 3 people who can't sit next to each other.
Let's seat them in the 3 spots we found earlier.
So there are 3 different ways to seat them.
So we have 3 * 4! * 3 = 3 * 24 * 3 = 216 ways to do it.
So the answer is $\boxed{216}$.
#### 216
```

Attempt 3:

```
So we have to find the number of ways to arrange 4 people around a round table
such that 3 of the people can't sit next to each other.
So let's start with the 3 people who can sit next to each other.
There are 3! ways to arrange them.
Now we have to arrange the other 4 people.
There are 4! ways to do that.
So the total number of ways to arrange the 7 people around the table is 3! * 4! =
6 * 24 = 144.
So the answer is $\boxed{144}$.
#### 144
```

Figure 18 | Revision model example 2. On the first attempt the model takes the incorrect approach, on the second attempt it gets closer but then makes a mistake towards the end. On the final attempt it gets to the correct answer.