

Figure C2. Same as Fig. 5, but for the mass bin $\log M/M_\odot \in [8.5, 9.0]$

$\log \nu L_{3500}$ (ergs/s)	$\log \dot{m}$	$kT_{e,w}$ (keV)	Γ_h	Γ_w	$r_{\rm hot}$	T_{hi} (K)	$[r_w(T_{hi})]$
			$\log M/M_\odot \in [7.5, 8.0]$				
[41.9, 42.3]	-1.64	†	1.60	†	520††	4913	[520]
[42.3, 42.7]	-1.79	†	1.71	†	351††	6022	[351]
[42.7, 43.1]	-1.82	†	1.90	†	292††	6719	[292]
[43.1, 43.5]	-1.64	0.97	1.62	2.20	55.3	8996	[227]
[43.5, 43.9]	-1.22	0.82	1.66	2.57	37.3	5913	[557]
[43.9, 44.3]	-0.93	0.57	1.77	3.08	18.8	5913*	[701]
[44.3, 44.7]	-0.38	0.07	1.99	2.90	11.6	5913*	[1085]
[44.7, 45.1]	-0.08	1	2.01	4.87	9.18	5913*	[1398]
			$\log M/M_\odot \in [8.5, 9.0]$				
[42.4, 42.98]	-2.26	†	1.57	†	157††	4578	[157]
[42.98, 43.56]	-2.27	†	1.73	†	131††	5209	[131]
[43.56, 44.14]	-2.15	0.19	1.86	2.45	74.4	3920	[212]
[44.14, 44.72]	-1.85	0.18	1.79	2.45	31.1	3920*	273
[44.72, 45.3]	-1.31	0.06	2.09	2.40	13.6	3920*	[422]

Table C1. Fit values for each of the luminosity bins (left column) for the stacked SEDs in the mass bin $\log_{10} M/M_\odot \in [7.5, 8.0]$ (top panel) and $\log_{10} M/M_\odot \in [8.5, 9.0]$ (bottom panel). The r_w values in the square brackets (rightmost column) show the radius where the disc temperature reaches T_{hi}.
† The warm Compton region is not required in this bin.
†† No warm Compton region is required so $r_w = r_h$, and $T_{hi} = T_{bb}(r_h)$.
∗ In these bins the standard outer disc is subdominant and T_{hi} cannot be well constrained. Hence we freeze it to the last bin where it could make an impact (i.e $[43.9, 44.3]$ for $\log_{10} M/M_\odot \in [7.5, 8.0]$ and $[43.56, 44.14]$ for $\log_{10} M/M_\odot \in [8.5, 9.0]$ respectively.)