- 1. What is the primary purpose of using histograms in graphical data analysis?
 - \circ Answer: B. To display the frequency of observations within categories of equal width
 - Histograms sort data into bins or categories of equal width to display
 the distribution of the data, particularly its central tendency, variability,
 and symmetry.
- 2. Which plot is particularly useful for assessing the fit of sample data to a theoretical distribution?
 - o Answer: D. Probability (Q-Q) Plot
 - Q-Q (Quantile-Quantile) plots compare the quantiles of a sample dataset against the quantiles of a theoretical distribution, making them particularly useful for assessing how well the sample data fit the theoretical distribution.
- 3. What does a boxplot primarily represent?
 - Answer: C. The median, interquartile range, and potential outliers in a dataset
 - Boxplots summarize the distribution of a dataset by showing its central tendency (median), variability (interquartile range), and outliers.
- 4. In the context of graphical data analysis, what advantage do quantile plots have over histograms?
 - Answer: C. Quantile plots display each data point, offering a more comprehensive view than boxplots.
 - Unlike histograms, which categorize data into bins, quantile plots display the cumulative distribution of the data, showing each data point and providing a more detailed view of the distribution.
- 5. Which of the following is NOT a correct application or characteristic of Q-Q plots based on the chapter summary?
 - Answer: B. Q-Q plots are used to evaluate the central tendency of a dataset.
 - The primary use of Q-Q plots is to assess the distributional fit of a dataset to a theoretical distribution, not to evaluate its central tendency, which is more directly assessed with measures like the mean or median.