

A boosting approach for prostate cancer detection using multi-parametric MRI

Guillaume Lemaître^{a,c} and Joan Massich^a and Robert Martí^c and Jordi Freixenet^c and Joan C. Vilanova^d and Paul M. Walker^b and Désiré D. Sidibé^a and Fabrice Mériaudeau^a

^aLE2I-UMR CNRS 6306, Université de Bourgogne, 12 rue de la Fonderie, 71200 Le Creusot, France;

^bLE2I-UMR CNRS 6306, Université de Bourgogne, Avenue Alain Savary, 21000 Dijon, France;

^cViCOROB, Universitat de Girona, Campus Montilivi, Edifici P4, 17071 Girona, Spain;

^dDepartment of Magnetic Resonance, Clinica Girona, Lorenzana 36, 17002 Girona, Spain

ABSTRACT

This document shows the desired format and appearance of a manuscript prepared for the Proceedings of the SPIE. It contains general formatting instructions and hints about how to use LaTeX. The LaTeX source file that produced this document, `article.tex` (Version 3.3), provides a template, used in conjunction with `spie.cls` (Version 3.3).

Keywords: Gradient boosting, multi-parametric MRI, prostate cancer, computer-aided diagnosis

1. INTRODUCTION

On a worldwide scale, prostate cancer (PCa) has been reported as the second most frequently diagnosed men cancers accounting for 13.6%.¹ Statistically, the estimated number of new diagnosed cases was 899,000 with no less than 258,100 estimated deaths.¹ In United States, aside from skin cancer, PCa was declared to be the most commonly diagnosed cancer among men, implying that around one in seven men will be diagnosed with PCa during their lifetime.²

Since its introduction in mid-1980s, prostate-specific antigen (PSA) is widely used for PCa screening³ and has shown to improve early detection of PCa.⁴ However, several trials conducted in Europe and United States conclude that PSA screening suffers from low specificity.⁵⁻⁷ Thus, current research focus on developing new screening methods to improve PCa detection. In this perspective, Magnetic resonance imaging (MRI) techniques have recently shown promising results for PCa detection. Furthermore, three different modalities are currently investigated: (i) T₂ Weighted (T₂-W) MRI, (ii) Dynamic Contrast-Enhanced (DCE) MRI and (iii) Apparent Diffusion Coefficient (ADC).

Several researches have been carried out in order to investigate the contributions of machine learning classifiers for PCa detection using 3T multi-parametric MRI such as Support Vector Machines (SVM),⁸⁻¹² probabilistic boosting tree¹³ or probabilistic neural network.¹³

In this research, we investigate the performance of gradient boosting for PCa detection using 3T multi-parametric MRI. Two different features extraction strategies have been chosen in order to feed the classifier: (i) voxel-based and (ii) 3D-texton-based. An evaluation of both strategies as well as the contribution of each modality is provided.

2. MATERIAL AND METHODS

2.1 Data

Here, right a paragraph on how the data were acquired and which features were extracted

Further author information: (Send correspondence to G.L.)
G.L.: E-mail: guillaume.lemaitre@udg.edu

2.2 Classification framework

2.2.1 Feature extraction strategies

- voxel-based
- 3d-texton-based

2.2.2 Gradient boosting

2.3 Validation model

k-cross validation

3. RESULTS

Include the two figures with one for voxel-based and the other one for texton-based and discuss briefly.

4. DISCUSSION

Discuss the results.

- Single modality is not working as good as multi-parametric
- Which single modality is better.
- What is the increase of the zone information.
- Voxel-based vs texton-based

5. CONCLUSION

5.1 Future works

- Check the difference with other features usually extracted.
- Check the results difference with a registration of the three modalities.

ACKNOWLEDGMENTS

This unnumbered section is used to identify those who have aided the authors in understanding or accomplishing the work presented and to acknowledge sources of funding.

REFERENCES

1. J. Ferlay, H. R. Shin, F. Bray, D. Forman, C. Mathers, and D. M. Parkin, "Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008," *Int. J. Cancer* **127**, pp. 2893–2917, Dec 2010.
2. R. Siegel, J. Ma, Z. Zou, and A. Jemal, "Cancer statistics, 2014," *CA: A Cancer Journal for Clinicians* **64**(1), pp. 9–29, 2014.
3. R. Etzioni, D. F. Penson, J. M. Legler, D. di Tommaso, R. Boer, P. H. Gann, and E. J. Feuer, "Overdiagnosis due to prostate-specific antigen screening: lessons from U.S. prostate cancer incidence trends," *J. Natl. Cancer Inst.* **94**, pp. 981–990, Jul 2002.
4. R. Chou, J. M. Croswell, T. Dana, C. Bougatsos, I. Blazina, R. Fu, K. Gleitsmann, H. C. Koenig, C. Lam, A. Maltz, J. B. Ruge, and K. Lin, "Screening for prostate cancer: a review of the evidence for the U.S. Preventive Services Task Force," *Ann. Intern. Med.* **155**, pp. 762–771, Dec 2011.
5. G. L. Andriole, E. D. Crawford, R. L. Grubb, S. S. Buys, D. Chia, T. R. Church, M. N. Fouad, E. P. Gelmann, P. A. Kvale, D. J. Reding, J. L. Weissfeld, L. A. Yokochi, B. O'Brien, J. D. Clapp, J. M. Rathmell, T. L. Riley, R. B. Hayes, B. S. Kramer, G. Izmirlian, A. B. Miller, P. F. Pinsky, P. C. Prorok, J. K. Gohagan, and C. D. Berg, "Mortality results from a randomized Prostate-cancer screening trial," *New England Journal of Medicine* **360**(13), pp. 1310–1319, 2009.

6. J. Hugosson, S. Carlsson, G. Aus, S. Bergdahl, A. Khatami, P. Lodding, C. G. Pihl, J. Stranne, E. Holmberg, and H. Lilja, "Mortality results from the Göteborg randomised population-based prostate-cancer screening trial," *Lancet Oncol.* **11**, pp. 725–732, Aug 2010.
7. F. H. Schröder, J. Hugosson, M. J. Roobol, T. L. Tammela, S. Ciatto, V. Nelen, M. Kwiatkowski, M. Lujan, H. Lilja, M. Zappa, L. J. Denis, F. Recker, A. Pez, L. Määttänen, C. H. Bangma, G. Aus, S. Carlsson, A. Villers, X. Rebillard, T. van der Kwast, P. M. Kujala, B. G. Blijenberg, U.-H. Stenman, A. Huber, K. Taari, M. Hakama, S. M. Moss, H. J. de Koning, and A. Auvinen, "Prostate-cancer mortality at 11 years of follow-up," *New England Journal of Medicine* **366**(11), pp. 981–990, 2012.
8. G. J. S. Litjens, P. C. Vos, J. O. Barentsz, N. Karssemeijer, and H. J. Huisman, "Automatic computer aided detection of abnormalities in multi-parametric prostate MRI," in *Proc. SPIE 7963, Medical Imaging 2011: Computer-Aided Diagnosis*, pp. 79630T–79630T–7, 2011.
9. G. Litjens, O. Debats, W. van de Ven, N. Karssemeijer, and H. Huisman, "A pattern recognition approach to zonal segmentation of the prostate on MRI," *Med Image Comput Comput Assist Interv* **15**(Pt 2), pp. 413–420, 2012.
10. G. Litjens, O. Debats, J. Barentsz, N. Karssemeijer, and H. Huisman, "Computer-aided detection of prostate cancer in MRI," *Medical Imaging, IEEE Transactions on* **33**, pp. 1083–1092, May 2014.
11. P. Liu, S. Wang, B. Turkbey, P. C. P. Grant, K. and Pinto, B. J. Wood, and R. M. Summers, "A prostate cancer computer-aided diagnosis system using multimodal magnetic resonance imaging and targeted biopsy labels," in *Proc. SPIE 8670, Medical Imaging 2013: Computer-Aided Diagnosis*, pp. 86701G–86701G–6, 2013.
12. Y. Peng, Y. Jiang, C. Yang, J. Brown, T. Antic, I. Sethi, C. Schmid-Tannwald, M. Giger, S. Eggener, and A. Oto, "Quantitative analysis of multiparametric prostate MR images: differentiation between prostate cancer and normal tissue and correlation with Gleason score—a computer-aided diagnosis development study," *Radiology* **267**, pp. 787–796, June 2013.
13. S. Viswanath, B. N. Bloch, J. Chappelow, P. Patel, N. Rofsky, R. Lenkinski, E. Genega, and A. Madabhushi, "Enhanced multi-protocol analysis via intelligent supervised embedding (EMPrAvISE): detecting prostate cancer on multi-parametric MRI," in *Proc. SPIE 7963, Medical Imaging 2011: Computer-Aided Diagnosis*, 2011.