

A bit of theory as far as relevant to the implementation

The necessary theory - for non-reduced axes - is found in [Ryssens et al, PHYSICAL REVIEW C 92, 064318 \(2015\)](#) section III.B Lagrange-mesh representation.

Non-reduced axes

(As from [Ryssens et al, PHYSICAL REVIEW C 92, 064318 \(2015\)](#) section III.B Lagrange-mesh representation)

Grid points

Using M evenly space grid point at distance dx , the boundaries of the box are $[-\frac{M}{2}dx, \frac{M}{2}dx]$. The width of the box is $L = Mdx$. We require $M = 2N$ to be even such that there are $N = \frac{M}{2}$ grid points on each side of the origin and the origin is avoided as a grid point.

For non-reduced axes (where the shift parameter σ is required be zero) the grid points are:
[eq 1]

$$\pm\frac{1}{2}dx, \pm\frac{3}{2}dx, \dots, \pm\frac{N-3}{2}dx, \pm\frac{N-1}{2}dx$$

or

[eq 2]

$$x_{\pm i} = \pm(\frac{1}{2} + i)dx$$
$$i = 0..N-1$$

in order:

[eq 3]

$$-\frac{2N-1}{2}dx, -\frac{2N-3}{2}dx, \dots, -\frac{3}{2}dx, -\frac{1}{2}dx, \frac{1}{2}dx, \frac{3}{2}dx, \dots, \frac{2N-3}{2}dx, \frac{2N-1}{2}dx$$

or

[eq 4]

$$x_i = -\frac{2N-1-2i}{2} = \frac{1}{2} + i - N, i = 0..2N-1$$

Basis functions

The basis functions are plane waves:

[eq 5]

$$\phi_k(x) = \frac{1}{\sqrt{L}} \exp\left(\frac{2\pi i}{L} kx\right)$$

$$k = \pm \frac{1}{2}, \pm \frac{3}{2}, \dots, \pm \frac{N-1}{2},$$

(Note that i is the imaginary unit here).

Interpolation

The Lagrange interpolation functions are:

[eq 6]

$$f_i(x) = \frac{1}{2N} \frac{\sin\left(\frac{\pi}{dx}(x - x_i dx)\right)}{\sin\left(\frac{\pi}{dx} \frac{x - x_i dx}{2N}\right)}$$

By construction, the Lagrange interpolation functions $f_i(x)$ have the property of being equal to 1 at the i -th mesh point, x_i , and 0 at all others:

[eq 7]

$$f_i(x_j) = \delta_{ij}$$

Note that the arguments of the two sine functions are the same, apart from a factor $1/2N$:

[eq 8]

$$f_i(x) = \frac{1}{2N} \frac{\sin(A_i(x))}{\sin\left(\frac{A_i(x)}{2N}\right)}, A_i(x) = \frac{\pi}{dx}(x - x_i dx)$$

This can be exploited for efficiency in a computation.

An arbitrary function $h(x)$ taking the values $h_i = h(x_i)$ on the grid points can be interpolated on an arbitrary point $x \in [-Ndx, Ndx]$ as:

[eq 9]

$$h(x) = \sum_{i=0}^{2N-1} f_i h(x_i)(x)$$

This is essentially a dot product $\mathbf{f} \cdot \mathbf{h}$.

Derivatives

The 1st and 2nd order derivatives of the Lagrange interpolation functions are:

[eq 10]

$$D_{ji}^{(1)} = \left. \frac{df_i(x)}{dx} \right|_{x=x_j} = \begin{cases} (-1)^{i-j} \frac{\pi}{2Ndx} \frac{1}{\sin(\pi(i-j)/2N)}, & \text{for } i \neq j. \\ 0, & \text{for } i = j. \end{cases}$$

$$D_{ji}^{(2)} = \left. \frac{d^2 f_i(x)}{dx^2} \right|_{x=x_j} = \begin{cases} (-1)^{i-j+1} 2 \left(\frac{\pi}{2Ndx} \right)^2 \frac{\cos(\pi(i-j)/2N)}{\sin^2(\pi(i-j)/2N)}, & \text{for } i \neq j. \\ -\frac{\pi^2}{3dx^2} \left(1 - \frac{1}{(2N)^2} \right), & \text{for } i = j. \end{cases}$$

By applying these to the expansion $\phi(x) = \sum_{i=0}^{2N-1} \phi(x_i) f_i(x)$ we obtain
[eq 11]

$$\frac{d\mathbf{h}}{d\mathbf{x}} = \frac{dh(x)}{dx} \Big|_{x=x_j} = \sum_{i=0}^{2N-1} \frac{df_i(x)}{dx} \Big|_{x=x_j} h(x_i) = \sum_{i=0}^{2N-1} D_{ji}^{(1)} h(x_i) = \mathbf{D}^{(1)} \mathbf{h}$$

So, the column vector $\frac{d\mathbf{h}}{d\mathbf{x}}$ of the derivatives of $\mathbf{h}$ at all grid points is found as a matrix product of $\mathbf{D}^{(1)}$ with the column vector of the values of $\mathbf{h}$ at all the gridpoints. As the 2nd derivative forms a matrix as well, we also have
[eq 12]

$$\frac{d^2\mathbf{h}}{d\mathbf{x}^2} = \mathbf{D}^{(2)} \mathbf{h}$$

Reduced axes

Grid points

On reduced axes only the N grid points to the right of the origin are kept:
[eq 13]

$$\frac{1}{2}dx, \frac{3}{2}dx, \dots, \frac{2N-3}{2}dx, \frac{2N-1}{2}dx$$

or
[eq 14]

$$x_i = (i + \frac{1}{2})dx$$

$$i = 0..N-1$$

(In the case of reduced axis a shift $\sigma < dx$ may be subtracted from each grid point.)

Interpolation

An arbitrary function $h(x)$, $x \in [0, Ndx]$ taking the values $h_i = h(x_i)$ on the grid points on a reduced axis can be interpolated as :
[eq 15]

$$h(x) = \sum_{i=0}^{2N-1} [h_i f_i(x) \pm h_i f_{-i}(x)]$$

where $+$, resp. $-$, is selected if $h(x)$ is symmetric, resp. antisymmetric, and $f_{-i}(x)$ is the Lagrange interpolation function corresponding to the i -th grid point to the left of the origin:
[eq 16]

$$f_{-i}(x) = \frac{1}{2N} \frac{\sin(\frac{\pi}{dx}(x - x_{-i}dx))}{\sin(\frac{\pi}{dx} \frac{x - x_{-i}dx}{2N})} = \frac{1}{2N} \frac{\sin(\frac{\pi}{dx}(x + (\frac{1}{2} + i)dx))}{\sin(\frac{\pi}{dx} \frac{x + (\frac{1}{2} + i)dx}{2N})}$$

This time we have a sum or a difference of two dot products $\mathbf{h} \cdot \mathbf{f}_+ \pm \mathbf{h} \cdot \mathbf{f}_-$

Again the arguments of the two sine functions are the same, apart from a factor $1/2N$.

Derivatives

The formula for the derivative of a function h expanded on a reduced grid is found easily by extending the column vector $\mathbf{h}$ (of length N) on the reduced grid as

[eq 17]

$$\begin{bmatrix} \pm \mathbf{g} \\ - \\ \mathbf{h} \end{bmatrix}$$

with $\mathbf{g}$ containing the same elements as $\mathbf{h}$ but in reverse order, if the function h is symmetric, and its negatives in reverse order, if it antisymmetric as if they were mirrored by a horizontal line between the two, and then forming the matrix product

[eq 18]

$$\mathbf{D} \begin{bmatrix} \pm \mathbf{g} \\ \mathbf{h} \end{bmatrix} = \begin{bmatrix} \mathbf{D}^- \\ \mathbf{D}^+ \end{bmatrix} \begin{bmatrix} \pm \mathbf{g} \\ \mathbf{h} \end{bmatrix} = \begin{bmatrix} \mathbf{D}^- \begin{bmatrix} \pm \mathbf{g} \\ \mathbf{h} \end{bmatrix} \\ - \\ \mathbf{D}^+ \begin{bmatrix} \pm \mathbf{g} \\ \mathbf{h} \end{bmatrix} \end{bmatrix}$$

The upper half of $\mathbf{D}$, $\mathbf{D}^-$, produces the derivatives on the negative x -axis, and the lower half of $\mathbf{D}$, $\mathbf{D}^+$, produces the derivatives on the positive x -axis. The upper part and the lower part are obviously related by symmetry: if the function $h(x)$ is symmetric, then the derivative is antisymmetric and *vice versa*. Thus we are only interested in the lower part. Writing $\mathbf{D}^+$ as $[\mathbf{D}_{ll} | \mathbf{D}_{lr}]$ (ll stands for lower-left and lr for lower-right quadrant of $\mathbf{D}$) we find:

[eq 19]

$$\mathbf{D}^+ \begin{bmatrix} \pm \mathbf{g} \\ \mathbf{h} \end{bmatrix} = [\mathbf{D}_{ll} | \mathbf{D}_{lr}] \begin{bmatrix} \pm \mathbf{g} \\ \mathbf{h} \end{bmatrix} = [\mathbf{E}_{ll} | \mathbf{D}_{lr}] \begin{bmatrix} \pm \mathbf{h} \\ \mathbf{h} \end{bmatrix}$$

where $\mathbf{E}_{ll}$ is obtained from $\mathbf{D}_{ll}$ by reversing the order of the rows. Reversing the order of the rows of $\mathbf{D}_{ll}$ forces us to also reverse the order of the elements of $\pm \mathbf{g}$, which then becomes, by definition $\pm \mathbf{h}$. Finally, we obtain:

[eq 20]

$$\pm \mathbf{E}_{ll} \mathbf{h} + \mathbf{D}_{lr} \mathbf{h} = [\mathbf{D}_{lr} \pm \mathbf{E}_{ll}] \mathbf{h}$$

or

[eq 21]

$$\frac{d\mathbf{h}}{dx} = [\mathbf{D}_{lr}^{(1)} \pm \mathbf{E}_{ll}^{(1)}] \mathbf{h}$$

where we have reintroduced the superscript (1) to indicate the first order derivative. By the same reasoning, we obtain for the 2nd derivative:

[eq 22]

$$\frac{d^2 \mathbf{h}}{d\mathbf{x}^2} = [\mathbf{D}_{lr}^{(2)} \pm \mathbf{E}_l^{(2)}] \mathbf{h}$$

N -dimensional grids

The full Cartesian 3D representation of a function $h(\mathbf{r})$, where $\mathbf{r} = [x \ y]$ (2D case), or $\mathbf{r} = [x \ y \ z]$ (3D case), is then provided (for the 3D case) by [eq 23]

$$h(\mathbf{r}) = \sum_{ijk} h_{ijk} f_i(x) f_j(y) f_k(z)$$

where the number of discretization points does not have to be the same in each direction. In this case, the derivative matrices $\mathbf{D}^{(1)}$ and $\mathbf{D}^{(2)}$ have to be set up separately for each direction, taking into account whether the axis is reduced or not.

Basis functions

The basis functions are plane wave products of the different axes:
[eq 24]

$$\begin{aligned} \Phi_{klm}(x, y, z) &= \phi_k(x) \phi_l(y) \phi_m(z) = \frac{1}{\sqrt{L_x}} \frac{1}{\sqrt{L_y}} \frac{1}{\sqrt{L_z}} \exp\left(\frac{2\pi i}{L_x} kx\right) \exp\left(\frac{2\pi i}{L_y} ly\right) \exp\left(\frac{2\pi i}{L_z} mz\right) \\ &= \frac{1}{\sqrt{L_x L_y L_z}} \exp\left(2\pi i \left(\frac{kx}{L_x} + \frac{ly}{L_y} + \frac{mz}{L_z}\right)\right) \\ &== \frac{1}{\sqrt{L_x L_y L_z}} \exp(2\pi i (\mathbf{k} \cdot \mathbf{r})) \\ \mathbf{k} &= \left[\frac{k}{L_x} \quad \frac{l}{L_y} \quad \frac{m}{L_z} \right] \\ \mathbf{x} &= [x \ y \ z] \end{aligned}$$