SV Evaluation

The figure in the geographical domain are sketched at the Earth's mean radius.

The models are analyzed in the following order:

1:'JAPAN','

2:NCEI',

3:'IPGP',

4:'BGS',

5:'Leeds',

6:'ISTerre',

7:'MPS',

8:'Strasbourg',

9:'GSCF',

10:'DTU',

11:'MPS',

12:'IZMIRAN',


13:'UCM',

14: 'GFZ'


The last column is the arithmetic mean model. Attached is the matrix of RMS between all models corresponding to Ciaran's plot (in a different order).

Summary: The histogram of RMS with respect to the arithmetic mean model indicates two groups of models. The candidates most different from the rest of the population are the STRASBOURG, GSFC and JAPAN models. For Strasbourg the difference is mostly on the degree 1 and more specifically on the g10. For JAPAN and GSFC the difference is distributed mostly on degrees 2 and 3. MPS, NCEI and IZMIRAN have positive residuals over Indonesia while most of others have negative residuals. Such structures are in general carried by the n=m harmonics, with n=2 in this case.


The power spectra of the difference to the mean are:



The azimuthal power spectrum is:




The coefficient difference is:




And the differences in the geographical domain for the radial component of each candidate model (at the Earth's mean radius) is:


JAPAN:




NCEI:




IPGP:




BGS:




LEEDS:




ISTERRE :




MPS :




STRASBOURG:




DTU:




MPS :



IZMIRAN :



UCM :



GFZ :

