

EEG Spectral Components Ratio Results and Discussion

In this study, we recruited 10 participants for the TloadDback experiment. Data collected from 0–20 minutes were classified as non-fatigue data, whereas those from 41–60 minutes were classified as fatigue data. These data segments were then used to extract EEG features, followed by PSD computation to obtain the band power (BP) characteristics. Subsequently, the BP features were compared and interpreted. Figures 1 to 10 illustrate the BP topographic maps for the non-fatigue and fatigue conditions across different frequency bands in all 10 participants. The differences in frequency band feature intensities between the non-fatigue and fatigue states for each participant will be described in the following sections.




Figure 1 Subject No. 1 frequency band power characteristics (a) non-fatigue characteristics 0 to 20 minutes (b) fatigue characteristics 41 to 60 minutes

Figure 1 shows the experimental results for Subject 1. In the θ/α ratio topographic map for this subject, we observe that power in the frontal and bilateral

temporal regions increases under the fatigue state. This elevation in θ -wave power may be significantly related to the onset of fatigue and drowsiness. In contrast, β -wave power increases more than α -wave power, leading to a decrease in the α/β ratio in the frontal area. This rise in β -wave power appears to be associated with sustained alertness in a prolonged concentration state.




Figure 2 Subject No. 2 frequency band power characteristics (a) non-fatigue characteristics 0 to 20 minutes (b) fatigue characteristics 41 to 60 minutes

Figure 2 shows the experimental results for Subject 2. In the θ/α ratio topographic map, an increase in frontal lobe intensity is observed under the fatigue condition, whereas the frontal lobe energy in the α/β ratio decreases. Additionally, from the $(\theta + \alpha)/(\alpha + \beta)$ map, both hemispheres originally show relatively high intensities, but after entering the fatigue state, the right hemisphere's intensity decreases

more noticeably than the left. A rise in α -wave power is also evident; however, the increase in alertness contributes to some growth in β -wave power, which leads to a decrease in the frontal α/β ratio.




Figure 3 Subject No. 3 frequency band power characteristics (a) non-fatigue characteristics 0 to 20 minutes (b) fatigue characteristics 41 to 60 minutes

Figure 3 presents the experimental results for Subject 3. In the θ/α topographic map, the frontal region shows increased intensity during fatigue, whereas frontal energy in the α/β ratio decreases. The growth of α -wave power in the frontal area is weaker than that of the β -wave (which rises due to heightened alertness). Additionally, compared with the $(\theta + \alpha)/\beta$ feature, the $(\theta + \alpha)/(\alpha + \beta)$ feature shows a noticeable decrease in parietal lobe energy.




Figure 4 Subject No. 4 frequency band power characteristics (a) non-fatigue characteristics 0 to 20 minutes (b) fatigue characteristics 41 to 60 minutes

Figure 4 shows the experimental results for Subject 4. Using the θ/α and α/β features, it is evident that slow-wave activity (θ and α) increases in the frontal region, while the β -wave exhibits enhanced power, likely due to high alertness during prolonged tasks. Furthermore, both the $(\theta + \alpha)/(\alpha + \beta)$ and $(\theta + \alpha)/\beta$ features display relatively high intensities overall.




Figure 5 Subject No. 5 frequency band power characteristics (a) non-fatigue characteristics 0 to 20 minutes (b) fatigue characteristics 41 to 60 minutes

Figure 5 presents the experimental results for Subject 5. In this subject's topographic map, the θ/α ratio shows increased power in the left frontal region but decreased power in the right prefrontal region. Additionally, the right occipital area exhibits higher power across multiple features, potentially due to the prolonged visual demand on the occipital lobe. Meanwhile, the α/β ratio also decreases in the prefrontal region, which may be related to a pronounced increase in β -wave power arising from an elevated state of alertness.




Figure 6 Subject No. 6 frequency band power characteristics (a) non-fatigue characteristics 0 to 20 minutes (b) fatigue characteristics 41 to 60 minutes

Figure 6 shows the experimental results for Subject 6. In this subject's topographic map, most features appear relatively unstable. Specifically, in the $(\theta + \alpha)/(\alpha + \beta)$ and $(\theta + \alpha)/\beta$ features, the initial intensity is primarily concentrated in both temporal lobes. However, when fatigue sets in, energy distribution shifts to the entire brain, resulting in an overall increase in intensity. Regarding the θ/α ratio, the non-fatigue state shows a more irregular distribution across the brain. Once fatigue occurs, the activity becomes mainly focused in the frontal region, indicating an overall increase in slow-wave activity (i.e., elevated θ -wave power associated with drowsiness).




Figure 7 Subject No. 7 frequency band power characteristics (a) non-fatigue characteristics 0 to 20 minutes (b) fatigue characteristics 41 to 60 minutes

Figure 7 presents the experimental results for Subject 7. In this subject's topographic map, the $(\theta + \alpha)/\beta$ feature exhibits a synchronous increase in power across the entire brain region. Meanwhile, the α/β feature follows a pattern similar to most subjects, showing a decrease in frontal energy after entering the fatigue state—likely due to increased β -wave activity. However, the θ/α feature indicates elevated θ -wave power in the temporal region under fatigue, suggesting that this subject experienced both drowsiness and heightened alertness simultaneously.




Figure 8 Subject No. 8 frequency band power characteristics (a) non-fatigue characteristics 0 to 20 minutes (b) fatigue characteristics 41 to 60 minutes

Figure 8 shows the experimental results for Subject 8. In the non-fatigue state, the θ/α , θ/β , and α/β features remain relatively stable across the entire scalp, with no pronounced channels. Upon entering the fatigue state, a rise in frontal θ/α power is observed, attributable to increased θ -wave activity caused by drowsiness, while a decline in frontal α/β power indicates reduced α -wave activity and elevated β -wave activity. This suggests that in the later stages of the experiment, the subject gradually overcame drowsiness and actively worked to maintain alertness.




Figure 9 Subject No. 9 frequency band power characteristics (a) non-fatigue characteristics 0 to 20 minutes (b) fatigue characteristics 41 to 60 minutes

Figure 9 presents the experimental results for Subject 9. In this subject's topographic map, the θ/α feature shows a marked increase in the anterior regions of the brain, while α/β decreases in intensity. A concurrent decline in $(\theta + \alpha)/\beta$ is also observed, suggesting that β -wave power may have increased due to heightened alertness from prolonged concentration.




Figure 10 Subject No. 10 frequency band power characteristics (a) non-fatigue characteristics 0 to 20 minutes (b) fatigue characteristics 41 to 60 minutes

Figure 10 shows the experimental results for Subject 10. After entering the fatigue state, all three features indicate a chaotic distribution across the brain, implying that fatigue may disrupt overall cognitive functioning. In the frontal region, the θ/α ratio increases whereas α/β decreases, suggesting simultaneous rises in both α - and β -wave activity—although β -wave intensity grows more prominently to maintain alertness and counteract fatigue.

In the integration of the five BP characteristics of the 10 subjects mentioned above, it can be found that most of them produce similar states, such as: 1. In the θ/α characteristic, it can be seen that the frontal lobe and both sides of the temporal lobe increase in intensity during the state of fatigue; 2. In the α/β , the prefrontal energy decreases, and it can be inferred that the beta wave increases significantly, which is

likely to be related to the alertness of the long-term focus on performing tasks; 3. In the θ/β characteristic, there is no obvious change in θ/β characteristic, which suggests that there is consistency in the size and direction of changes in θ and β wave activities, which means that there will be a state of high sleepiness and high alertness at the same time when the subject is fatigued. There is no significant change in θ/β characteristics before and after fatigue, which suggests that the size and direction of the changes in θ and β wave activities are consistent, which means that when fatigue occurs, the subjects will be in a state of high sleepiness and high alertness at the same time. The results are consistent with some of the characteristics mentioned in the literature, such as 1. the increase in θ -wave energy and the slight increase in α -wave energy, and the predominance of slow-wave activity in the overall fatigue state[1]; and 2. the increase in β -wave power is associated with psychological fatigue and alertness when mental fatigue occurs[2][3][4].

Although similar phenomena were found in most subjects, a few subjects did not respond in the same way to this experiment, and the brain intensity confusion after fatigue was initially due to the long duration of the experiment, which prevented subjects from effectively recovering their brain function after entering the fatigue state, and caused their load level to be overloaded, which in turn affected the presentation of the characteristics. How to effectively adjust the experiment duration is also a major

issue for the future.

References

- [1] A. M. Strijkstra, D. G. M. Beersma, B. Drayer, N. Halbesma, S. Daan, Subjective sleepiness correlates negatively with global alpha (8-12 Hz) and positively with central frontal theta (4-8 Hz) frequencies in the human resting awake electroencephalogram, *Neuroscience Letters*, 340 (1) (2003), 17-20. [https://doi.org/10.1016/S0304-3940\(03\)00033-8](https://doi.org/10.1016/S0304-3940(03)00033-8).
- [2] S. K. Lal, A. Craig, A critical review of the psychophysiology of driver fatigue. *Biological Psychology*, 55(3) (2001), 173-194. [https://doi.org/10.1016/S0301-0511\(00\)00085-5](https://doi.org/10.1016/S0301-0511(00)00085-5).
- [3] X. Fan, Q. Zhou, Z. Liu, F. Xie, Electroencephalogram assessment of mental fatigue in visual search, *Bio-medical materials and engineering*, 26(s1) (2015), s1455–s1463. <http://dx.doi.org/10.3233/BME-151444>.
- [4] A. Craig, Y. Tran, N. Wijesuriya, H. Nguyen, Regional brain wave activity changes associated with fatigue. *Psychophysiology*, 49(4) (2012), 574–582. <https://doi.org/10.1111/j.1469-8986.2011.01329.x>.