

EXPLORING CRITICAL CONFORMATIONS:
STATE SEARCHING AND SAMPLING IN BOTH
GERMANIUM CHAINS AND ICE INTERFACES
{EARLY DRAFT}

By

GENTRY H. SMITH

B.S.

Southern Nazarene University

Bethany, OK, USA

2016

Submitted to the Faculty of the
Graduate College of
Oklahoma State University
in partial fulfillment of
the requirements for
the Degree of
Master of Science
November 2018

EXPLORING CRITICAL CONFORMATIONS:
STATE SEARCHING AND SAMPLING IN BOTH
GERMANIUM CHAINS AND ICE INTERFACES
{EARLY DRAFT}

Thesis Approved:

Thesis Advisor

ACKNOWLEDGMENTS

Acknowledgments reflect the views of the author and are not endorsed by committee members or Oklahoma State University.

ACKNOWLEDGMENTS

To Oklahoma State University, for providing the environment in which I have been able to study, teach, and research.

To the HPCC and the individuals within (**get first/last names**) for providing a powerful cluster for computations and continuous support for technical issues.

To my advisor, who instructed and assisted me in research

To my parents, by blood and marriage, who have always encouraged me toward higher goals.

To my wife, Miranda, who has supported me for over 5 years.

Acknowledgments reflect the views of the author and are not endorsed by committee members or Oklahoma State University.

Name: GENTRY H SMITH

Date of Degree: November 2018

Title of Study: EXPLORING CRITICAL CONFORMATIONS

Major Field: COMPUTATIONAL CHEMISTRY

Abstract: Molecular conformation plays a critical role in the properties of systems in either the condensed or vapor states. The ensemble of conformations dictates structural properties, energies, heat capacities, and other thermodynamic and dynamic quantities. Here, we explore the role of conformation in proton ordering and orientational defect formation in ice as well as strategies for exhaustive conformer searching for molecules using Group IV element backbones. In the ice systems, we show algorithmic strategies for seeking optimized proton disordered crystals that satisfy the Bernal-Fowler ice rules. In the Group IV molecule investigations, we develop an automated strategy for seeking the optimal low energy conformer and uncover previously unreported deficiencies in common computational software used in investigating Germanium complex energies.

TABLE OF CONTENTS

Chapter	Page
1 Introduction	1
1.1 Reason for Study	1
1.1.1 Generation of Ice I_h Crystal Structure	2
1.1.2 Conformation Landscapes of Group IV Chains	2
1.1.3 Ice Interfaces in Two Dimensions	2
1.2 Ice Annealing	3
1.2.1 Bernal - Fowler Ice Rules	3
1.2.2 Structures of Ice	3
1.2.3 Residual Entropy of Ice I_h and Ice XI	4
1.2.4 Hydrogen Bond Defects in Ice Crystals	4
1.2.5 Literature Review on Relevant Works	4
1.3 Conformation Landscapes	4
1.3.1 A Brief History of Conformation Landscapes	5
1.3.1.1 Levinthal's Paradox	5
1.3.2 Computational Modeling	5
1.3.2.1 History of Chemical Modeling	5
1.3.2.2 Hardware: Oklahoma State University's Cowboy Cluster	5
1.3.2.3 Software	5
1.3.2.4 Programming Languages	6
1.3.3 Literature Review on Relevant Works	6
1.4 Modeling Germanium Compounds	6
1.4.1 Tools for Modeling Germanium	6
1.4.2 Computational Complexity of Germanium Compounds	6
1.4.3 Literature Review on Relevant Works	6
1.5 Two-Dimensional Water	7
1.5.1 Lennard-Jones Potential	7
1.5.2 Modeling Water in Two Dimensions	8
1.5.2.1 Mercedes-Benz Model	8
1.5.2.2 Rose Potential Model	9
1.5.2.3 Two-Dimensional Modeling	9
1.5.3 Literature Review on Relevant Works	9
2 On Algorithms for Building and Sampling Disordered Crystal States	10
2.1 States and Properties of Ice	10
2.1.1 Bernal-Fowler Ice Rules	10

2.1.2	Forms of Ice	10
2.1.3	Ice I_h	11
2.1.4	Efforts to Generate Ice I_h	11
2.1.5	Comparison between Ice XI and Ice I_h	11
2.2	Method Design	12
2.2.1	Overview	12
2.2.2	Selection of Software Tools	12
2.2.3	Generation of Source Ice XI	12
2.2.4	Source Ingestion	12
2.2.5	Identifying Neighboring Molecules	13
2.2.6	Defining Tetrahedral Positions	13
2.2.7	Pseudorandom Rearrangement of Water Molecules	14
2.2.8	Detecting Hydrogen Bond Defects	15
2.2.9	Correcting Hydrogen Bond Defects	15
2.3	Results of Method	16
2.4	Comparison with Other Methods	18
2.4.1	benefits of own method over others	18
2.4.2	benefits of other methods over this	18
2.5	Comments on Limitations and Proposed Improvements	18
3	Crystal and Liquid 2D Water at Interfaces	19
3.1	Two-Dimensional Water	19
3.1.1	Lennard-Jones Potential	19
3.1.2	Modeling Water in Two Dimensions	20
3.1.2.1	Mercedes-Benz Model	20
3.1.2.2	Rose Potential Model	21
3.1.2.3	Two-Dimensional Modeling	21
3.1.3	Literature Review on Relevant Works	22
3.2	Goal of Project	22
3.3	Tools and Terms	22
3.3.1	OOPSE in 2D	22
3.3.2	Reduced Terms: 2D analogues	22
3.4	Designing System	23
3.4.1	Defining the Surface	23
3.4.1.1	Manipulation of LJ Potential	23
3.4.1.2	Manipulation of bead spacing	23
3.5	Results	23
3.5.1	Future Work	23
4	Germanium Compounds and QM Concerns	24
4.1	Modeling Germanium Compounds	24
4.1.1	Tools for Modeling Germanium	24
4.1.2	Computational Complexity of Germanium Compounds	24
4.1.3	Literature Review on Relevant Works	24
4.2	The Initial Problem: Germanium Study	25

4.2.1	Parameters of Work and Previous Collaborator's Results . . .	25
4.2.2	Design and Approach to Solution	28
4.2.2.1	Design 1: Occam's Smallest Razer	28
4.2.2.2	Design 2: A Blunt Effort	29
4.2.2.3	Design 3: Death by 1.59 Million Cuts	30
4.2.3	Scale Reduction Efforts	32
4.2.4	junk section	33
4.2.5	Attempts at Simplification	33
4.2.6	Discovery of Program Flaw	33
5	Sampling Conformation Landscapes by Rotatable Bond Degrees of Freedom	34
5.1	Introduction to Topic	34
5.1.1	A Brief History on Conformation Landscapes	34
5.1.1.1	Levinthal's Paradox	34
5.1.2	Computational Modeling	34
5.1.2.1	History of Chemical Modeling	34
5.1.2.2	Hardware: Oklahoma State University's Cowboy Cluster	35
5.1.2.3	Software	35
5.1.2.4	Programming Languages	35
5.1.3	Literature Review on Relevant Works	35
5.2	Design of System	37
5.2.1	Variation of Theory and Basis Set Usage by System Size and largest atom type	38
5.2.2	Computational Optimization by Varying Resolution	38
5.2.3	Inherent Complications	38
5.3	Results	38
5.3.1	Problems	38
5.3.2	Anticipated Approaches for Future Work	39
	References	40
A	Ice Ih to Ice XI Conversion	41
A.1	Code: PDBDisorganize.py	41
B	Conformation Landscapes	53
C	Germanium Landscape	54
C.1	Code: hexagermane-transall.pdb	54
C.2	Code: ge4h.pdb	63
D	Two-Dimensional Rose-Potential Water	65

LIST OF TABLES

Table		Page
4.1	Collaborator's Hexagermane Energies by Conformation (density functional theory, unknown basis set, energy in Hartrees and KJ/mol) . .	25

LIST OF FIGURES

Figure		Page
2.1	"Before" image of Ice XI	17
2.2	"After" image of generated ice Ih	17
4.1	Fully trans configuration of pentagermanium-based compound.	26
4.2	Fully trans configuration of hexagermanium-based compound.	27
5.1	example VR chart (exploding like molecule/cell/fiber/muscle)	36
5.2	Flow of method design for variable resolution conformation landscape search.	37

NOMENCLATURE

Variables

ϵ	Lennard-Jones Potential well depth
r	Lennard-Jones Potential distance between center of two particles
σ	Lennard-Jones Potential intermolecular contact distance
V	Lennard-Jones Potential intermolecular potential

Subscripts/Superscripts

0	Initial condition
---------	-------------------

CHAPTER 1

Introduction

I want a first paragraph that inspires the reader to continue reading. Maybe something with a quote or a question. Maybe not.

The idea for an introduction is to interest the reader and provide general background information. For my work, the interesting part is how impactful computational science (specifically chemistry) has been in research and on society. The background information will start extremely generic and then go into some overarching themes. Generically, I'll include the development of computational science in the 20th century and the vast applications of computational chemistry specifically. Overarching themes are related, but don't have to be explicitly relevant. For example, Levinthal's paradox is a fun example showing the problem of conformation landscape searches. Also,

1.1 Reason for Study

The studies conducted that comprise this work were determined as a combination of collaborative efforts and larger research group goals with novel discoveries worth reporting. Collectively, they explore conformations of internal bond dihedrals, molecule orientations of microstates, and properties of interfacing substrates. These efforts are categorized and separated into three categories: ice crystal states, conformation landscapes, and ice interfaces. A brief introduction of each and a literature review of relevant information is given below.

1.1.1 Generation of Ice I_h Crystal Structure

Ice crystals can take many forms based on properties like temperature and pressure. The proton-disorganized orthorhombic form of ice known as I_h is the form of ice most commonly found on earth (general understanding, CITATION NEEDED). Due to the inherent randomness of the disorganization of the molecules within the crystal, computational efforts are limited in scope or instead utilize the proton-ordered orthorhombic form of ice XI (general understanding, CITATION NEEDED). This project explores a method to produce a high quality pseudorandom ice I_h crystal structure.

1.1.2 Conformation Landscapes of Group IV Chains

Any molecule with a chain length of at least four contains at least one dihedral. In small molecules, the steric hindrance between the head and tail atoms are usually minimized in the fully gauche conformation to produce the lowest-energy conformer. In larger and more bulky molecules, additional interactions may cause the dihedral to take other conformations in search of the lowest-energy conformation. This project details the search for the lowest-energy conformer of a bulky hexagermane molecule in collaboration with Oklahoma State University’s Charles Weinert and the complications and curiosities found within.

1.1.3 Ice Interfaces in Two Dimensions

Ice is capable of interfacing with hydrophobic and amphipathic molecules. These interfaces often have a specific range of polarity and **physicochemical** properties like atom type, charge, and the spacing of interfaces. While usually considered in the biochemical sense, molecular interfaces with water have additional applications. If, for example, an interface had the properties to encourage ice growth, then it may be possible to observe significant ice growth above the expected freezing temperature.

Modeling water systems can be computationally intensive and require significant resources or a limited scope of system. One proven method of simplifying water modeling is to reduce the system by one dimension, creating a two-dimensional water. The 'Mercedes Benz' BN2D model and the more recent rose potential model are two examples of two-dimensional water models. Exploring and studying a two-dimensional water system allows for computationally-efficient investigating and can provide relevant information for a similar simulation with three-dimensional water. This project covers the usage of the two-dimensional rose potential model to observe ice growth on a substrate with varying properties.

1.2 Ice Annealing

EACH SUBSECTION: DEFINITION OF TERMS

1.2.1 Bernal - Fowler Ice Rules

Citation! Bernal and Fowler (1933)

There have been many works published on this topic, so I will have no problem obtaining a cohesive review. I plan to include the various forms of ice and focus on I_h and XI. This includes differences between the two in prevalence, environment, and synthesis (is synthesis the correct term here?).

I expect to focus on many thermodynamic properties and to focus on entropy as a driving force of difference in the modeled system. I will include publications of the effort to model a truly proton-disordered system and focus on the computational aspect.

1.2.2 Structures of Ice

Ice contains many structures. These structures are typically orthorhombic, which means rectangular at non-90 angles. They can also be hexagonal, cubic, else. These

different structures form based on temperature and pressure. Ice I_h is the most prevalent form of ice found normally on earth, forming at pressures around 1 atm and temperatures around 0C. (source needed?).

More detail on ice I_h .

More detail on ice XI.

1.2.3 Residual Entropy of Ice I_h and Ice XI

Residual entropy is a thermodynamic property that greatly differs between ice I_h and XI. In general, entropy can be calculated for a system of N molecules as $S = Nk \ln(w)$, where k is the Boltzmann constant and w is the number of real microstates corresponding to any macrostate. Residual entropy differs in calculation from entropy in that it generally refers to the entropy of a crystal near zero kelvin. Linus Pauling (1935) described the w of very low temperature crystals to approach the number of orientations possible for each molecule with consideration to immediate neighbors. Since ice I_h is a

For a proton-disordered ice I_h crystal, w becomes $\frac{x}{24}$ where x is the number of acceptable orientations within the crystal.

1.2.4 Hydrogen Bond Defects in Ice Crystals

Speak here about defects and how they quantitatively harm stability and why the defects need to be reduced in general ice I_h structures.

1.2.5 Literature Review on Relevant Works

1.3 Conformation Landscapes

EACH SUBSECTION: DEFINITION OF TERMS

For ROUGHLY forty years, computational programs have allowed investigators to model chemical systems with high accuracy to determine their physical properties.

1.3.1 A Brief History of Conformation Landscapes

1.3.1.1 Levinthal's Paradox

Discuss history of Levinthal and his paradox. Provide the non-paradoxical solution.

Next: Levinthal golf courses by Ken Dill.

1.3.2 Computational Modeling

Introduce importance and impact. Bigly important.

1.3.2.1 History of Chemical Modeling

Here I will introduce molecular modeling. This will begin with a brief history of the development of the field. It will continue through to mention the styles and goals of molecular modeling. Upon reaching modern techniques, I will discuss the benefits and costs associated with the major types of calculations (QM, MD, MC, etc).

1.3.2.2 Hardware: Oklahoma State University's Cowboy Cluster

It would also be appropriate to mention the computational capabilities of OSU's Cowboy cluster.

1.3.2.3 Software

Software to mention:

Visualizing Avogadro, UCSF Chimera(?),

Computing GAMESS(?), Gaussian, NWChem(?), OOPSE

This also includes brief pros and cons about the programs and the general purpose of use in case.

1.3.2.4 Programming Languages

Another hugely important portion of this will include a choice in the programming languages used (mostly python (Cython-compiled!), some Perl and Bash).

1.3.3 Literature Review on Relevant Works

1.4 Modeling Germanium Compounds

EACH SUBSECTION: DEFINITION OF TERMS

This will be an interesting section as there is extremely little in terms of Ge computational work. Perhaps a broader search will yield interesting results. For sake of thoroughness, I will also include work on computational energy optimization in general and work through complications brought by the size of Ge. I might also include a portion on the statistical spread of conformations at a given temperature (internal energy?) I may include a sentence or paragraph on Gaussian-based publications.

1.4.1 Tools for Modeling Germanium

Computational Requirements and reasons for those requirements.

Germanium is not the most-studied atom in computational works. The majority of Germanium studies are done with Gaussian (citation needed?).

1.4.2 Computational Complexity of Germanium Compounds

Draw-backs of modeling Germanium. Uncommon but still necessary for wetwork.

1.4.3 Literature Review on Relevant Works

Make note of various Germanium modeling research. Make note of tools and methods used.

1.5 Two-Dimensional Water

EACH SUBSECTION: DEFINITION OF TERMS

Rose water is fairly new on the computational scene and so I may also include a review on the Mercedez-Benz water system as well as any other attempts to model water in two dimensions. For the rose potential system, I will review the Lennard-Jones potential as well as any other equations/systems related to the rose potential.

1.5.1 Lennard-Jones Potential

The Lennard-Jones Potential well is a soft-sphere model of interaction between two spheres described with

$$V_{LJ} = 4\epsilon[(\frac{\sigma}{r})^{12} - (\frac{\sigma}{r})^6] \quad (1.1)$$

where V is the potential, r is the distance between the center of two particles, σ is the specific distance between the two particles where the potential is zero, and $-\epsilon$ is the minimum potential of the plot. **REFINE WORDING:** The plot is defined in $[0, \infty)$. As two particles approach from infinity, their interaction becomes negative - which is an attractive force - and will approach the global minimum of $-\epsilon$. The r of this interaction is slightly larger than the combined radii of the two particles - which means they aren't quite touching - and is the equilibrium distance between the two particles. As r decreases beyond the minimum and toward σ , the interaction strength increases and reaches zero as $r = \sigma$. At $r < \sigma$,

Potential digression: In a "hard-sphere" model, a particle's radius is firm, which is to say that the interaction potential is infinite at r less than σ . Basically a vertical line between two discrete values (usually ϵ and ∞) as the potential shifts from $r \geq \sigma$ to $r < \sigma$ (maybe include image?). The Lennard-Jones potential is a "soft-sphere" model, which blurs the line and replaces the vertical line with a functional representation.

This breaks with reality as the particles become "squishy" and the potential ramps up toward infinity as r decreases. The benefit to the soft-sphere model is that modeling programs can more-easily account for overlaps in particles during time steps than with hard-sphere models. For example, a hard-sphere model of two particles interacting will likely not have a position where $r = \sigma$ and will potentially overlap. At this overlap, the potential is infinity and will introduce a nearly infinite force at that instant of time. Computer systems do not like having infinitely large repulsions suddenly introduced into a simulation.

1.5.2 Modeling Water in Two Dimensions

Modeling in two dimensions sacrifice the "realism" of models in three dimensions, but reduce the computational load significantly. This allows researchers (scientists, chemists, digital magicians?) to test more simple designs in two dimensions as well as a higher volume of simulations at the same time/computational cost.

1.5.2.1 Mercedes-Benz Model

The "Mercedes-Benz" BN2D model of water first proposed by Ben-Naim (1971) as "waterlike particles" are a popular two-dimensional representation of water. ROUGH: details of shape of MB water

The mathematical model used in the BN2D model is generated from the Percus-Yevick equation by substituting the approximation

$$c(X_1, X_2) = y(X_1, X_2)f(X_1, X_2) \quad (1.2)$$

into the Percus-Yevick equation obtained from the Ornstein-Zernike relation

$$h(X_1, X_2) = c(X_1, X_2) + \frac{\rho}{2\pi} \int c(X_1, X_3)h(X_3, X_2)dX_3 \quad (1.3)$$

to produce the overall relation

$$y(X_1, X_2) = 1 + \frac{\rho}{2\pi} \int y(X_1, X_3) f(X_1, X_3) \times \left[y(X_3, X_2) f(X_3, X_2) + y(X_3, X_2) - 1 \right] dX_3 \quad (1.4)$$

1.5.2.2 Rose Potential Model

The rose potential is another model first introduced by Williamson et al. (2017). This model, while similar to the three-pronged BN2D, is notably different in that the rose potential model simplifies the model by use of a radial sinusoidal plot to make the three "prongs" of the particle.

1.5.2.3 Two-Dimensional Modeling

Something other than OOPSE? (not seeing obvious answer other than "custom code modified/forked from existing 3D tools")

The Object Oriented Parallel Simulation Engine (OOPSE) was introduced by A. et al. (n.d.) as a relatively light-weight molecular dynamics simulation package focused on "efficiently integrating equations of motion for atom types with orientational degrees of freedom" (from abstract). While OOPSE was further developed and renamed OpenMD, a fork of OOPSE was developed specifically to model water in two dimensions.

1.5.3 Literature Review on Relevant Works

TODO: Work with MB water,

CHAPTER 2

On Algorithms for Building and Sampling Disordered Crystal States

2.1 States and Properties of Ice

Ice is cool. Ice has many forms, each with unique environments and structures that give rise to similar and unique properties.

2.1.1 Bernal-Fowler Ice Rules

Bernal-Fowler Ice Rules are the basic rules for how water molecules interact in an ice structure. DETAILS ON BF PAPER

Basically, water’s tetrahedral structure allows for four interactions on each molecule. The two protons allow for a hydrogen bond with a lone pair from a neighboring oxygen atom. Similarly, the oxygen atom’s two lone pairs allow for a hydrogen bond with a neighboring proton. These rules are fairly rigid in the sense that every water molecule can interact with two oxygen atoms and two protons from four surrounding water molecules. These are also free-form in the sense that each of the four attached water molecules can occupy one of three rotational microstates, allowing for 81 possible configurations (including rotational duplicates).

2.1.2 Forms of Ice

While ubiquitous in the ‘ I_h ’ form, ice water has many known forms. As of the writing of this work, there are 17 established forms of ice. These forms usually occur in cubic, hexagonal, and orthorhombic crystal structures. The relationship between external pressure and temperature are the primary defining characteristics of which form will

form in a given system. Do other characteristics come into play???????

2.1.3 Ice I_h

As the most commonly found form on earth, ice I_h is a highly desired form for computational studies involving ice systems.

2.1.4 Efforts to Generate Ice I_h

Has anyone else published efforts to generate Ice I_h ? I'm sure someone has.

2.1.5 Comparison between Ice XI and Ice I_h

While ice I_h is known as the most common form of ice found on the planet, it is much more difficult to computationally generate than an ice XI crystal. The ease of generation of an ice XI structure stems from the repetition of a unit cell with consistent layering and orientation throughout the crystal lattice.

With ice I_h crystals, the proton-disordered form introduces entropy by way of rotational disorder. As the protons and lone pairs are no longer consistently ordered, hydrogen bonds may no longer form properly at all interaction sites. The interaction of proton with proton or lone pair with lone pair are not hydrogen bonds and are considered defects in the lattice. An ice structure of randomly oriented molecules without consideration of hydrogen bonds will likely produce defects at many interaction sites across the lattice and weaken the integrity of the system, leading to stability problems while running simulations. In generating the crystal, the cause of these defects must be considered and countered effectively.

2.2 Method Design

2.2.1 Overview

The big idea is to convert an easy-to-make ice XI crystal into an ice Ih crystal. Because the key difference in structure is the proton-orderedness, it might be possible to rearrange the water molecule orientations in a pseudorandom way to create an ice Ih crystal. This section walks through the method developed to convert ice XI into ice Ih, the results of initial testing, and imperfections discovered in the design.

2.2.2 Selection of Software Tools

Python was chosen as the language of the tool due to the versatility of the language and the ease of development due to the "pseudocode" written style of the language and the availability of scientific packages including SciPy and NumPy. Python version 2.7 was specifically chosen due to familiarity with the language. Crystal files were defined and saved as Protein Data Bank (.pdb) files as this format allows for defining multiple molecules within a larger structure with a simple X, Y, Z grid position format.

2.2.3 Generation of Source Ice XI

This is Dr. Fennell's method to create an ice XI pdb file. Basically, the ice XI unit cell of eight water molecules is repeated as desired to create a sufficiently large crystal. The primarily used crystal consists of a 3 x 3 x 6 unit cell repetition totaling 432 water molecules.

2.2.4 Source Ingestion

It is important that the crystal be read and stored in an efficient method to keep relevant information about each molecule easily accessible. As the file is read in, each

molecule is stored as an entry in a multidimensional array where the first index is the molecule number. Further, the second index defines the molecule number where 0 is oxygen and 1 and 2 are the protons. The third, fourth, and fifth indices define the X, Y, and Z position coordinates.

2.2.5 Identifying Neighboring Molecules

Identifying the neighboring molecules proved computationally difficult. The most effective method is to find the closest four molecules by computing a distance calculation between every two oxygen atoms. This ensures every molecule is considered, but also presents significant hurdles. First, a distance calculation utilizes an extremely computationally-inefficient square root calculation, which can be ignored by instead calculating the squared-distance between molecules and finding the lowest values.

Second, molecules on the walls and edges of the molecule will not have four neighbors in the non-periodic crystal. This is accounted for by shifting all six sides to make a pseudo-periodicity for these edge cases. Those periodically-neighboring molecules are flagged with a shifting value in the neighboring atom array by specifying a translation in the x, y, or z axis values. Unfortunately, the necessary code to implement the periodically-neighboring molecule detections requires a major rewrite of the entire tool and has not yet been implemented.

Once these closest neighboring oxygen atoms have been discovered, the appropriate interacting tetrahedral position is identified by finding the closest of the four tetrahedral positions using the same squared-distance calculation with the four defined tetrahedral positions detailed in the next subsection.

2.2.6 Defining Tetrahedral Positions

An important aspect of pseudorandom selection is the existence of a bank of options. Using the ingestion portion to calculate and store all tetrahedral possibilities proves

useful. For each water molecule, the first two tetrahedral positions are known by the positions of the two hydrogen atoms. The other two positions are found by rotating one hydrogen atom 120° twice about the vector from the oxygen atom through the other hydrogen atom and storing the resulting positions as tetrahedral positions three and four.

This does not produce an exactly correct tetrahedral position of potential hydrogen atoms due to the slight acuteness of the H-O-H bond created by the variance in repulsive forces between the two lone pairs of electrons and two hydrogen atoms. Fortunately, this difference is sufficiently small for visualization programs like Avogadro to still recognize hydrogen bonds between a rotated hydrogen atom and corresponding neighboring lone pair. Currently, the method relies on a very soft annealing process by a simulation package to minimize the effect of this hydrogen bond imperfection. Future versions of this method may account for the variations.

2.2.7 Pseudorandom Rearrangement of Water Molecules

Once the tetrahedral positions have been defined, each water molecule is ready to rotate. What may seem the most crucial step in this methods ends up being the most simple. As designed, the rotation of water molecules is as simple as using a stepwise iterator to pseudorandomly select two tetrahedral positions for the hydrogen bonds and store the new positions in a new crystal array. An extremely important note is that this rearrangement does not consider the orientations of neighboring molecules and likely introduces defects of hydrogen - hydrogen and lone pair - lone pair interactions. The likelihood of a defect-free interaction lattice forming is nearly zero and is assumed to have a great deal of defects within the lattice.

2.2.8 Detecting Hydrogen Bond Defects

After all water molecules have been rearranged, defects between incorrectly-interacting hydrogen bonds must be found and corrected. Discovering the defects relies on the detection of neighboring molecules and the appropriate interacting hydrogen atom or electron lone pair. As previously discussed, the initial data ingest records and detects the nearest water molecules and determines the tetrahedral position containing the interacting space, be it electron lone pair or hydrogen atom. From that data, the detection of a valid hydrogen bond is as simple as checking both all interacting tetrahedral positions and confirming that they both do not contain or lack a hydrogen atom. Additionally, each water molecule keeps a count of how many defects are present among the four positions. This allows for contextual changes during the correction step.

2.2.9 Correcting Hydrogen Bond Defects

Once the hydrogen bond defects have been discovered and marked, each needs to be corrected. The most direct approach to this is to sequentially walk through each defect and repeat the pseudorandom rotation until the number of defective regions is zero or a user-specified value. The current implementation sorts the defect list by the number of defects and attempts to fix the most defective molecules first. The most defective molecules may include defects impossible to solve by simple rotation, specifically when neighboring molecules have collectively directed three or four hydrogen atoms or electron lone pairs at the target water. These can only be solved by adjusting one or more of the neighboring molecules until the number of hydrogen atoms and electron lone pairs have balanced. Unfortunately, this high-defect problem can quickly escalate if the neighboring molecules contain the same problem of unbalanced hydrogen atoms and electron lone pairs. The current solution is to recursively check for and fix these impossible interactions first, but has not yet yielded a defect-free crystal in testing.

The current design of the method allows for the user to specify a threshold of defects as an average per molecule. For example, a threshold of 2.5 will allow a maximum of 3 defects on any given molecule and will continue to correct defects until the average number of defects per molecule is equal to or below 2.5. Because each of these defects will be counted twice, once for each molecule, the total number of defects in a crystal can be determined by multiplying the average defect value by the number of molecules and dividing by two. As of the current implementation, the method cannot reliably produce a crystal with a threshold below 2 as it will continue to search until the system runs out of available memory and crash without recording any new structure.

2.3 Results of Method

When supplied with an input ice XI crystal, an output structure with rotated water molecule orientations strictly consistent with ice Ih describes a success at the most basic level. An example before and after of the method is given in figures 2.1 and 2.2. The generated ice Ih has been rotated. When following the subsequent layers in the crystal, patterns emerge. Inconsistently, some rows of waters remain consistent. Some of these are a uniform rotation of both hydrogen atoms, while others are just one consistently placed hydrogen atom. Multiple trials yield internally unique results, yet all contain these strange consistencies. This may be due to some accidental pattern in the method's implementation.




Figure 2.1: "Before" image of Ice XI




Figure 2.2: "After" image of generated ice Ih

2.4 Comparison with Other Methods

2.4.1 benefits of own method over others

2.4.2 benefits of other methods over this

2.5 Comments on Limitations and Proposed Improvements

During the hydrogen bond defect correction step, a weakness in the design is that any clustering or regions of high defect density will not be noticed. This allows the existence of a highly-defective region within the larger structure that could potentially cause problems when the crystal is used in simulations. The prevalence and occurrence of these defects have not been studied, but seem a natural inevitability of statistics. A potential solution with partial development will score regions based on the number of defects as a weighted function expanding out from a central molecule for N connections. For example, consider a given molecule defined as level 1. The neighboring four molecules are defined as level 2, and continued onward excepting already-defined molecules out to an N^{th} level. The number of defects in each level can be counted and averaged. Then a depressive factor along the lines of $\frac{1}{level}$ can be used to diminish the value of defects further away from the first-level molecule. This would create a value for each molecule that shows the relative density of defects centered about that specific molecule and could even be plotted as a gradient change within the crystal. The equation would be something like:

$$Value = \sum_{l=1}^{N_{levels}} \left[\frac{1}{l} * \frac{1}{N_{molecules}} * \sum_{m=1}^{N_{molecules}} [N_{defects,m}] \right] \quad (2.1)$$

CHAPTER 3

Crystal and Liquid 2D Water at Interfaces

3.1 Two-Dimensional Water

EACH SUBSECTION: DEFINITION OF TERMS

Rose water is fairly new on the computational scene and so I may also include a review on the Mercedez-Benz water system as well as any other attempts to model water in two dimensions. For the rose potential system, I will review the Lennard-Jones potential as well as any other equations/systems related to the rose potential.

3.1.1 Lennard-Jones Potential

The Lennard-Jones Potential well is a soft-sphere model of interaction between two spheres described with

$$V_{LJ} = 4\epsilon[(\frac{\sigma}{r})^{12} - (\frac{\sigma}{r})^6] \quad (3.1)$$

where V is the potential, r is the distance between the center of two particles, σ is the specific distance between the two particles where the potential is zero, and $-\epsilon$ is the minimum potential of the plot. **REFINE WORDING:** The plot is defined in $[0, \infty)$. As two particles approach from infinity, their interaction become negative - which is an attractive force - and will approach the global minimum of $-\epsilon$. The r of this interaction is slightly larger than the combined radii of the two particles - which means they aren't quite touching - and is the equilibrium distance between the two particles. As r decreases beyond the minimum and toward σ , the interaction strength

increases and reaches zero as $r = \sigma$. At $r < \sigma$,

Potential digression: In a "hard-sphere" model, a particle's radius is firm, which is to say that the interaction potential is infinite at r less than σ . Basically a vertical line between two discrete values (usually ϵ and ∞) as the potential shifts from $r \geq \sigma$ to $r < \sigma$ (maybe include image?). The Lennard-Jones potential is a "soft-sphere" model, which blurs the line and replaces the vertical line with a functional representation. This breaks with reality as the particles become "squishy" and the potential ramps up toward infinity as r decreases. The benefit to the soft-sphere model is that modeling programs can more-easily account for overlaps in particles during time steps than with hard-sphere models. For example, a hard-sphere model of two particles interacting will likely not have a position where $r = \sigma$ and will potentially overlap. At this overlap, the potential is infinity and will introduce a nearly infinite force at that instant of time. Computer systems do not like having infinitely large repulsions suddenly introduced into a simulation.

3.1.2 Modeling Water in Two Dimensions

Modeling in two dimensions sacrifice the "realism" of models in three dimensions, but reduce the computational load significantly. This allows researchers (scientists, chemists, digital magicians?) to test more simple designs in two dimensions as well as a higher volume of simulations at the same time/computational cost.

3.1.2.1 Mercedes-Benz Model

The "Mercedes-Benz" BN2D model of water first proposed by Ben-Naim (1971) as "waterlike particles" are a popular two-dimensional representation of water. ROUGH: details of shape of MB water

The mathematical model used in the BN2D model is generated from the Percus-Yevick equation by substituting the approximation

$$c(X_1, X_2) = y(X_1, X_2)f(X_1, X_2) \quad (3.2)$$

into the Percus-Yevick equation obtained from the Ornstein-Zernike relation

$$h(X_1, X_2) = c(X_1, X_2) + \frac{\rho}{2\pi} \int c(X_1, X_3)h(X_3, X_2)dX_3 \quad (3.3)$$

to produce the overall relation

$$y(X_1, X_2) = 1 + \frac{\rho}{2\pi} \int y(X_1, X_3)f(X_1, X_3) \times \left[y(X_3, X_2)f(X_3, X_2) + y(X_3, X_2) - 1 \right] dX_3 \quad (3.4)$$

3.1.2.2 Rose Potential Model

The rose potential is another model first introduced by Williamson et al. (2017). This model, while similar to the three-pronged BN2D, is notably different in that the rose potential model simplifies the model by use of a radial sinusoidal plot to make the three "prongs" of the particle.

3.1.2.3 Two-Dimensional Modeling

Something other than OOPSE? (not seeing obvious answer other than "custom code modified/forked from existing 3D tools")

The Object Oriented Parallel Simulation Engine (OOPSE) was introduced by A. et al. (n.d.) as a relatively light-weight molecular dynamics simulation package focused on "efficiently integrating equations of motion for atom types with orientational degrees of freedom" (from abstract). While OOPSE was further developed and renamed OpenMD, a fork of OOPSE was developed specifically to model water in two dimensions.

3.1.3 Literature Review on Relevant Works

intro: Paragraph refreshing current work. Emphasize computational efficiency of Rose-Potential over Mercedez-Benz model.

3.2 Goal of Project

The objective of this work was to model two-dimensional water with a surface designed to discourage crystal growth at freezing temperatures. The design of the surface was the primary focus. Successfully designing a surface capable of discouraging water ice formation at freezing temperatures would provide valuable information in designing a three-dimensional model of the same type at a reduced computational cost.

Idea: adjust freezing point depression to be freezing point modification.

3.3 Tools and Terms

Either a refresh from intro or a detailed explanation of OOPSE and the reduced terms.

3.3.1 OOPSE in 2D

Detail differences in computation system that Dr. Fennell developed to allow 2D MD(?) on a 3D program.

3.3.2 Reduced Terms: 2D analogues

Detail differences in dimensionality and define the reduced dimensions. Still working on the understanding/equations.

3.4 Designing System

Include: ensemble, thermodynamic variables, box attributes (size, pressure, temp, etc), number of waters, surface (size, spacing between beads of surface, charges, LJ values, etc)

3.4.1 Defining the Surface

Explain how to develop the surface in the program and how to build a custom surface

3.4.1.1 Manipulation of LJ Potential

Manipulate σ and ϵ values to effectively adjust the radius and interaction strength of surface beads.

3.4.1.2 Manipulation of bead spacing

Detail design of optimizing bead spacing for freezing encouragement or disruption.

3.5 Results

Success of freezing point elevation, pending results for freezing point depression

3.5.1 Future Work

Continued efforts to disrupt crystal growth on the surface.

CHAPTER 4

Germanium Compounds and QM Concerns

4.1 Modeling Germanium Compounds

EACH SUBSECTION: DEFINITION OF TERMS

This will be an interesting section as there is extremely little in terms of Ge computational work. Perhaps a broader search will yield interesting results. For sake of thoroughness, I will also include work on computational energy optimization in general and work through complications brought by the size of Ge. I might also include a portion on the statistical spread of conformations at a given temperature (internal energy?) I may include a sentence or paragraph on Gaussian-based publications.

4.1.1 Tools for Modeling Germanium

Computational Requirements and reasons for those requirements.

Germanium is not the most-studied atom in computational works. The majority of Germanium studies are done with Gaussian (citation needed?).

4.1.2 Computational Complexity of Germanium Compounds

Draw-backs of modeling Germanium. Uncommon but still necessary for wetwork.

4.1.3 Literature Review on Relevant Works

Make note of various Germanium modeling research. Make note of tools and methods used.

Conformation	Energy (E_h)	Δ Energy (E_h)	Δ Energy ($\frac{KJ}{mol}$)
Trans-coplanar	-15014.8403143	0.0066255	17.39525025
Cis-Trans-Cis	-15014.7983311	0.0486087	127.6221418
Trans-Cis-Trans	-15014.8469398	0.0000000	0.0000000
Cis-Trans-Trans	-15014.8246918	0.0222480	58.412124

Table 4.1: Collaborator’s Hexagermane Energies by Conformation
(density functional theory, unknown basis set, energy in Hartrees and KJ/mol)

4.2 The Initial Problem: Germanium Study

During Fall 2017, Dr. Christopher Fennell was approached by Dr. Charles Weinert of OSU to continue a collaborative effort in sampling conformation energies of two germanium-based compounds of interest to Dr. Weinert’s work. Seen as an opportunity to train a new graduate student in conformational calculations, this project was delegated to me. The initial focus was to create the two compounds in a 3D modeling program, save a file of each, run a conformation optimization program on a supercomputer, and read the output to report the findings. As detailed below, this work led to impossibilities, curiosities, and inconsistencies that resulted in a general solution and a discovery of a flaw in a popular computational program.

4.2.1 Parameters of Work and Previous Collaborator’s Results

The two subject germanium-based compounds are very similar: a germanium backbone with terminal isopropyl groups and internal phenyl rings. One compound constituted a pentagermane chain while the other a hexagermane backbone. The molecular formula for both is $Pr_3^iGe(GePh_2)_nGePr_3^i$ where n equals 3 for the pentagermane or 4 for the hexagermane compounds, respectively. An example image of both compounds in their fully-trans configurations are provided in figures 4.1 and 4.2.

Dr. Weinert had worked previously with an additional collaborator who provided conformation data supplied in table 4.1. If I want to cite somebody, all I do is type in the citation for Bernal and Fowler (1933).

The approach of labeling the conformation shape of each compound, given the




Figure 4.1: Fully trans configuration of pentagermanium-based compound.




Figure 4.2: Fully trans configuration of hexagermanium-based compound.

many points of torsion, focuses on the backbone structure. As the raw data from the collaborator was not available, the general dihedral angles of cis and trans proved a vexing focus for initial efforts at conformer design. Using Newman projections like in figure REF! as a visual guide, each Ge-Ge bond was defined as cis or trans based on the relative angle produced by the two adjacent bonded Ge atoms to each subject Ge. Specifically, the bonds are marked cis if the most acute angle is 90° or fewer, and likewise trans if greater than 90° up to the maximum 180° . Effectively the cis and trans angles coincide with gauche and anti in organic structure nomenclature. Terminal germanium atoms are not considered as a part of the conformation state. This is partly due to the definition in labeling where the terminal germanium does not have an adjacent germanium for the measured relative angle, in addition to the assumed C_3 symmetry of the terminal Ge with three isopropyl groups reducing the relative effects of terminal germanium rotation. Effectively, only dihedrals formed by four consecutive Ge are given a cis or trans label.

4.2.2 Design and Approach to Solution

The initial approach involved an attempt at basic replication of the collaborative results. This design gradually became more complex as I

4.2.2.1 Design 1: Occam’s Smallest Razer

With each non-terminal Ge-Ge dihedral initially labeled cis or trans for 0° or 180° , about 3 unique pentagermane and 6 unique hexagermane structures were built visually on a 3D visualization program (Avogadro). These were rotated without consideration for the phenyl rings populating the non-terminal Ge atoms. Each molecule was subjected to an energy minimization in Gaussian 09 with the B3LYP hybrid function and STO-3G basis set as a single particle in a vacuum at otherwise default settings. See code in REF for a sample job command file.

Unsurprisingly, only the fully trans conformers successfully converged (a 22% success rate) into a stable form. These troubles were likely caused by the poor design of the initial conformers. With initial results, the conformer design was altered into a more systematic approach with some consideration for the phenyl rings.

4.2.2.2 Design 2: A Blunt Effort

In the second iteration of the conformer design process, a greater number of backbone conformers were generated. Instead of the simple 180° opposition between the cis and trans conformers, more intentional initial angles seen in Newman projections were selected. Specifically, the anti and both gauche angles were chosen for the natural local minima in a non-bulky molecule, with both gauche angles (60 and 300) labeled as cis and the anti angle (180) as trans. For initial conformer design, these backbone angles were limited to three positions: 60° , 180° , or 300° . For the hexagermane compound, these structures were sequentially labeled trans-trans-trans, trans-trans-cis, trans-cis-trans, et cetera until all major unique conformers were produced. For clarity, each conformer was identified by the dihedral angles (60-60-60, 60-60-180) in increasing order (Ge 1-2-3-4, Ge 2-3-4-5, Ge 3-4-5-6 dihedral). The phenyl rings on the non-terminal Ge atoms were left untouched from an initial steepest-descent minimization available from Avogadro ran in the fully trans conformer.

To prevent potentially strong interactions between adjacent phenyl rings, an additional steepest-descent minimization from Avogadro was initially ran with the conformer-defining Ge-Ge dihedral angles locked in place. Additionally, a visual inspection of the phenyl rings and manual adjustments were utilized on Avogadro to reduce the chance of a relatively high energy local minima conformer. The phenyl rings usually were settled in a form of pi stacking or some kind of perpendicular ring interaction, based on relative energy stability according to the immediate simple minimization available.

To further avoid backbone rotation restrictions, variations of the bulky molecules were also produced. These included versions where the phenyl rings were replaced by methyl groups and also where the isopropyl ends were additionally replaced by methyl groups. The intention in these designs were to observe the shift in relative energy between the sets of conformers to determine how significant of a role the phenyl rings and isopropyl groups played. These variations, along with the original form structures, were subject to the same calculations as in the first design: Gaussian 09, B3LYP hybrid functional, STO-3G basis set, no angle restrictions, single particle in a vacuum, otherwise default parameters. The results of these calculations are tabulated in table REFNAME

Immediately obvious in the table are the considerable number of nonconverged results. An unexpected bulkiness trend followed that a fully methylated variation of the structure was most likely to converge to a stable state, while the fully internal phenyl structures with methyl ends slightly reduced convergence and the original fully internal phenyl structures with isopropyl ends drastically reduced convergence. The common-sense expectation that the addition of the phenyl ends would reduce stability was not realized in these results. A deeper exploration into the change of stability is a promising avenue for future investigation, but was not further explored in this work. As highlighted in table REF, the lowest energy conformer for each structure form varied greatly, but never included the fully trans conformer and only once the collaborator-reported trans-cis-trans conformer as the most stable. Still, given the considerable amount of nonconverged conformers, a new design was necessary to further improve the scope of the lowest energy conformation search.

4.2.2.3 Design 3: Death by 1.59 Million Cuts

In the final version of the conformer generation effort, additional creation efforts were focused on the individual phenyl rings. The unfavorable interactions between the

phenyl rings were considerable hurdle in the previous designs and a potential explanation for the large number of nonconverged structures, including the possibility that the terminal isopropyl hexagermane structures contained particularly unfavorable interactions among the phenyl rings. This third design sought to remove the uncertainty in phenyl ring bulkiness by applying the same approach as the backbone generation: create unique conformers of every backbone torsion and phenyl ring, limiting each torsion to one of three rotational positions following the Newman projection style. Unfortunately, this task proved prohibitively large.

As an explanation for the insurmountability of the problem, consider the hexagermane structure. The germanium dihedrals represent three rotatable bonds each with three initial positions. To include the phenyl rings would require the inclusion of eight new rotatable bonds each with three initial positions. Additionally, considering each terminal germanium’s rotation while ignoring each isopropyl’s rotatable bonds adds two initial positions each with three initial positions. Together, this creates a structure with 13 rotatable bonds each with three initial positions. The number of conformers follows as $3^{13} = 1,594,323$ initial conformers. Now we must consider the computational aspect of this many conformers. At 10 conformers rotated and generated per second and 16 KB per conformer, the initial conformers would require 44.3 hours and generate 25.49 GB of data just in the initial structures. At an average of 72 minutes per computation and 73.7 MB produced at B3LYP hybrid functional and STO-3G basis set and access to all 255 regular nodes of Oklahoma State University’s Cowboy cluster running in parallel, the complete computation would generate 117.5 TB of data and require 312 days of continuous computation to determine a possible lowest energy conformer of this one molecule at a relatively low level basis set and theory. A request to utilize 100% of university supercomputer resources for nearly a year for the sake of determining the lowest energy conformer of one molecule would likely be rejected, so this task would likely require a time scale of years or even

decades to produce with shared access to university resources. While conventionally considered a small molecule, the scale of conformers and computational requirements pushes this problem into the realm of Levinthal’s paradox.

While this third design would have likely revealed the lowest energy conformer, or at least one considerably close to the exactly lowest energy conformer, the effort ultimately fails under its own weight. Even with efforts to truncate duplicate forms, the problem of scale remains. A reduction by 50% still requires a computation effort in the timescale of years or decades for the calculation of a single molecule. For an effective computational outlook, this system needs to be reduced by several orders of magnitude.

4.2.3 Scale Reduction Efforts

For a system with conformers on the millions scale and computations on the hour scale, a magnitude reduction in either aspect would improve the practicality of this design approach. For example, by simplifying the computational method from 72 minutes on average to 5 minutes on average, the overall computational requirement would be reduced by 92%, a full order of magnitude. Unfortunately, reducing the complexity of the method sacrifices the reliability of data. A potential solution here would be to create rounds of calculations at different complexities, where each sequential round restricts the pool of potential conformers. Ideally, the balance of the increasing computational complexity and the decreasing pool size would maintain a consistent computational requirement. For example, a new round using a higher functional theory and basis set at 5x computational requirement would ideally be paired with a reduction in conformer pool size by a factor of 5. This would produce a series of calculation sets with additive computational requirement instead of a magnitudinal expansion.

The natural next question lies within the reliability of basis sets and functional

theories. It naturally follows that a less-accurate method should not be relied on while better methods exist. However, considering the scale of the conformer pool, it follows that a less accurate method would still produce energy values with a roughly similar internal consistency. For example, a 180° - 0° - 180° angle form of the hexagermane compound with parallel phenyl rings as modeled in figure REF will have intense syn interactions between some phenyl rings and will likely not yield a desirable energy value at any level of calculation while a fully trans form with perfect pi stacking phenyl rings will likely have a lower energy value at all levels of calculation. Therefore, at lower levels of accuracy, the extremely high energy conformers can be pruned from the pool early and drastically reduce overall computational requirements. A generic effort at producing a method in this style is detailed in chapter 2, while the remainder of this chapter concludes the efforts of calculating these germanium compounds.

4.2.4 junk section

4.2.5 Attempts at Simplification

One potential avenue of simplifying the process is removing all non-backbone groups.

4.2.6 Discovery of Program Flaw

This is a junk sentence.

CHAPTER 5

Sampling Conformation Landscapes by Rotatable Bond Degrees of Freedom

5.1 Introduction to Topic

EACH SUBSECTION: DEFINITION OF TERMS

For ROUGHLY forty years, computational programs have allowed investigators to model chemical systems with high accuracy to determine their physical properties.

5.1.1 A Brief History on Conformation Landscapes

5.1.1.1 Levinthal's Paradox

Discuss history of Levinthal and his paradox. Provide the non-paradoxical solution.

Next: Levinthal golf courses by Ken Dill.

5.1.2 Computational Modeling

Introduce importance and impact. Bigly important.

5.1.2.1 History of Chemical Modeling

Here I will introduce molecular modeling. This will begin with a brief history of the development of the field. It will continue through to mention the styles and goals of molecular modeling. Upon reaching modern techniques, I will discuss the benefits and costs associated with the major types of calculations (QM, MD, MC, etc).

5.1.2.2 Hardware: Oklahoma State University's Cowboy Cluster

It would also be appropriate to mention the computational capabilities of OSU's Cowboy cluster.

5.1.2.3 Software

Software to mention:

Visualization Avogadro, UCSF Chimera

Computing GAMESS, Gaussian, NWChem, Octopus

This also includes brief pros and cons about the programs and the general purpose of use in case.

5.1.2.4 Programming Languages

Another hugely important portion of this will include a choice in the programming languages used (mostly python (Cython-compiled!), some Perl and Bash).

5.1.3 Literature Review on Relevant Works

intro: Reference introduction: inherent complexity of "objective" or "exhaustive" search, levinthal's paradox as it applies to non-biological systems, ELSE?

Run through a set of dihedral positions at a constant interval. Selection of lowest-energy optimization organized on dihedral values. Quick determination of importance of dihedral based on how heavily it impacts internal energy. Splitting "best" dihedral into smaller interval to repeat the process.

This method produces an interesting visual plot with varying resolution at different X values (if plotting energy vs. dihedral). Example given in figure 5.1.




Figure 5.1: example VR chart (exploding like molecule/cell/fiber/muscle)




Figure 5.2: Flow of method design for variable resolution conformation landscape search.

5.2 Design of System

System designed in Python for ease of development and compiled via Cython for computational efficiency. Utilizes Gaussian and UCSF Chimera, but can be redesigned for any computational programs that accomplish the desired tasks. Overview of system flow given in figure 5.2.

This design, with implementation being a current work in progress, but shouldTM work as a cascade toward the lowest energy conformer in each case.

5.2.1 Variation of Theory and Basis Set Usage by System Size and largest atom type

System will have inherent restrictions. Give an example of large system with simple atoms, small system with complex atoms. System estimates quantity and cost of calculations based on computational limits defined by user for various theory-basis set pairings. System optimizes calculations for the scale of run (is it the first broad-scope search, or a final near-exact search).

Good to have: data on how theory and basis set alter computational requirement. Available in literature? Create data from runs?

5.2.2 Computational Optimization by Varying Resolution

Extant work not optimized for a general search (negative claim: make sure literature has nothing). Design should work with additional development (primary focus this semester) as a general search tool.

5.2.3 Inherent Complications

Complications of size and atom type, impossible conformers, duplications, limited computational resources.

5.3 Results

Current success: finding accepted lowest energy conformer of a two-dihedral system by manually cranking each step. Self-running is still a work in progress.

5.3.1 Problems

Difficulty in defining an abstract system based on arbitrary hardware limitations. Propose a test-run to determine efficiency and resource availability.

5.3.2 Anticipated Approaches for Future Work

Putting system into a single cohesive program. Further optimizing Theory/Basis Set determination by computational efficiency as well as system size (determine an upper-limit of computation?)

References

A., M. M., F., V. C., Teng, L., J., F. C. and Daniel, G. J. (n.d.), ‘Oops: An object-oriented parallel simulation engine for molecular dynamics’, *Journal of Computational Chemistry* **26**(3), 252–271.

URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/jcc.20161>

Ben-Naim, A. (1971), ‘Statistical mechanics of ”waterlike” particles in two dimensions. i. physical model and application of the percus-yevick equation’, *The Journal of Chemical Physics* **54**(9), 3682–3695.

URL: <https://doi.org/10.1063/1.1675414>

Bernal, J. D. and Fowler, R. H. (1933), ‘A theory of water and ionic solution, with particular reference to hydrogen and hydroxyl ions’, *The Journal of Chemical Physics* **1**(8), 515–548.

URL: <https://doi.org/10.1063/1.1749327>

Pauling, L. (1935), ‘The structure and entropy of ice and of other crystals with some randomness of atomic arrangement’, *Journal of the American Chemical Society* **57**(12), 2680–2684.

URL: <https://doi.org/10.1021/ja01315a102>

Williamson, C. H., Hall, J. R. and Fennell, C. J. (2017), ‘Two-dimensional molecular simulations using rose potentials’, *Journal of Molecular Liquids* **228**(Complete), 11–18.

APPENDIX A

Ice Ih to Ice XI Conversion

Listed below is the source code utilized in the conversion of a PDB Ice Ih structure into an Ice XI structure. This code is functional in a Python 2.7 environment with NumPy and SciPy packages included.

A.1 Code: PDBDisorganize.py

```
1
2 #!/usr/bin/python
3
4 # Author = Gentry Smith
5 # Copyright 2016, all rights reserved
6
7 # this reads in a .PDB file , takes an argument for deformities per
   molecules , and randomly organizes the crystal
8 # structure into a disordered proton formation
9
10 # import sample: python PDBDisorganize.py arg1 arg2 arg3
11 # where:
12 # arg1 = source pdb file to be read (ex: acetone.pdb or acetone)
13 # arg2 = number of defects per molecule (in H2O, num of non-hydrogen-
   bonds. from 0 to 4)
14 # arg3 = desired output pdb file name
15
16 import sys
17 print sys.path
18 import string
19 import numpy as np
20 import math
21 import random
22
23 sys.setrecursionlimit(10000000) # maximum recursive depth. Set to
   (10,000,000) as under maximum
24
25
26 pdbIN = file(sys.argv[1])      # source PDB file
27 maxErr = int(sys.argv[2])      # max errors allowed
28 pdbOUT = str(sys.argv[3])      # output file name
29 finalData = [ [ [ 0 for i in range(3) ] for j in range(3) ] for k in
   range(300) ]
30
31 # looks at args validity
32 def checkArgs(arg1, arg2, arg3):
33     returnBool = False
```

```

34     if type(arg1) != file: # check arg1
35         print "Bad arg", arg1, " must be a file "
36         returnBool = True
37     if type(arg3) != str: # check arg3
38         print "Bad arg", arg3, ", must be a file name"
39         checkPDBSuffix(arg3)
40         print arg3
41         returnBool = True
42     if type(arg2) != int: # check arg2 type
43         print "Bad arg2: ", arg2, " is not an int."
44         returnBool = True
45     elif type(arg2) == int:
46         if arg2 < 0 or arg2 > 4: # check arg2 range
47             print "arg2 is not in a valid range 0 <= arg2 <= 4"
48             returnBool = True
49     return returnBool
50
51 def checkPDBSuffix(pdbFile):
52     if string.find(pdbFile, '.pdb', 0, len(pdbFile)) == -1:
53         print("did not find 'pdb' in ", pdbFile, ". Appending...")
54         pdbFile += '.pdb'
55
56
57
58 # reads in file ,
59 def readFile(fileName):
60     print "Reading file..."
61     # gets number of atoms
62     atoms = 0
63     for line in fileName:
64         data = line.split()
65         if len(data) > 0:
66             if data[0] != "CONNECT" and data[0] != "END":
67                 atoms += 1
68     # print "atoms: ", atoms
69     numMol = atoms / 3 # assumes 3-atom water molecule
70     dataTable = [ [ [ 0 for i in range(3) ] for j in range(3) ] for k in
71                     range(numMol) ]
72     fileName.seek(0)
73     iter0 = 0
74     iter1 = 0
75     pdbType = -1
76     for line in fileName:
77         data = line.split()
78         if pdbType == -1:
79             if data[0] == "ATOM":
80                 pdbType = 0
81             elif data[0] == "HETATM":
82                 pdbType = 1
83         # print "LineTuple=", data
84         if len(data) > 1 and ( data[0] == "ATOM" or data[0] == "HETATM"
85                                ):
86             if data[0] == "ATOM":
87                 newData = getDataATOM(data)

```

```

86         for i in range(3):
87             #data[molecule][atom][X/Y/Z]
88             dataTable[iter0][iter1 % 3][i] = newData[i]
89         elif data[0] == "HETATM":
90             dataTable[iter0][iter1 % 3] = getDataHETATM(data)
91         if iter1 == 2:
92             iter0 += 1
93             iter1 = 0
94         elif iter1 != 2:
95             iter1 += 1
96     # print "DataTable: ", dataTable
97     print "File read"
98     return dataTable, pdbType
99
100
101     # Split by index
102     # if having a problem with reading data, check .pdb to see if data
103     # has a space between each value
104 # reads XYZ coordinate data from ATOM-type pdb
105 def getDataATOM(strLine):
106     # print "Getting ATOM Data..."
107     dataLine = strLine[5:8]
108     # print "dataline: ", dataLine
109     i = 0
110     while i < 3:
111         # print "dataline[", i, "]: ", dataLine[i]
112         dataLine[i] = float(dataLine[i])
113         # print "dataline[", i, "]" type: ", type(dataLine[i])
114         i += 1
115     return dataLine
116
117
118 # reads XYZ coordinate data from HETATM-type pdb
119 def getDataHETATM(strLine):
120     # print "Getting HETATM Data..."
121     dataLine = strLine[5:8]
122     # print "dataline: ", dataLine
123     i = 0
124     while i < 3:
125         # print "dataline[", i, "]: ", dataLine[i]
126         dataLine[i] = float(dataLine[i])
127         # print "dataline[", i, "]" type: ", type(dataLine[i])
128         i += 1
129     return dataLine
130
131
132 # gets all four position vectors of hydrogen/lone pair as offset of
133 # oxygen molecule
134 def getOrientations( molecule ):
135     # 120 degrees = ( 2 * pi ) / 3 radians
136     theta = ( ( 2 * math.pi ) / 3 )
137     newMol = zeroOrientation(molecule)
138     returnInt1 = rotateMolecule(newMol[1], newMol[2], theta)

```

```

138     returnInt2 = rotateMolecule(newMol[1], newMol[2], (-1 * theta) )
139     return [returnInt1, returnInt2]
140
141
142 # randomly selects new orientation, returns two unique ints, from 0 to 3
    inclusively
143 def newRandOrientation( positions ):
144     # print "Changing orientation"
145     randVal1 = random.randint(0,3)
146     randVal2 = random.randint(0,3)
147     while randVal1 == randVal2:
148         randVal2 = random.randint(0,3)
149     newMol = [ [ 0, 0, 0 ],
150               positions[ randVal1 ],
151               positions[ randVal2 ] ]
152     return newMol
153
154 # selects new orientation from list. Reduces computational overhead in
    re-orientation option traversal
155 def newSetOrientation( positions, pos1, pos2 ):
156     newMol = [ [ 0, 0, 0 ],
157               positions[ pos1 ],
158               positions[ pos2 ] ]
159     return newMol
160
161
162 # sets molecule coordinates so that oxygen is the origin
163 def zeroOrientation(source):
164     # print "Zeroing Molecule..."
165
166     oxy = source[0]
167     hyd1 = source[1]
168     hyd2 = source[2]
169
170     # print "Oxygen pos: ", oxy
171     # print "Hydrogen 1: ", hyd1
172     # print "Hydrogen 2: ", hyd2
173
174     zeroedOrigin = [0, 0, 0]
175     zeroedHyd1 = [0, 0, 0]
176     zeroedHyd2 = [0, 0, 0]
177     for i in range(3):
178         zeroedHyd1[i] = hyd1[i] - oxy[i]
179         zeroedHyd2[i] = hyd2[i] - oxy[i]
180
181     # print "Zeroed Hydrogen 1: ", zeroedHyd1
182     # print "Zeroed Hydrogen 2: ", zeroedHyd2
183
184     # return new molecule position
185     newMol = [zeroedOrigin, zeroedHyd1, zeroedHyd2]
186     return newMol
187
188 # resets the zeroed molecule to the original oxygen position
189 def resetOrientation(oxygenPos, molecule):

```

```

190 # print "Resetting molecule..."
191 rO = oxygenPos
192 rH1 = [0,0,0]
193 rH2 = [0,0,0]
194 newMol = []
195 for i in range(3):
196     rH1[i] = molecule[1][i] + rO[i]
197     rH2[i] = molecule[2][i] + rO[i]
198     newMol = [rO, rH1, rH2]
199 # print "Rebuilt Molecule: ", newMol
200 return newMol
201
202 # rotates vector about axis for theta degrees
203 # Handler for rotationMatrix function below
204 def rotateMolecule(vector, axis, theta):
205     rotMatx = rotationMatrix(axis, theta)
206     return np.dot(rotMatx, vector)
207
208
209 # Creates Rotation matrix for a given axis and theta
210 # from stackoverflow user unutbu
211 # page: http://stackoverflow.com/questions/6802577/python-rotation-of-3d-vector
212 def rotationMatrix(axis, theta):
213     """
214
215     :type axis: list
216     :type theta: union
217     """
218     axis = np.asarray(axis)
219     theta = np.asarray(theta)
220     axis /= math.sqrt(np.dot(axis, axis))
221     a = math.cos(theta/2.0)
222     b, c, d = -axis*math.sin(theta/2.0)
223     aa, bb, cc, dd = (a * a), (b * b), (c * c), (d * d)
224     bc, ad, ac, ab, bd, cd = (b * c), (a * d), (a * c), (a * b), (b * d), (c * d)
225     return np.array([ [ (aa + bb - cc - dd), ( 2 * ( bc + ad ) ), ( 2 *
226         ( bd - ac ) ) ],
227         [ ( 2 * ( bc - ad ) ), (aa + cc - bb - dd), ( 2 *
228         ( cd + ab ) ) ],
229         [ ( 2 * ( bd + ac ) ), ( 2 * ( cd - ab ) ), (aa +
230         dd - bb - cc) ] ] )
231
232 # gets results from rotateAboutAxis plus two Hydrogens to get the
233 # tetrahedron positions
234 def getTetrahedronPositions(molecule):
235     positions = [ [ 0 for i in range(3) ] for j in range(4) ]
236     newMol = zeroOrientation(molecule) # zero molecule
237     positions[0] = newMol[1]
238     positions[1] = newMol[2]
239     newPos = getOrientations(molecule) # get final two positions
240     positions[2] = list(newPos[0])

```



```

238     positions[3] = list(newPos[1])
239     return positions                                # return all four positions
240
241
242 # checks distance of new positions from zero
243 def checkDist(posArray):
244     distance = [0 for i in range(len(posArray))]
245     for i in range(len(posArray)):
246         distance[i] = ( (posArray[i][0] * posArray[i][0]) +
247                         (posArray[i][1] * posArray[i][1]) +
248                         (posArray[i][2] * posArray[i][2]) )
249         # print "Distance", i, ": ", distance[i]
250     avg = 0
251     for i in range(len(posArray)):
252         avg += distance[i]
253     averageDistance = ( avg / len(posArray) )
254     # print "Average Distance: ", averageDistance
255     return averageDistance
256
257
258 # prints data given a 3D table of water molecules
259 def printData(data):
260     print "Data: "
261     strData = [" O", "H1", "H2"]
262     dimData = ["X", "Y", "Z"]
263     bigAvg = 0
264     numAtoms = 0
265     for mol in range(len(data)):
266         for atom in range(len(data[mol])):
267             printStr = str(mol) + ": " + strData[atom] + ": "
268             for dimension in range(3):
269                 printStr += dimData[atom] + ":" + "{:7.3f}".format(data[
270 mol][atom][dimension]) + "\t"
271             print printStr
272             bigAvg += checkDist(zeroOrientation(data[mol])[1:])
273             numAtoms += 1
274             print ""
275     print "total average distance: ", bigAvg / numAtoms
276
277 # checks validity of molecule
278 def isDefectiveCheck(err, neighborData, posData, index):
279     # find nearby molecules (avg oxygen distance???)
280     print "checking for defects at index", index, "..."
281     print "neighbor indices: ", neighborData[index]
282     returnBool = False
283     neighbors = 4
284     for i in range(4): # count real neighbors
285         if neighborData[index][1][i] == -1:
286             neighbors -= 1
287     if neighbors <= err: # de facto good if num(neighbors) <
288 maxErrAllowed
289     # print "Fewer neighbors than allowed errors. de facto Good
290 Orientation"

```

```

289     returnBool = True
290     elif neighbors > err: # enough neighbors to require check
291         # print "More neighbors than error threshold"
292         defectCount = 0
293         for neighbor in range(4): # check each neighbor
294             if neighborData[index][1][neighbor] != -1: # skip over non-
existant neighbors
295                 molA = posData[index]
296                 molB = posData[ neighborData[index][1][neighbor] ]
297                 oxyDist = getDistBetweenAtoms(molA[0], molB[0])
298
299                 if minHydrogenDistance(molA, molB) > oxyDist: # check
for facing lone pairs
300                     print "Double Lone Pair defect"
301                     defectCount += 1
302                     break
303                 else: # check for facing protons
304                     smallerHydrogenDistanceCount = 0
305                     isDefective = False
306                     for first in range(2):
307                         if not isDefective:
308                             for second in range(2):
309                                 newDist = getDistBetweenAtoms(molA[ first
+ 1], molB[second + 1])
310                                 if newDist < oxyDist:
311                                     smallerHydrogenDistanceCount += 1
312                                 if smallerHydrogenDistanceCount > 1:
313                                     print "Double Hydrogen defect"
314                                     defectCount += 1
315                                     isDefective = True
316                     # print "Defects found:", defectCount
317                     if defectCount > 4:
318                         print "IMPOSSIBLE AMOUNT OF DEFECTS DETECTED!!!!"
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
319                     if defectCount > err:
320                         # print "Found a bad molecule!"
321                         returnBool = False
322                     else:
323                         # print "Molecule is within parameters."
324                         returnBool = True
325
326     return returnBool
327
328
329 # randomly re-reorients molecule and neighbors, rechecks all
330 def rerunMolAndNeighbors(err, neighborData, posData, index):
331     # print "Re-reordering molecule at", index
332     # err - max errors allowed
333     # neighborData - int[4] of neighbor indices
334     # posData - array of all molecule position vectors
335     # index - location of focus molecule in posData
336     isGood = False
337     timeCount = 0
338     while not isGood:

```

```

339         # re-rotate molecule through all positions (iterated through all
orientations)
340         positions = getTetrahedronPositions(posData[index])
341         zeroedMol = newRandOrientation(positions)
342         # print "isGood CHECK", isGood
343         isGood, posData = iterThroughRotations(err, neighborData,
posData, index)
344         posData[index] = resetOrientation(posData[index][0], zeroedMol)
345         if timeCount >= 13: # { (1 - 1/6)^n < 0.05 } says n = 17
346             # BROKEN - need to rebuild
347             # 0. evaluated molecule has too many defects
348             # 1. reorient molecule statistically probable amount of
times to cover all orientations
349             # 2. Repeat 1. with neighbor 1
350             # 2a repeat 1. with original molecule
351             # 3. Repeat 2. with neighbor 2, 3, 4, as/if necessary
352             for neighborIndex in range(4):
353                 if neighborData[index][1][neighborIndex] != -1:
354                     positions = getTetrahedronPositions(posData[
neighborIndex])
355                     zeroedMol = newRandOrientation(positions)
356                     posData[neighborIndex] = resetOrientation(posData[
neighborIndex][0], zeroedMol)
357                     # isGood = isDefectiveCheck(err, neighborData,
posData, neighborIndex)
358                     isGood = isDefectiveCheck(err, neighborData, posData, index)
359                     if not isGood:
360                         isGood, posData = rerunMolAndNeighbors(err, neighborData
, posData, neighborData[index][1][neighborIndex])
361                     finalData = posData
362                     return True, finalData
363
364 # iterates molecule through all possible rotations
365 def iterThroughRotations(err, neighborData, posData, index):
366     isGood = False
367     pos1 = 0 # tetrahedral position for H1
368     pos2 = 0 # tetrahedral position for H2
369     while not isGood or (pos1 != 3 and pos2 != 3): # iterates through
all orientations, stops if good orientation
370         if pos1 != pos2:
371             posData[index] = newSetOrientation(posData[index][0], pos1,
pos2)
372             isGood = isDefectiveCheck(err, neighborData, posData, index)
373             if pos2 < 3:
374                 pos2 += 1
375             elif pos2 == 3:
376                 if pos1 < 3:
377                     pos1 += 1
378                 pos2 = 0
379             return isGood, posData
380 # determines minimum hydrogen distance between two atoms
381 def minHydrogenDistance(mol1, mol2):
382     minDist = 100
383     for first in range(2):

```

```

384         for second in range(2):
385             newDist = getDistBetweenAtoms(mol1[first+1], mol2[second+1])
386             if newDist < minDist:
387                 minDist = newDist
388         return minDist
389
390
391
392
393
394 # finds neighboring molecules of each molecule
395 def getNeighbors(data):
396     returnData = [ [ [ 0 for i in range(4) ] for j in range(2) ] for k
397                     in range(len(data)) ] # data[molecule][distance,index][four values]
398     for mol1 in range(len(data)):
399         minDist = [100, 100, 100, 100]
400         minIndex = [0, 0, 0, 0]
401         for mol2 in range(len(data)):
402             if mol1 != mol2:
403                 newMin = getDistBetweenAtoms(data[mol1][0], data[mol2
404                                                 ][0])
405
406                 bigIndex = indexOfBiggest(minDist)
407                 if newMin < minDist[bigIndex]:
408                     minDist[bigIndex] = newMin
409                     minIndex[bigIndex] = mol2
410
411                 for i in range(4):
412                     if minDist[i] >= 9:
413                         minDist[i] = -1
414                         minIndex[i] = -1
415
416                 # print "Four smallest Distances of", mol1, ": ", minDist
417                 # print "Four smallest Indices of", mol1, ": ", minIndex
418                 returnData[mol1] = [minDist, minIndex]
419         return returnData
420
421
422
423
424
425 # finds distance between oxygen atoms
426 def getDistBetweenAtoms( mol1, mol2 ):
427     distance = ( ( ( mol1[0] - mol2[0] ) * ( mol1[0] - mol2[0] ) ) +
428                 ( ( mol1[1] - mol2[1] ) * ( mol1[1] - mol2[1] ) ) +
429                 ( ( mol1[2] - mol2[2] ) * ( mol1[2] - mol2[2] ) ) )
430     return distance
431
432
433
434 # gets index of largest item from a list
435 def indexOfBiggest(check):
436     bigIndex = 0
437     for i in range(len(check)):
438         if check[i] > check[bigIndex]:
439             bigIndex = i
440     return bigIndex
441
442
443
444 # writes data to PDB file
445 def writeDataPDB(data, pdbType):

```

```

436     print "Writing Data to", str(pdbOUT)
437     fileName = str(pdbOUT)
438     output = open(fileName, 'w')
439     if pdbType == 0:
440         writeDataPDBATOM(data, output)
441     elif pdbType == 1:
442         writeDataPDBHETATM(data, output)
443     output.close()
444
445
446 # Writes data to PDB file style = ATOM
447 def writeDataPDBATOM(data, inFile):
448     iterator = 0
449     for molecule in range(len(data)):
450         for atom in range(3):
451             iterator += 1
452             outStr = "ATOM "
453             outStr += str(iterator)
454             while len(outStr) < 11:
455                 outStr = outStr[:6] + " " + outStr[6:]
456             outStr += " "
457             if atom == 0:
458                 outStr += " O " + " WAT"
459             elif atom == 1:
460                 outStr += " H1 " + " WAT"
461             elif atom == 2:
462                 outStr += " H2 " + " WAT"
463             outStr += str(molecule)
464             while len(outStr) < 26:
465                 outStr = outStr[:20] + " " + outStr[20:]
466             outStr += " "
467             outStr += "{:8.3f}".format(data[molecule][atom][0])
468             outStr += "{:8.3f}".format(data[molecule][atom][1])
469             outStr += "{:8.3f}".format(data[molecule][atom][2])
470             outStr += " 1.00" + " 0.00"
471             outStr += " "
472             if atom == 0:
473                 outStr += " O "
474             elif atom == 1:
475                 outStr += " H "
476             elif atom == 2:
477                 outStr += " H "
478             outStr += "\n"
479             inFile.write(outStr)
480
481
482 # Writes data to PDB file style = HETATOM
483 def writeDataPDBHETATM(data, inFile):
484     iterator = 0
485     for molecule in range(len(data)):
486         for atom in range(3):
487             iterator += 1
488             outStr = "HETATM"
489             outStr += str(iterator)

```

```

490         while len(outStr) < 11:
491             outStr = outStr[:6] + " " + outStr[6:]
492         outStr += " "
493         if atom == 0:
494             outStr += " O " + " WAT"
495         elif atom == 1:
496             outStr += " H1 " + " WAT"
497         elif atom == 2:
498             outStr += " H2 " + " WAT"
499         outStr += str(molecule)
500         while len(outStr) < 26:
501             outStr = outStr[:20] + " " + outStr[20:]
502         outStr += " "
503         outStr += "{:8.3f}".format(data[molecule][atom][0])
504         outStr += "{:8.3f}".format(data[molecule][atom][1])
505         outStr += "{:8.3f}".format(data[molecule][atom][2])
506         outStr += " 1.00" + " 0.00"
507         outStr += " "
508         if atom == 0:
509             outStr += " O "
510         elif atom == 1:
511             outStr += " H "
512         elif atom == 2:
513             outStr += " H "
514         outStr += "\n"
515         inFile.write(outStr)
516
517
518 # runs program
519 def testRun(inFile, err, outFile):
520     print "Running Test Version of Program..."
521
522
523 # this is the parent runner for the program
524 def runPgm(inFile, err):
525     print "Running Program..."
526     data, pdbType = readFile(inFile)
527     newData = [ [ [ 0 for i in range(3) ] for j in range(3) ] for k in
528                 range(len(data)) ]
529     print "Reordering Molecules..."
530     for i in range(len(data)):
531         positions = getTetrahedronPositions(data[i])
532         zeroedMol = newRandOrientation(positions)
533         newMol = resetOrientation( data[i][0], zeroedMol )
534         newData[i] = newMol
535     print "Molecules Reordered"
536     connectedMolecules = getNeighbors(newData) # -1 index = not
537     neighboring
538     finalData = newData
539     for i in range(len(connectedMolecules)):
540         # print "check defects"
541         isFine = isDefectiveCheck(err, connectedMolecules, finalData, i)
542         # print "isFINE CHECK", isFine
543         if not isFine:

```

```

542         # print "fixing defects"
543         while not isFine:
544             # print "RerunMol"
545             isFine, finalData = rerunMolAndNeighbors(err,
connectedMolecules, finalData, i)
546             # print "rerunDone"
547             writeDataPDB(finalData, pdbType)
548             # printData(newData)
549
550
551 badArgs = checkArgs(pdbIN, maxErr, pdbOUT) # stop in case of bad
argument
552
553 # check input args
554 if not badArgs: # stop in case of bad argument
555     print "Good Arguments, Initializing Reorientation with", maxErr, "
maximum defects"
556     # testRun(pdbIN, maxErr, pdbOUT)
557     runPgm(pdbIN, maxErr)
558 elif badArgs:
559     print "Bad Arguments, Quitting..."

```

APPENDIX B

Conformation Landscapes

APPENDIX C

Germanium Landscape

Listed below are two example Germanium PDB files. The first is for the end-goal hexagermane in the trans-trans-trans conformation with isopropyl groups on the terminal Ge atoms. The second is for the simplified butagermane with fully protonated Germanium atoms.

C.1 Code: hexagermane-transall.pdb

```
1 HEADER
2 REMARK Title: hexagermane-transall system
3 HETATM 1 Ge 1 -4.399 0.008 0.355 0.00 0.00
   Ge
4 HETATM 2 Ge 1 -1.965 0.138 -0.022 0.00 0.00
   Ge
5 HETATM 3 C 1 -4.822 1.886 0.961 0.00 0.00
   C
6 HETATM 4 C 1 -5.008 -1.297 1.715 0.00 0.00
   C
7 HETATM 5 C 1 -5.256 -0.261 -1.445 0.00 0.00
   C
8 HETATM 6 C 1 -1.213 1.435 1.157 0.00 0.00
   C
9 HETATM 7 Ge 1 -0.756 -1.988 0.223 0.00 0.00
   Ge
10 HETATM 8 C 1 -1.297 -2.917 1.805 0.00 0.00
   C
11 HETATM 9 Ge 1 1.647 -1.496 0.371 0.00 0.00
   Ge
12 HETATM 10 C 1 -1.182 -3.010 -1.339 0.00 0.00
   C
13 HETATM 11 C 1 2.131 -0.425 1.877 0.00 0.00
   C
14 HETATM 12 C 1 2.111 -0.634 -1.269 0.00 0.00
   C
15 HETATM 13 Ge 1 2.889 -3.585 0.738 0.00 0.00
   Ge
16 HETATM 14 C 1 2.287 -4.358 2.378 0.00 0.00
   C
17 HETATM 15 Ge 1 5.327 -3.386 1.080 0.00 0.00
   Ge
18 HETATM 16 C 1 2.766 -4.685 -0.813 0.00 0.00
   C
19 HETATM 17 C 1 5.688 -2.615 2.887 0.00 0.00
   C
```

20	HETATM	18	C	1	6.239	-2.415	-0.417	0.00	0.00
		C							
21	HETATM	19	C	1	5.893	-5.324	0.888	0.00	0.00
		C							
22	HETATM	20	C	1	-3.527	2.543	1.328	0.00	0.00
		C							
23	HETATM	21	C	1	-5.754	1.844	2.133	0.00	0.00
		C							
24	HETATM	22	H	1	-5.303	2.355	0.072	0.00	0.00
		H							
25	HETATM	23	H	1	-5.269	1.358	2.999	0.00	0.00
		H							
26	HETATM	24	H	1	-6.679	1.287	1.913	0.00	0.00
		H							
27	HETATM	25	H	1	-6.047	2.856	2.449	0.00	0.00
		H							
28	HETATM	26	H	1	-3.043	2.019	2.171	0.00	0.00
		H							
29	HETATM	27	H	1	-3.683	3.585	1.642	0.00	0.00
		H							
30	HETATM	28	H	1	-2.818	2.559	0.490	0.00	0.00
		H							
31	HETATM	29	H	1	-4.336	-1.167	2.589	0.00	0.00
		H							
32	HETATM	30	C	1	-4.907	-2.680	1.154	0.00	0.00
		C							
33	HETATM	31	C	1	-6.417	-0.909	2.051	0.00	0.00
		C							
34	HETATM	32	H	1	-3.858	-2.985	0.982	0.00	0.00
		H							
35	HETATM	33	H	1	-5.433	-2.773	0.187	0.00	0.00
		H							
36	HETATM	34	H	1	-5.349	-3.420	1.836	0.00	0.00
		H							
37	HETATM	35	H	1	-6.488	0.167	2.302	0.00	0.00
		H							
38	HETATM	36	H	1	-6.802	-1.477	2.909	0.00	0.00
		H							
39	HETATM	37	H	1	-7.103	-1.094	1.205	0.00	0.00
		H							
40	HETATM	38	C	1	5.200	-6.127	1.944	0.00	0.00
		C							
41	HETATM	39	C	1	7.384	-5.453	0.966	0.00	0.00
		C							
42	HETATM	40	H	1	5.523	-5.590	-0.126	0.00	0.00
		H							
43	HETATM	41	H	1	7.790	-4.974	1.874	0.00	0.00
		H							
44	HETATM	42	H	1	7.885	-4.994	0.099	0.00	0.00
		H							
45	HETATM	43	H	1	7.691	-6.509	0.992	0.00	0.00
		H							
46	HETATM	44	H	1	5.502	-5.821	2.960	0.00	0.00
		H							

47	HETATM	45	H	1	5.436	-7.197	1.849	0.00	0.00
		H							
48	HETATM	46	H	1	4.106	-6.027	1.879	0.00	0.00
		H							
49	HETATM	47	C	1	6.243	-1.232	2.746	0.00	0.00
		C							
50	HETATM	48	C	1	6.612	-3.524	3.636	0.00	0.00
		C							
51	HETATM	49	H	1	4.684	-2.582	3.376	0.00	0.00
		H							
52	HETATM	50	H	1	7.535	-3.731	3.068	0.00	0.00
		H							
53	HETATM	51	H	1	6.139	-4.497	3.853	0.00	0.00
		H							
54	HETATM	52	H	1	6.913	-3.088	4.599	0.00	0.00
		H							
55	HETATM	53	H	1	7.243	-1.234	2.279	0.00	0.00
		H							
56	HETATM	54	H	1	6.347	-0.742	3.725	0.00	0.00
		H							
57	HETATM	55	H	1	5.589	-0.589	2.128	0.00	0.00
		H							
58	HETATM	56	C	1	5.630	-1.055	-0.555	0.00	0.00
		C							
59	HETATM	57	H	1	6.024	-3.039	-1.315	0.00	0.00
		H							
60	HETATM	58	C	1	7.712	-2.342	-0.145	0.00	0.00
		C							
61	HETATM	59	H	1	7.923	-1.890	0.839	0.00	0.00
		H							
62	HETATM	60	H	1	8.227	-1.728	-0.898	0.00	0.00
		H							
63	HETATM	61	H	1	8.188	-3.335	-0.163	0.00	0.00
		H							
64	HETATM	62	H	1	4.573	-1.106	-0.861	0.00	0.00
		H							
65	HETATM	63	H	1	6.155	-0.455	-1.314	0.00	0.00
		H							
66	HETATM	64	H	1	5.675	-0.486	0.391	0.00	0.00
		H							
67	HETATM	65	H	1	-5.890	-1.163	-1.302	0.00	0.00
		H							
68	HETATM	66	C	1	-4.220	-0.487	-2.505	0.00	0.00
		C							
69	HETATM	67	C	1	-6.093	0.945	-1.729	0.00	0.00
		C							
70	HETATM	68	H	1	-6.841	1.122	-0.939	0.00	0.00
		H							
71	HETATM	69	H	1	-6.644	0.838	-2.676	0.00	0.00
		H							
72	HETATM	70	H	1	-5.478	1.858	-1.818	0.00	0.00
		H							
73	HETATM	71	H	1	-3.754	-1.481	-2.414	0.00	0.00
		H							

74	HETATM	72	H	1	-3.411	0.262	-2.459	0.00	0.00
		H							
75	HETATM	73	H	1	-4.659	-0.429	-3.512	0.00	0.00
		H							
76	HETATM	74	C	1	-1.706	1.681	2.429	0.00	0.00
		C							
77	HETATM	75	C	1	-0.128	2.155	0.679	0.00	0.00
		C							
78	HETATM	76	H	1	0.268	1.941	-0.323	0.00	0.00
		H							
79	HETATM	77	C	1	0.451	3.147	1.465	0.00	0.00
		C							
80	HETATM	78	C	1	-1.134	2.678	3.216	0.00	0.00
		C							
81	HETATM	79	C	1	-0.058	3.415	2.731	0.00	0.00
		C							
82	HETATM	80	H	1	-1.525	2.873	4.219	0.00	0.00
		H							
83	HETATM	81	H	1	1.306	3.716	1.086	0.00	0.00
		H							
84	HETATM	82	H	1	0.391	4.199	3.349	0.00	0.00
		H							
85	HETATM	83	C	1	-1.557	-4.274	1.694	0.00	0.00
		C							
86	HETATM	84	C	1	-0.365	-3.088	-2.455	0.00	0.00
		C							
87	HETATM	85	C	1	-2.027	-4.985	2.796	0.00	0.00
		C							
88	HETATM	86	H	1	-1.378	-4.792	0.742	0.00	0.00
		H							
89	HETATM	87	C	1	-1.446	-2.277	3.025	0.00	0.00
		C							
90	HETATM	88	C	1	-0.752	-3.866	-3.544	0.00	0.00
		C							
91	HETATM	89	H	1	0.592	-2.548	-2.482	0.00	0.00
		H							
92	HETATM	90	C	1	-2.386	-3.699	-1.304	0.00	0.00
		C							
93	HETATM	91	C	1	-2.219	-4.336	4.011	0.00	0.00
		C							
94	HETATM	92	H	1	-2.237	-6.056	2.707	0.00	0.00
		H							
95	HETATM	93	C	1	-1.915	-2.983	4.130	0.00	0.00
		C							
96	HETATM	94	H	1	-1.159	-1.217	3.132	0.00	0.00
		H							
97	HETATM	95	C	1	-2.771	-4.484	-2.388	0.00	0.00
		C							
98	HETATM	96	H	1	-3.043	-3.610	-0.422	0.00	0.00
		H							
99	HETATM	97	C	1	-1.952	-4.568	-3.509	0.00	0.00
		C							
100	HETATM	98	H	1	-0.105	-3.928	-4.425	0.00	0.00
		H							

101	HETATM	99	H	1	-3.721	-5.027	-2.358	0.00	0.00
		H							
102	HETATM	100	H	1	-2.253	-5.182	-4.364	0.00	0.00
		H							
103	HETATM	101	H	1	-2.596	-4.891	4.876	0.00	0.00
		H							
104	HETATM	102	H	1	-2.041	-2.474	5.091	0.00	0.00
		H							
105	HETATM	103	C	1	2.487	-3.679	3.571	0.00	0.00
		C							
106	HETATM	104	C	1	1.701	-5.563	-0.935	0.00	0.00
		C							
107	HETATM	105	C	1	3.733	-4.618	-1.807	0.00	0.00
		C							
108	HETATM	106	H	1	0.940	-5.615	-0.140	0.00	0.00
		H							
109	HETATM	107	C	1	1.598	-6.382	-2.057	0.00	0.00
		C							
110	HETATM	108	C	1	1.690	-5.609	2.382	0.00	0.00
		C							
111	HETATM	109	C	1	2.102	-4.259	4.776	0.00	0.00
		C							
112	HETATM	110	H	1	2.956	-2.680	3.567	0.00	0.00
		H							
113	HETATM	111	C	1	1.520	-5.523	4.784	0.00	0.00
		C							
114	HETATM	112	H	1	2.260	-3.721	5.716	0.00	0.00
		H							
115	HETATM	113	C	1	1.311	-6.197	3.585	0.00	0.00
		C							
116	HETATM	114	H	1	1.504	-6.131	1.431	0.00	0.00
		H							
117	HETATM	115	C	1	2.562	-6.313	-3.057	0.00	0.00
		C							
118	HETATM	116	H	1	0.754	-7.074	-2.153	0.00	0.00
		H							
119	HETATM	117	C	1	3.630	-5.430	-2.933	0.00	0.00
		C							
120	HETATM	118	H	1	4.590	-3.931	-1.700	0.00	0.00
		H							
121	HETATM	119	H	1	4.391	-5.376	-3.718	0.00	0.00
		H							
122	HETATM	120	H	1	2.481	-6.954	-3.941	0.00	0.00
		H							
123	HETATM	121	H	1	1.223	-5.984	5.731	0.00	0.00
		H							
124	HETATM	122	H	1	0.844	-7.187	3.587	0.00	0.00
		H							
125	HETATM	123	C	1	1.878	0.732	-1.306	0.00	0.00
		C							
126	HETATM	124	C	1	1.530	-0.534	3.120	0.00	0.00
		C							
127	HETATM	125	C	1	2.642	-1.289	-2.370	0.00	0.00
		C							

128	HETATM	126	C	1	2.179	1.455	-2.458	0.00	0.00
		C							
129	HETATM	127	H	1	1.444	1.239	-0.432	0.00	0.00
		H							
130	HETATM	128	C	1	3.179	0.461	1.679	0.00	0.00
		C							
131	HETATM	129	C	1	2.005	0.227	4.186	0.00	0.00
		C							
132	HETATM	130	H	1	0.661	-1.197	3.265	0.00	0.00
		H							
133	HETATM	131	C	1	2.940	-0.568	-3.524	0.00	0.00
		C							
134	HETATM	132	H	1	2.840	-2.370	-2.334	0.00	0.00
		H							
135	HETATM	133	C	1	2.710	0.804	-3.567	0.00	0.00
		C							
136	HETATM	134	H	1	1.989	2.533	-2.491	0.00	0.00
		H							
137	HETATM	135	H	1	3.358	-1.081	-4.396	0.00	0.00
		H							
138	HETATM	136	H	1	2.944	1.370	-4.475	0.00	0.00
		H							
139	HETATM	137	C	1	3.067	1.105	3.998	0.00	0.00
		C							
140	HETATM	138	H	1	1.534	0.140	5.170	0.00	0.00
		H							
141	HETATM	139	C	1	3.650	1.229	2.740	0.00	0.00
		C							
142	HETATM	140	H	1	3.633	0.553	0.682	0.00	0.00
		H							
143	HETATM	141	H	1	4.480	1.926	2.585	0.00	0.00
		H							
144	HETATM	142	H	1	3.439	1.703	4.836	0.00	0.00
		H							
145	HETATM	143	C	1	-2.039	0.838	-1.804	0.00	0.00
		C							
146	HETATM	144	C	1	-1.525	0.195	-2.916	0.00	0.00
		C							
147	HETATM	145	C	1	-2.655	2.077	-1.927	0.00	0.00
		C							
148	HETATM	146	C	1	-1.618	0.802	-4.168	0.00	0.00
		C							
149	HETATM	147	H	1	-1.048	-0.789	-2.818	0.00	0.00
		H							
150	HETATM	148	C	1	-2.746	2.686	-3.175	0.00	0.00
		C							
151	HETATM	149	H	1	-3.084	2.566	-1.036	0.00	0.00
		H							
152	HETATM	150	C	1	-2.223	2.047	-4.296	0.00	0.00
		C							
153	HETATM	151	H	1	-1.210	0.296	-5.049	0.00	0.00
		H							
154	HETATM	152	H	1	-3.229	3.663	-3.275	0.00	0.00
		H							

155	HETATM	153	H		1		-2.292	2.524	-5.279	0.00	0.00
		H									
156	HETATM	154	H		1		-2.539	1.081	2.827	0.00	0.00
		H									
157	CONECT	3	1	20	21	22					
158	CONECT	4	1	29	30	31					
159	CONECT	5	1	65	66	67					
160	CONECT	6	74	75	2						
161	CONECT	8	83	87	7						
162	CONECT	10	84	90	7						
163	CONECT	11	124	128	9						
164	CONECT	12	123	125	9						
165	CONECT	14	103	108	13						
166	CONECT	16	104	105	13						
167	CONECT	17	15	47	48	49					
168	CONECT	18	56	57	58	15					
169	CONECT	19	15	38	39	40					
170	CONECT	20	3	26	27	28					
171	CONECT	21	3	23	24	25					
172	CONECT	30	4	34	32	33					
173	CONECT	31	4	35	36	37					
174	CONECT	38	19	44	45	46					
175	CONECT	39	19	41	42	43					
176	CONECT	47	54	55	17	53					
177	CONECT	48	17	50	51	52					
178	CONECT	56	62	63	64	18					
179	CONECT	58	59	60	61	18					
180	CONECT	66	71	72	73	5					
181	CONECT	67	68	69	70	5					
182	CONECT	74	78	154	6						
183	CONECT	75	76	77	6						
184	CONECT	77	75	79	81						
185	CONECT	78	74	79	80						
186	CONECT	79	77	78	82						
187	CONECT	83	85	86	8						
188	CONECT	84	88	89	10						
189	CONECT	85	83	91	92						
190	CONECT	87	93	94	8						
191	CONECT	88	84	97	98						
192	CONECT	90	95	96	10						
193	CONECT	91	85	93	101						
194	CONECT	93	87	91	102						
195	CONECT	95	90	97	99						
196	CONECT	97	88	95	100						
197	CONECT	103	109	110	14						
198	CONECT	104	106	107	16						
199	CONECT	105	117	118	16						
200	CONECT	107	104	115	116						
201	CONECT	108	113	114	14						
202	CONECT	109	103	111	112						
203	CONECT	111	109	113	121						
204	CONECT	113	108	111	122						
205	CONECT	115	107	117	120						
206	CONECT	117	105	115	119						

207	CONECT	123	126	127	12
208	CONECT	124	129	130	11
209	CONECT	125	131	132	12
210	CONECT	126	123	133	134
211	CONECT	128	139	140	11
212	CONECT	129	124	137	138
213	CONECT	131	125	133	135
214	CONECT	133	126	131	136
215	CONECT	137	129	139	142
216	CONECT	139	128	137	141
217	CONECT	143	144	145	2
218	CONECT	144	143	146	147
219	CONECT	145	143	148	149
220	CONECT	146	144	150	151
221	CONECT	148	145	150	152
222	CONECT	150	146	148	153
223	CONECT	1	2	3	4
224	CONECT	2	1	143	6
225	CONECT	7	2	8	9
226	CONECT	9	7	11	12
227	CONECT	13	9	14	15
228	CONECT	15	13	17	18
229	CONECT	22	3		
230	CONECT	23	21		
231	CONECT	24	21		
232	CONECT	25	21		
233	CONECT	26	20		
234	CONECT	27	20		
235	CONECT	28	20		
236	CONECT	29	4		
237	CONECT	32	30		
238	CONECT	33	30		
239	CONECT	34	30		
240	CONECT	35	31		
241	CONECT	36	31		
242	CONECT	37	31		
243	CONECT	40	19		
244	CONECT	41	39		
245	CONECT	42	39		
246	CONECT	43	39		
247	CONECT	44	38		
248	CONECT	45	38		
249	CONECT	46	38		
250	CONECT	49	17		
251	CONECT	50	48		
252	CONECT	51	48		
253	CONECT	52	48		
254	CONECT	53	47		
255	CONECT	54	47		
256	CONECT	55	47		
257	CONECT	57	18		
258	CONECT	59	58		
259	CONECT	60	58		
260	CONECT	61	58		


```

261 CONECT    62    56
262 CONECT    63    56
263 CONECT    64    56
264 CONECT    65     5
265 CONECT    68    67
266 CONECT    69    67
267 CONECT    70    67
268 CONECT    71    66
269 CONECT    72    66
270 CONECT    73    66
271 CONECT    76    75
272 CONECT    80    78
273 CONECT    81    77
274 CONECT    82    79
275 CONECT    86    83
276 CONECT    89    84
277 CONECT    92    85
278 CONECT    94    87
279 CONECT    96    90
280 CONECT    98    88
281 CONECT    99    95
282 CONECT   100    97
283 CONECT   101    91
284 CONECT   102    93
285 CONECT   106   104
286 CONECT   110   103
287 CONECT   112   109
288 CONECT   114   108
289 CONECT   116   107
290 CONECT   118   105
291 CONECT   119   117
292 CONECT   120   115
293 CONECT   121   111
294 CONECT   122   113
295 CONECT   127   123
296 CONECT   130   124
297 CONECT   132   125
298 CONECT   134   126
299 CONECT   135   131
300 CONECT   136   133
301 CONECT   138   129
302 CONECT   140   128
303 CONECT   141   139
304 CONECT   142   137
305 CONECT   147   144
306 CONECT   149   145
307 CONECT   151   146
308 CONECT   152   148
309 CONECT   153   150
310 CONECT   154    74
311 END

```

The above molecule contains 154 atoms and 153 bonds, making it extremely computationally expensive for regular QM calculations. This made utilizing the large

molecule as a trial system unreasonable due to the prohibitively long computation time for each conformation, assuming the conformation calculation would complete at all.

The below PDB file is the simplified butagermane with fully protonated Germanium atoms. As a significantly smaller system with only 14 atoms and 13 bonds, the relatively short computation time allowed the trial system to move with relative ease.

C.2 Code: ge4h.pdb

```

1 COMPND      UNNAMED
2 AUTHOR      GENERATED BY OPEN BABEL 2.3.90
3 HETATM      1 GE      UNL      1      -3.520      1.842      -0.078      1.00      0.00
               Ge3—
4 HETATM      2 GE      UNL      1      -1.368      2.888      -0.034      1.00      0.00
               Ge2—
5 HETATM      3 GE      UNL      1      0.324      1.200      0.059      1.00      0.00
               Ge3—
6 HETATM      4 GE      UNL      1      2.475      2.248      0.099      1.00      0.00
               Ge
7 HETATM      5 H      UNL      1      -4.622      2.930      -0.135      1.00      0.00
               H
8 HETATM      6 H      UNL      1      -3.699      0.985      1.202      1.00      0.00
               H
9 HETATM      7 H      UNL      1      -3.621      0.932      -1.328      1.00      0.00
               H
10 HETATM     8 H      UNL      1      -1.258      3.797      1.217      1.00      0.00
               H
11 HETATM     9 H      UNL      1      -1.178      3.740      -1.314      1.00      0.00
               H
12 HETATM    10 H      UNL      1      0.213      0.288      -1.189      1.00      0.00
               H
13 HETATM    11 H      UNL      1      0.135      0.352      1.342      1.00      0.00
               H
14 HETATM    12 H      UNL      1      2.655      3.095      -1.186      1.00      0.00
               H
15 HETATM    13 H      UNL      1      3.578      1.161      0.165      1.00      0.00
               H
16 HETATM    14 H      UNL      1      2.574      3.167      1.343      1.00      0.00
               H
17 CONECT      1      2      5      6      7
18 CONECT      2      1      3      8      9
19 CONECT      3      2      4     10     11
20 CONECT      4      3     12     13     14
21 CONECT      5      1
22 CONECT      6      1
23 CONECT      7      1
24 CONECT      8      2
25 CONECT      9      2
26 CONECT     10      3
27 CONECT     11      3
28 CONECT     12      4
29 CONECT     13      4
30 CONECT     14      4

```

31	MASTER	0	0	0	0	0	0	0	0	14	0	14	0
32	END												

APPENDIX D

Two-Dimensional Rose-Potential Water

words

VITA

Gentry H. Smith

Candidate for the Degree of
Master of Science

Thesis: EXPLORING CRITICAL CONFORMATIONS

Major Field: Chemistry

Biographical:

Personal Data: Born in Olathe, KS in November 1993.

Education:

Received a Bachelors of Science in Chemistry at Southern Nazarene University in May 2016.

Completed the requirements for the degree of Master of Science with a major in Chemistry at Oklahoma State University in May 2018.

Experience:

Professional Affiliations:

American Chemical Society

Awards

Colonel Andre Whitely Scholarship in Chemistry