

# Index Fit Report

## North Atlantic Swordfish MSE

Adrian Hordyk [adrian@bluematterscience.com](mailto:adrian@bluematterscience.com)

May 28, 2020

## Contents

<b>Introduction</b>	<b>1</b>
<b>Indices and Fits</b>	<b>1</b>
Fleet & Survey Indices . . . . .	1
Combined Index . . . . .	9
<b>Summary of Statistical Properties</b>	<b>9</b>
Fleet & Survey Indices . . . . .	9
Combined Index . . . . .	13

## Introduction


This report summarizes the fits across the 288 OM's from the uncertainty grid to the fleet & survey 14 indices from the North Atlantic swordfish.

It also shows plots of the combined index (see SCRS/2017/137) against the predicted total biomass from each of the OM's.


## Indices and Fits

### Fleet & Survey Indices


The following plots show the observed index (black solid dots - left panel) and the observed index adjusted for an assumed 1% annual increase in catchability (black solid dots - right panel) for the 14 fleet and survey indices, with the model fits from each OM shown as colored lines (red = no environmental covariate, blue = an environmental covariate).




### Canada – Early




### Canada – Late




### Japan – Early




### Japan – Mid





# Japan – Late




# Portugal







### Age-2 Survey




### Age-3 Survey



### Age-4 Survey




### Age-5+ Survey





## Combined Index

The following plot show the combined index (black solid dots) with the standardized predicted total biomass from each OM shown as colored lines (red = no environmental covariate, blue = an environmental covariate). The OMs that were fitted to fleet & survey indices assuming a 1% annual increase in catchability are shown in the right panel.




## Summary of Statistical Properties

### Fleet & Survey Indices


#### No Increase in Catchability

Histograms of the statistical properties of the log-deviations from the fits to the 14 indices, including auto-correlation (ac), correlation (cor), root-mean-square-error (rmse), and standard deviation (sd), for the 144 OMs using the observed indices.



### **1% Annual Increase in Catchability**

Histograms of the statistical properties of the log-deviations from the fits to the 14 fleet & survey indices, including auto-correlation (ac), correlation (cor), root-mean-square-error (rmse), and standard deviation (sd), for the 144 OMs using the observed indices adjusted for an assumed 1% annual increase in catchability.



## Combined Index

Histograms of the statistical properties of the log-deviations from combined index and the predicted total biomass, including auto-correlation (ac), correlation (cor), root-mean-square-error (rmse), and standard deviation (sd). The bottom row shows the 144 OMs that were fitted to the observed indices that were adjusted for an assumed 1% annual increase in catchability.

