



# Normalized Difference Vegetation Index (NDVI)

## Disclaimer

---

- Though SSC Pacific makes every effort to perform quality assurance on its training materials, the material in this presentation may inadvertently include technical inaccuracies or other errors. We would be grateful if users notify us of any errors or inaccuracies they may find.
- The presentation contains references to links and to third-party websites. These are provided for the convenience and interest of users and this implies neither responsibility for, nor approval of, information contained in these websites on the part of the U.S. Government. The USG makes no warranty, either express or implied, as to the accuracy, availability or content of information, text, graphics in the links/third party websites. The USG has not tested any software located at these sites and does not make any representation as to the quality, safety, reliability or suitability of such software, nor does this presentation serve to endorse the use of such sites.


# Overview of Talk

---


- Overview/History of NDVI
- NDVI Calculation
- NDVI Uses
- OSSIM Band Algebra

# What is NDVI?

- The Normalized Difference Vegetation Index (NDVI) is a simple graphical indicator that can be used to assess whether the target being observed contains green vegetation or not
- Determines the density of green on a patch of land
- Based on the principle that when light strikes an object, light of different wavelengths are absorbed and reflected



average NDVI of October 2003



# NDVI Calculation

- Plants absorb light for photosynthesis
  - Absorbs greatly in the 400nm – 700nm range (VIS)
  - Reflects greatly in the 700nm – 1100nm range (NIR)
- How many leaves a plant has affects the amount of absorption/reflection

$$NDVI = \frac{(NIR - VIS)}{(NIR + VIS)}$$

- Calculations for NDVI always between [-1, 1]
  - No green leaves  $\approx 0$
  - Highest possible density  $\approx +1$
  - Clouds and snow  $< 0$


# NDVI Calculation - Sensitivities

---

- Atmospheric effects – specifically water vapor and aerosols
- Clouds – large clouds are easy to filter, but thin clouds may affect measurements, as well as the shadow of those clouds
- Soil Effects – reflectance of soil is directly related to water content; wet soil can effect measurements
- Anisotropic Effects – target geometry and observation time may effect measurements
- Spectral Effects – measurements will vary depending on the sensor used due to the inherent differences between each type of imaging sensor

## NDVI Used as an Indicator

- NDVI measurements allow for the quantification of absorbed radiation by vegetation
- Photosynthesis is directly related to absorbed photosynthetically active radiation
  - The more visible light absorbed, the more productive the plant
- A region's absorption and reflection of photosynthetically active radiation over a period of time can help to characterize the productivity of that region's vegetation
- When these characteristics are compared to data repositories, the health of a region for a particular time period can be compared that region's norm value





## NDVI Example Image

- LandSat7 Image –  
    `..\Day02\imagenes\*_HRF.fst`

### LandSat 7 Imagery

Band 1 Visible (0.45 – 0.52  $\mu\text{m}$ ) 30 m

Band 2 Visible (0.52 – 0.60  $\mu\text{m}$ ) 30 m

Band 3 Visible (0.63 – 0.69  $\mu\text{m}$ ) 30 m


Band 4 Near-Infrared (0.77 – 0.90  $\mu\text{m}$ ) 30 m

Band 5 Near-Infrared (1.55 – 1.75  $\mu\text{m}$ ) 30 m

Band 61 Low Gain Thermal (10.40 – 12.50  $\mu\text{m}$ ) 60 m

Band 62 High Gain Thermal (10.40 – 12.50  $\mu\text{m}$ ) 60 m

Band 7 Mid-Infrared (2.08 – 2.35  $\mu\text{m}$ ) 30 m





# NDVI Example Code

---

- `..\Day06\codigo\bandAlgebra.zip`
- Need to modify the batch files
- Build the project using Cmake
- Compile the solution using Visual Studio
- Copy and modify the runProject.bat file
- Run the project

## Resources and References

---

- Nasa Earth Observatory  
[http://earthobservatory.nasa.gov/Features/MeasuringVegetation/measuring\\_vegetation\\_1.php](http://earthobservatory.nasa.gov/Features/MeasuringVegetation/measuring_vegetation_1.php)
- Tucker, C.J. 1979. Red and photographic infrared linear combinations for monitoring vegetation. Remote Sensing of the Environment, v. 8, p.127-150.
- Jackson, R.D., P.N. Slater, and P.J. Pinter, 1983. Discrimination of growth and water stress in wheat by various vegetation indices through clear and turbid atmospheres. Remote Sensing of the Environment, v. 15, p.187-208.
- Tucker, C. J., W. W. Newcomb, S. O. Los, and S. D. Prince, 1991. Mean and inter-year variation of growing-season normalized difference vegetation index for the Sahel 1981-1989. International Journal of Remote Sensing, v. 12, p.1113-1115.

# Questions?

---

## Normalized Difference Vegetation Index (NDVI)

Mr. John Stastny  
SPAWAR Systems Center,  
Pacific  
Phone: 619-553-4564  
Email: [john.stastny@navy.mil](mailto:john.stastny@navy.mil)

Mr. Bryan Bagnall  
SPAWAR Systems Center,  
Pacific  
Phone: 619-553-4061  
Email: [bryan.bagnall@navy.mil](mailto:bryan.bagnall@navy.mil)

Mr. Lucas Keenan  
SPAWAR Systems Center,  
Pacific  
Phone: 619-553-3686  
Email: [lucas.keenan@navy.mil](mailto:lucas.keenan@navy.mil)