



Research report

The role of NMDA and AMPA/Kainate receptors in the consolidation of catalepsy sensitization

K. Riedinger^{a,*}, A. Kulak^{a,b,1}, W.J. Schmidt^a, A. von Ameln-Mayerhofer^a^a University of Tübingen, Neuropharmacology, Auf der Morgenstelle 28E, 72076 Tübingen, Baden-Wuerttemberg Germany^b Graduate School of Neural and Behavioural Sciences, International Max-Planck Research School, University of Tübingen, Tübingen, Germany

ARTICLE INFO

Article history:

Received 27 August 2010

Received in revised form

25 November 2010

Accepted 28 November 2010

Keywords:

Catalepsy sensitization

Consolidation

Excitatory amino acids

Haloperidol

NMDA-receptors

AMPA/Kainate-receptors

ABSTRACT

Daily injection of the dopamine D₂ receptor antagonist haloperidol is associated with the development of catalepsy sensitization in rats, which leads to a day to day increase of rigor and akinesia. The process of catalepsy sensitization incorporates different learning stages. Here we investigated the mechanisms underlying the consolidation of catalepsy sensitization. In particular, we asked whether NMDA- and non-NMDA (AMPA- and Kainate) receptors play a role in the consolidation of catalepsy sensitization. Accordingly, rats received post-training injections of the NMDA receptor antagonist MK-801 (single injection of either 0.1 mg/kg or 0.25 mg/kg; or a double injection of 0.1 mg/kg immediately and 30 min after test cessation) or of the AMPA/Kainate receptor antagonist GYKI 52466 (single injection of 5 mg/kg). Our results showed that the consolidation of catalepsy sensitization was decelerated by both glutamatergic AMPA/Kainate- and NMDA-receptor antagonists. With the higher MK-801 dosage, the deceleration was stronger, suggesting a dose dependent mechanism. We hence affirmed a role for the ionotropic glutamate receptors in the consolidation process of catalepsy sensitization.

© 2010 Elsevier B.V. All rights reserved.

1. Introduction

Haloperidol induced catalepsy (HIC) is a well accepted animal model for the Parkinsonian symptoms akinesia and rigor and can be reliably quantified by movement initiation latency measurement [1,2]. Antagonism of dopamine (DA) D₂ receptors with haloperidol leads to hyperactivity of the indirect, striatofugal pathway within the dorsal basal ganglia loop [3,4], disinhibition of the major output structures of the basal ganglia, and finally to an increased inhibition of thalamo-cortical neurons. Under constant conditions, repeated daily catalepsy testing with a preceding haloperidol injection leads to a gradual decline in motility, corresponding to a gradual increase in HIC [5]. Previous work has shown that this gradual increase in HIC is based on a specific learning process that is commonly referred to as behavioural sensitization. While the classically understood sensitization refers to the gradual increase of a behavioural effect to the repeated administration of a DA agonist – usually a psychostimulant [6,7] – sensitization processes are also involved in the progressive aggravation of rigor and akinesia in Parkinson's disease: repeated administration of a DA antagonist results in a day to day decrease in activity, i.e. an increase in catalepsy [8].

Thus, sensitization to stimulants and HIC might follow similar rules but in opposite directions, facilitating appropriate and suppressing inappropriate motor commands [9–11].

A recent study classified HIC-sensitization as a form of the so called “no-go” learning [12], one of two learning processes that are important during the completion of planned actions. Frank et al. [13] developed a series of neurocomputational models in which they demonstrated that two main populations of medium spiny neurons in the striatum have opposing effects. Whereas an activity of the so called “go” neurons facilitate the execution of a motor action (synaptically modulated by the DA D₁ Pathway), the activity of the “no-go” (modulated via the DA D₂ Pathway) neurons prevent or suppress the execution of competing motor actions. The comprehension of the selection of “go” (in our model the leaving of an inconvenient position) versus “no-go” (in our model persisting in an inconvenient position) mediated movements is important not only for control of daily movements but also for the understanding of diseases with disturbed action selection (i.e. in Parkinson's disease and schizophrenia).

Furthermore HIC-sensitization is described as an associative learning process, which is strongly dependent on the context [14–16] and/or the animal's internal state (interoceptive cues) [5]. The robustness of this association process was already shown [14,15]. The described properties of HIC-sensitization lead to the presumption that this learning paradigm follows the same rules as other associative learning processes [17,18]. In general, learning phases can be subdivided into acquisition (development of

* Corresponding author. Tel.: +49 70712974572; fax: +49 7071295144.

E-mail address: Katrin.Riedinger@web.de (K. Riedinger).¹ Present address: Center for Psychiatric Neuroscience, Department of Psychiatry, Lausanne University Hospital, Site de Cery, CH-1008 Prilly-Lausanne, Switzerland.

an association), consolidation (conversion from labile to more stable memory storage and hence the induction of long-term-potential (LTP) or long-term-depression (LTD)) and expression (retrieval of learned associations). In this study, we explicitly focussed on the consolidation of HIC-sensitization and the role of ionotropic glutamate receptors during that learning stage, since these receptors are evidently involved in processes of neuronal plasticity which plays an essential role in acquisition, consolidation and reconsolidation [19–26]. In the present study, we administered the selective non-competitive NMDA-receptor-antagonist MK-801 or the non-competitive AMPA/Kainate receptor-antagonist GYKI 52466 [27–29] directly after cessation of the catalepsy test to unequivocally assess whether the respective receptors are necessary for the consolidation of HIC-sensitization.

2. Materials and methods

2.1. Subjects

Male Sprague–Dawley rats (250–280 g bodyweight (BW), Charles River, Sulzfeld, Germany), were used in all experiments. Animals were housed in groups of four per cage under a 12/12 h light–dark cycle with restricted access to food (Sniff special-diet – standard dried pellets 12 g per animal per day) and water *ad libitum*. For the first experiment we used 7 or 8 animals per group, for the second experiment 10 animals per group and for the third experiment 12 animals per group.

2.2. Substances

Haloperidol (Janssen-Cilag GmbH, Neuss, Germany) was diluted in saline with a concentration of 0.5 mg/ml in the first and second experiment and in aqua injectabilia (Ampuwa, Fresenius Kabi, Bad Homburg, Germany) in a concentration of 0.25 mg/ml in the third experiment.

The NMDA receptor-antagonist [(5R,10S)-(+)-5-Methyl-10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5,10-imine hydrogen maleate] (dizocilpine hydrogen maleate (INN), MK-801, Sigma–Aldrich, Steinheim, Germany) was dissolved in saline. For the first experiment a concentration of 0.1 mg/ml (once or twice) was used and for the second experiment concentrations of 0.1 mg/ml and 0.25 mg/ml were used. All concentrations were related to free base. The above-named substances were administered intraperitoneally in an injection volume of 1 ml/kg BW.

The AMPA/Kainate receptor antagonist GYKI 52466 (a generous gift from Dr. Tarnawa, Budapest, Hungary) which was used in the third experiment was diluted in aqua injectabilia to a concentration of 5.0 mg/ml and administered intraperitoneally as 1.0 mg per kg BW. The concentration of GYKI 52466 was related to free base.

2.3. Behavioural testing

Catalepsy was first assessed on a horizontal bar (18 cm length, 0.8 cm diameter, 8 cm above the test table surface) by placing the animal's forepaws onto the bar. Immediately afterwards animals were tested on a vertical wire-mesh grid (25 cm × 40 cm surface area; 1 cm mesh width) by placing the animal with all four paws onto the grid. For both tests, the time was measured until the animal actively displaced one paw (descent latency). Testing took place during the light phase on nine consecutive days. The respective glutamate receptor antagonist was administered directly after the catalepsy test. Animals were tested 60 min after haloperidol or vehicle injection. To avoid physical fatigue of the animals, every catalepsy test was cut off after 180 s. As mentioned above, the injection of one glutamate receptor antagonist was applied directly after the grid test. For the first and second experiment this second injection was the NMDA receptor antagonist MK-801 or its vehicle. In the first experiment, one group received an additional injection 30 min after that second injection. For the third experiment, the AMPA receptor antagonist GYKI 52466 or its vehicle was applied.

2.3.1. Experiment 1

In the first experiment, a total of 23 animals were used which were subdivided into three groups. The first group (positive control group, $n = 8$) received a vehicle injection directly after the catalepsy test. The second group (1 × MK-801 group, $n = 8$) received one MK-801 injection (0.1 mg/kg BW) directly after the catalepsy test. The third group (2 × MK-801 group, $n = 7$) received two MK-801 injections, one directly after the catalepsy test and the second injection 30 min later.

2.3.2. Experiment 2

For the second experiment, we used four groups ($n = 10$ in each group). The positive control group received a haloperidol injection before and a vehicle injection after the catalepsy test.

The two groups which were used to test the effect of MK-801 had a haloperidol treatment first. A second injection was applied after the catalepsy test consisting of a MK-801 injection of 0.1 mg/kg BW within one treatment group (MK-801 0.1 group)

or 0.25 mg/kg BW in the other treatment group (MK-801 0.25 group). The negative control group received a vehicle injection at each time point instead of haloperidol and a post-test vehicle injection, respectively.

2.3.3. Experiment 3

For the third experiment, we used a total of 24 animals separated in two groups ($n = 12$ for each group). Like in the first and second experiment we had a positive control group that received only vehicle directly after the catalepsy test. The testing group (GYKI group) received the AMPA receptor antagonist GYKI 52466 5.0 mg/kg BW directly after the catalepsy test.

2.4. Statistical analysis

For all statistical tests we used the software GB STAT professional, version 7.0 (Dynamic Microsystems, Inc.). All data are presented as means ± standard errors (SEM). Data were analysed using a two-way analysis of variance (ANOVA, treatment × time) followed by post hoc Fisher's LSD test when appropriate.

3. Results

3.1. Experiment 1

All groups of the first experiment showed a significant development of HIC-sensitization in the bar test (Fig. 1A, two-way ANOVA: $F(8, 160)_{\text{time}} = 48.75$, $p < 0.0001$). However dependent on treatment, the development was different between groups ($F(16, 160)_{\text{treatment} \times \text{time}} = 1.99$, $p = 0.016$). For the grid test a significant effect over time was observed (Fig. 1B, two-way ANOVA: $F(8, 16)_{\text{time}} = 18.71$, $p < 0.0001$).

The descent latencies extracted from both testing apparatuses indicated an obvious delay of HIC-sensitization in the MK-801 treated groups compared to the positive control group with the stronger effect in the 2 × MK-801 group.

3.2. Experiment 2

In the second experiment the comparison between the negative control group and the positive control group was carried out separately in order to test the validity of the positive control group. Two-way ANOVA revealed a significant difference between the negative and positive control groups in the bar test ($F(1, 18)_{\text{treatment}} = 27.98$, $p < 0.0001$, $F(8, 144)_{\text{time}} = 6.53$, $p < 0.0001$, $F(8, 144)_{\text{treatment} \times \text{time}} = 6.44$, $p < 0.0001$) as well as in the grid test ($F(1, 8)_{\text{treatment}} = 19.75$, $p = 0.0003$). The positive control group showed significantly higher decent latency from day three to day nine ($p < 0.01$) in the bar test and from day two to day nine ($p < 0.01$) in the grid test according to post hoc test Fisher's LSD (significances not shown in Fig. 2A and B).

The positive control group and both MK-801 0.1 group and MK-801 0.25 group developed significant HIC-sensitization over time in the bar (Fig. 2A) and the grid (Fig. 2B) test (two-way ANOVA: bar: $F(8, 216)_{\text{time}} = 15.83$, $p < 0.0001$; grid: $F(8, 216)_{\text{time}} = 3.87$, $p = 0.0003$). Additionally, treatment with MK-801 led to a trend in reduction of descent latency compared to the positive control group in the bar test (two-way ANOVA: $F(2, 27)_{\text{treatment}} = 3.15$, $p = 0.058$).

Thus the higher MK-801 dosage led to a stronger deceleration of HICs than the lower MK-801 dose. Furthermore there was generally a weaker effect in the grid test concerning the HICs over all haloperidol treated groups.

3.3. Experiment 3

The comparison of the positive control group and GYKI group showed significant differences for the treatment and development of HIC-sensitization over time (two-way ANOVA: $F(1, 22)_{\text{treatment}} = 6.91$, $p = 0.0153$, $F(8, 176)_{\text{time}} = 13.48$, $p < 0.0001$) in the bar test (Fig. 3A)). In the grid test (Fig. 3B) a significant difference was observed over time ($F(8, 176)_{\text{time}} = 27.76$, $p < 0.0001$)




Fig. 1. Means \pm SEM of descent latency after daily injection of 0.5 mg/kg haloperidol tested on bar (A) and afterwards on grid (B). Post-test treatment: MK-801 (0.1 mg/kg) was administered either immediately after the test on the grid (half filled circles, 1 \times MK-801 group), or immediately after the test on the grid and 30 min later (filled circles, 2 \times MK-801 group). Positive control group received vehicle instead of MK-801 (open circles). Statistic: post hoc Fisher's LSD test, significant differences between the positive control and 1 \times MK-801 groups are marked by plus signs (* $p < 0.05$ and ** $p < 0.01$) and differences between positive control and 2 \times MK-801 groups are marked by asterisks (* $p < 0.05$ and ** $p < 0.01$).

and a trend in reduction of decent latency due to treatment with GYKI 52466 ($F(1, 22)_{\text{treatment}} = 3.49, p = 0.075$). Whereas the bar test showed a strong deceleration of HIC-sensitization due to treatment with GYKI 52466, this effect was not apparent in the grid test.

4. Discussion

Post-training administration of both NMDA and AMPA/Kainate receptor antagonists decelerated the development of HIC-sensitization supporting the notion that the glutamatergic system is involved in the consolidation of this learning process.

There is substantial evidence that the ionotropic NMDA- and AMPA/Kainate-receptors are essential for both the consolidation of learning [19,21,23–25,30–33] and the development of behavioural sensitization [34–36]. Concerning NMDA receptors, a significant number of studies describe serious learning impairments due to blockade of these receptors [31,32,37]. An impairment induced by NMDA receptor antagonists was especially shown in spatial mem-




Fig. 2. Descent latency after daily injection of vehicle (rectangles) or 0.5 mg/kg haloperidol (circles) tested on bar (A) and afterwards on grid (B) displayed as means \pm SEM. Post-test treatment: vehicle (rectangles, negative control group and blank circles, positive control group), 0.1 mg/kg MK-801 (half filled circles, MK-801 0.1 group) or 0.25 mg/kg MK-801 (filled circles, MK-801 0.25 group). Statistics: post hoc Fisher's LSD test, significant differences between the positive control and MK-801 0.1 groups are marked by plus signs (* $p < 0.05$ and ** $p < 0.01$) and differences between the positive control and MK-801 0.25 groups are marked by asterisks (* $p < 0.05$ and ** $p < 0.01$).

ory tasks [38,39], in visuo-spatial tasks [40], in fear conditioning [41], as well as in behavioural sensitization (reviewed in [36]). Concerning the blockade of behavioural sensitization by MK-801 most recent antagonist studies targeted the acquisition of learning. It should be noted that many learning processes develop in a state-dependent manner, i.e. the recall of stored information is possible only in the presence of a specific state which must be identical to the state during the storage of information, namely during learning [42,43]. Many substances and MK-801 explicitly do produce a strong and definite endogenous state, thus the expression of particular learned behavioural responses (like lever pressing or behavioural sensitization) can vary significantly, dependent on the state present on the test day [44,45]. Nevertheless there is most likely an involvement of NMDA receptors in sensitization development in specific and in behavioural plasticity processes in general, despite the risk of over-interpreting some results of co-administration studies. Given that MK-801 injections in our experiments were given after completion of the testing




Fig. 3. Means \pm SEM of descent latency after daily injection of 0.25 mg/kg haloperidol treated groups tested on bar (A) and grid (B). Post-test treatment: vehicle (blank circles, positive control group) or 5.0 mg/kg GYKI 52466 (filled circles, GYKI group). Statistics: post hoc Fisher's LSD test showed significances between both groups marked by asterisks (* $p < 0.05$ and ** $p < 0.01$).

procedure and thus did not co-occur with haloperidol injections, we can disregard an effect of state dependency. Worth mentioning is that previous studies of our group already excluded a role of NMDA-receptors during the acquisition phase of HIC-sensitization: neither non-competitive nor competitive NMDA-receptor antagonists blocked HIC-sensitization when administered previously to the test [5,46]. However, our approach to target explicitly the consolidation phase and to avoid the acquisition phase of sensitization learning cannot fully exclude the possibility that consolidation processes start already before or during the testing. Explicit targeting of consolidation or reconsolidation by pharmacological manipulation after testing is a well established method [47–51]. In this regard it should be made clear that our hypothesis describes corticostriatal synaptic plasticity which occurs due to activation of cortical inputs to the “no-go”-medium spiny neurons and which is isochronous with the haloperidol related disinhibition of this particular neuronal population. From our perspective, the cortical activation is generated by exposure of the animals to the test apparatus in a specific context (cf. [12]). Both catalepsy testing methods (bar and grid) are accepted in the catalepsy research; however, we obtained clearer results using the bar test apparatus compared to the grid test apparatus. This might be due to the fact that there is a difference between the effort for the animals in latch-

ing onto the grid with all four paws or only holding on the bar with the forepaws, while the hind paws are lying calmly on the table surface.

An important advantage of our treatment design (i.e. to administer the glutamatergic antagonists after testing), is that we can absolutely avoid side effects of the NMDA-antagonist administration. There is a bunch of evidence that NMDA-receptor antagonists (e.g. amantadine, memantine) do possess strong therapeutic potential for the treatment of Parkinson's disease [52–56]. MK-801 which has been used in our study reduces experimental parkinsonism in numerous studies as well [57–61]. Thus, by administration of MK-801 after the test, we could avoid the acute anticataleptic effects of the substance which would have blurred the results.

It should be stated at this point, that in case of a complete blockade of the consolidation of HIC-sensitization no “absolute therapy” effect was expected. In other words, even a severe pharmacological interference with the memory consolidation process and therefore sensitization blockade would result in a stable basal catalepsy response. It is not very probable that the basal firing rate of the striatal “no-go”-neurons would decline in the presence of a D2-receptor antagonist. Nevertheless, we observe only a marginal sensitization antagonism or more correctly, a deceleration: there is a slowdown rather than a blockade of the sensitization process, since most animals reached catalepsy levels comparable to positive controls at the end of each experiment. This might have occurred for three reasons: first, the used dose of MK-801 might be insufficient for a full blockade effect. In this case, higher concentrations of MK-801 might have been effective. However, we decided to use these dosages *a priori*, in order to avoid severe adverse or neuropathological effects (cf. [62,63]). With the dose regimen in experiment 1, we assumed that the stronger deceleration of HIC-sensitization in the animal group which received MK-801 twice was due to an accumulation of MK-801 rather than due to a temporal mechanism [this interpretation is also supported by pharmacokinetics data on MK-801 [64,65]]. We hence designed our second experiment in order to assess this possibility and animals were treated once immediately after the catalepsy-test with the higher dose of 0.25 mg/kg MK-801. The results of both experiments account consequently for a dose dependent attenuation or retardation of the catalepsy. Second, the investigated glutamatergic receptors might not be the only mechanistic players within the consolidation of HIC-sensitization. This idea is supported by the absence of effects of NMDA-antagonists during the acquisition of HIC-sensitization, i.e. parts of HIC-sensitization related neuronal plasticity occurs independently of NMDA receptors [5,46]. Third, blockade of NMDA-receptors only after the test, i.e. after test-induced cortical activation, might be simply insufficient for a full blockade of the related consolidation process. Then, consequently, the consolidation process would have started already before or during testing (or at least the role of NMDA-receptors within). As already discussed above it would then not be possible to distinguish between acquisition and consolidation phases.

In the third experiment we used the selective AMPA/Kainate receptor antagonist GYKI 52466. We used a low dose as well in order to avoid severe side effects [66]. As it turned out, this dose was already effective in attenuating the consolidation of HIC-sensitization, although the effect was not strong. Interestingly, the literature search does not deliver a clear-cut antiparkinsonian effect of AMPA-receptor antagonists. Systemic administration of AMPA receptor antagonists NBQX and GYKI 52466 did not have any direct anti-cataleptic potential when tested in an acute model of haloperidol treated rats [67]. Opposite results have been observed for NBQX [68] and a co-administration of NBQX and L-DOPA [69]. In another previous study, the AMPA/Kainate receptor antagonist GYKI 52466 did not alter the HIC acutely while it significantly antagonised the MK-801 induced locomotor hyperactivity [58].

5. Conclusion

HIC-sensitization is a robust and context dependent [15] learning process. Here we ascertain that the mechanisms underlying the consolidation of HIC-sensitization differ from the mechanisms that are active during acquisition with regards to NMDA and AMPA/Kainate receptor involvement. Whereas pre-test administration of glutamate antagonists had no effect on the acquisition phase of HIC-sensitization, we show for the first time that post-test administration of a glutamate antagonist interferes with the consolidation phase and decelerates the previously proposed “no-go”-learning within basal ganglia, a learning process evidently associated with the pathophysiology of Parkinson's disease and schizophrenia [70,71]. Hence, treatment with NMDA- or AMPA/Kainate receptor antagonists might be useful not only as acute therapeutics [72] but might additionally slow down the progressive aggravation of Parkinson symptomatology. Our results further provide an important contribution to the current knowledge of basal ganglia plasticity, especially that one mechanistic element of Parkinsonism-associated “no-go” learning is (during consolidation and not acquisition) dependent on functional NMDA and AMPA/Kainate receptors.

Acknowledgements

This manuscript is dedicated to our deceased supervisor and friend Werner J. Schmidt (deceased head of the department of neuropharmacology in Tuebingen, Germany). We thankfully remember his great efforts in research and his excellent considerate support, his good advices and ideas which led to the progression of our work. We wish to thank Mareike Fromm and Ulrich Ruess for technical assistance and Dr. I. Tarnawa (Inst. Drug Research, Hungary) for kindly providing GYKI 52466. Furthermore we wish to thank Nadine Lettfuß and Sonja Seeger-Armbruster for proof-reading and constructive discussion.

This work was supported by the Deutsche Forschungsgemeinschaft (SFB 550), K.R. was supported by a stipend of the Land Baden-Württemberg (Landesgraduiertenförderung).

References

- [1] Sanberg PR. Haloperidol-induced catalepsy is mediated by postsynaptic dopamine receptors. *Nature* 1980;284:472–3.
- [2] Sanberg PR, Bunsey MD, Giordano M, Norman AB. The catalepsy test: its ups and downs. *Behav Neurosci* 1988;102:748–59.
- [3] Blandini F, Nappi G, Tassorelli C, Martignoni E. Functional changes of the basal ganglia circuitry in Parkinson's disease. *Prog Neurobiol* 2000;62:63–88.
- [4] Schmidt WJ, Kretschmer BD. Behavioural pharmacology of glutamate receptors in the basal ganglia. *Neurosci Biobehav Rev* 1997;21:381–92.
- [5] Schmidt WJ, Tzschentke TM, Kretschmer BD. State-dependent blockade of haloperidol-induced sensitization of catalepsy by MK-801. *Eur J Neurosci* 1999;11:3365–8.
- [6] Robinson TE, Becker JB. Enduring changes in brain and behavior produced by chronic amphetamine administration: a review and evaluation of animal models of amphetamine psychosis. *Brain Res* 1986;396:157–98.
- [7] Stewart J, Badiani A. Tolerance and sensitization to the behavioral effects of drugs. *Behav Pharmacol* 1993;4:289–312.
- [8] Schmidt WJ, Beninger RJ. Behavioural sensitization in addiction, schizophrenia, Parkinson's disease and dyskinesia. *Neurotox Res* 2006;10:161–6.
- [9] Grillner S, Hellgren J, Menard A, Saitoh K, Wikstrom MA. Mechanisms for selection of basic motor programs—roles for the striatum and pallidum. *Trends Neurosci* 2005;28:364–70.
- [10] Kropotov JD, Etlinger SC. Selection of actions in the basal ganglia-thalamocortical circuits: review and model. *Int J Psychophysiol* 1999;31:197–217.
- [11] Mink JW. The basal ganglia: focused selection and inhibition of competing motor programs. *Prog Neurobiol* 1996;50:381–425.
- [12] Wiecki TV, Riedinger K, von Ameln-Mayerhofer A, Schmidt WJ, Frank MJ. A neurocomputational account of catalepsy sensitization induced by D2 receptor blockade in rats: context dependency, extinction, and renewal. *Psychopharmacology (Berl)* 2009;204:265–77.
- [13] Frank MJ, Scheres A, Sherman SJ. Understanding decision-making deficits in neurological conditions: insights from models of natural action selection. *Philos Trans R Soc Lond B Biol Sci* 2007;362:1641–54.
- [14] Amtage J, Schmidt WJ. Context-dependent catalepsy intensification is due to classical conditioning and sensitization. *Behav Pharmacol* 2003;14:563–7.
- [15] Klein A, Schmidt WJ. Catalepsy intensifies context-dependently irrespective of whether it is induced by intermittent or chronic dopamine deficiency. *Behav Pharmacol* 2003;14:49–53.
- [16] Banjaw MY, Mayerhofer A, Schmidt WJ. Anticataleptic activity of cathinone and MDMA (Ecstasy) upon acute and subchronic administration in rat. *Synapse* 2003;49:232–8.
- [17] Abel T, Lattal KM. Molecular mechanisms of memory acquisition, consolidation and retrieval. *Curr Opin Neurobiol* 2001;11:180–7.
- [18] Inda MC, Delgado-Garcia JM, Carrion AM. Acquisition, consolidation, reconsolidation, and extinction of eyelid conditioning responses require de novo protein synthesis. *J Neurosci* 2005;25:2070–80.
- [19] Ben Mamou C, Gamache K, Nader K. NMDA receptors are critical for unleashing consolidated auditory fear memories. *Nat Neurosci* 2006;9:1237–9.
- [20] Contractor A, Swanson G, Heinemann SF. Kainate receptors are involved in short- and long-term plasticity at mossy fiber synapses in the hippocampus. *Neuron* 2001;29:209–16.
- [21] Cooke SF, Bliss TV. Plasticity in the human central nervous system. *Brain* 2006;129:1659–73.
- [22] Cotman CW, Monaghan DT, Ganong AH. Excitatory amino acid neurotransmission: NMDA receptors and Hebb-type synaptic plasticity. *Annu Rev Neurosci* 1988;11:61–80.
- [23] Riedel G, Platt B, Micheau J. Glutamate receptor function in learning and memory. *Behav Brain Res* 2003;140:1–47.
- [24] Sadler R, Herzig V, Schmidt WJ. Repeated treatment with the NMDA antagonist MK-801 disrupts reconsolidation of memory for amphetamine-conditioned place preference. *Behav Pharmacol* 2007;18:699–703.
- [25] Sprengel R. Role of AMPA receptors in synaptic plasticity. *Cell Tissue Res* 2006;326:447–55.
- [26] Winters BD, Bussey TJ. Glutamate receptors in perirhinal cortex mediate encoding, retrieval, and consolidation of object recognition memory. *J Neurosci* 2005;25:4243–51.
- [27] Bleakman D, Ballyk BA, Schoepp DD, Palmer AJ, Bath CP, Sharpe EF, et al. Activity of 2,3-benzodiazepines at native rat and recombinant human glutamate receptors in vitro: stereospecificity and selectivity profiles. *Neuropharmacology* 1996;35:1689–702.
- [28] Paternain AV, Morales M, Lerma J. Selective antagonism of AMPA receptors unmasks kainate receptor-mediated responses in hippocampal neurons. *Neuron* 1995;14:185–9.
- [29] Vizi ES, Mike A, Tarnawa I. 2,3-Benzodiazepines (GYKI 52466 and analogs): negative allosteric modulators of ampa receptors. *CNS Drug Rev* 1996;2:91–126.
- [30] Bortolotto ZA, Lauri S, Isaac JT, Collingridge GL. Kainate receptors and the induction of mossy fibre long-term potentiation. *Philos Trans R Soc Lond B Biol Sci* 2003;358:657–66.
- [31] Collingridge GL, Bliss TVP. NMDA receptors—their role in long-term potentiation. *Trends Neurosci* 1987;10:288–93.
- [32] Danysz W, Wroblewski JT, Costa E. Learning impairment in rats by N-methyl-D-aspartate receptor antagonists. *Neuropharmacology* 1988;27:653–6.
- [33] Rao VR, Finkbeiner S. NMDA and AMPA receptors: old channels, new tricks. *Trends Neurosci* 2007;30:284–91.
- [34] Boudreau AC, Wolf ME. Behavioral sensitization to cocaine is associated with increased AMPA receptor surface expression in the nucleus accumbens. *J Neurosci* 2005;25:9144–51.
- [35] Sripada S, Gaytan O, Al-rahim S, Swann A, Dafny N. Dose-related effects of MK-801 on acute and chronic methylphenidate administration. *Brain Res* 1998;814:78–85.
- [36] Wolf ME. The role of excitatory amino acids in behavioral sensitization to psychomotor stimulants. *Prog Neurobiol* 1998;54:679–720.
- [37] Collingridge G. Synaptic plasticity. The role of NMDA receptors in learning and memory. *Nature* 1987;330:604–5.
- [38] Morris RG, Anderson E, Lynch GS, Baudry M. Selective impairment of learning and blockade of long-term potentiation by an N-methyl-D-aspartate receptor antagonist, AP5. *Nature* 1986;319:774–6.
- [39] Ward L, Mason SE, Abraham WC. Effects of the NMDA antagonists CPP and MK-801 on radial arm maze performance in rats. *Pharmacol Biochem Behav* 1990;35:785–90.
- [40] Harder JA, Aboobaker AA, Hodgetts TC, Ridley RM. Learning impairments induced by glutamate blockade using dizocilpine (MK-801) in monkeys. *Br J Pharmacol* 1998;125:1013–8.
- [41] Liu JL, Li M, Dang XR, Wang ZH, Rao ZR, Wu SX, et al. A NMDA receptor antagonist, MK-801 impairs consolidating extinction of auditory conditioned fear responses in a Pavlovian model. *PLoS One* 2009;4:e7548.
- [42] Colpaert FC. State dependency as a mechanism of central nervous system drug action. *NIDA Res Monogr* 1991;245–66.
- [43] Overton. State-dependent learning produced by depressant and atropine-like drugs. *Psychopharmacologia* 1966;6–31.
- [44] Jackson A, Koek W, Colpaert FC. NMDA antagonists make learning and recall state-dependent. *Behav Pharmacol* 1992;3:415–21.
- [45] Tzschentke TM, Schmidt WJ. Blockade of behavioral sensitization by MK-801: fact or artifact? A review of preclinical data. *Psychopharmacology (Berl)* 2000;151:142–51.

- [46] Lanis A, Schmidt WJ. NMDA receptor antagonists do not block the development of sensitization of catalepsy, but make its expression state-dependent. *Behav Pharmacol* 2001;12:143–9.
- [47] Borta A, Schwarting RK. Post-trial treatment with the nicotinic agonist metanicotine: differential effects in Wistar rats with high versus low rearing activity. *Pharmacol Biochem Behav* 2005;80:541–8.
- [48] Fricks-Gleason AN, Marshall JF. Post-retrieval beta-adrenergic receptor blockade: effects on extinction and reconsolidation of cocaine-cue memories. *Learn Mem* 2008;15:643–8.
- [49] Garg M, Holland HC. Consolidation and maze learning: the effects of post-trial injections of a depressant drug (pentobarbital sodium). *Psychopharmacologia* 1968;12:127–32.
- [50] McDonald RJ, Hong NS, Craig LA, Holahan MR, Louis M, Muller RU. NMDA-receptor blockade by CPP impairs post-training consolidation of a rapidly acquired spatial representation in rat hippocampus. *Eur J Neurosci* 2005;22:1201–13.
- [51] Rose JK, Rankin CH. Blocking memory reconsolidation reverses memory-associated changes in glutamate receptor expression. *J Neurosci* 2006;26:11582–7.
- [52] Danielczyk W. Twenty-five years of amantadine therapy in Parkinson's disease. *J Neural Transm Suppl* 1995;46:399–405.
- [53] Danysz W, Parsons CG, Kornhuber J, Schmidt WJ, Quack G. Aminoadamantanes as NMDA receptor antagonists and antiparkinsonian agents—preclinical studies. *Neurosci Biobehav Rev* 1997;21:455–68.
- [54] Klockgether T, Jacobsen P, Loschmann PA, Turski L. The antiparkinsonian agent bupropion is an N-methyl-D-aspartate antagonist. *J Neural Transm Park Dis Dement Sect* 1993;5:101–6.
- [55] Kornhuber J, Quack G, Danysz W, Jellinger K, Danielczyk W, Gsell W, et al. Therapeutic brain concentration of the NMDA receptor antagonist amantadine. *Neuropharmacology* 1995;34:713–21.
- [56] Uitti RJ, Rajput AH, Ahlskog JE, Offord KP, Schroeder DR, Ho MM, et al. Amantadine treatment is an independent predictor of improved survival in Parkinson's disease. *Neurology* 1996;46:1551–6.
- [57] Elliott PJ, Close SP, Walsh DM, Hayes AG, Marriott AS. Neuroleptic-induced catalepsy as a model of Parkinson's disease. II. Effect of glutamate antagonists. *J Neural Transm Park Dis Dement Sect* 1990;2:91–100.
- [58] Hauber W, Andersen R. The non-NMDA glutamate receptor antagonist GYKI 52466 counteracts locomotor stimulation and anticataleptic activity induced by the NMDA antagonist dizocilpine. *Naunyn Schmiedeberg Arch Pharmacol* 1993;348:486–90.
- [59] Kaur S, Ozer H, Starr M. MK 801 reverses haloperidol-induced catalepsy from both striatal and extrastriatal sites in the rat brain. *Eur J Pharmacol* 1997;332:153–60.
- [60] Papa SM, Engber TM, Boldry RC, Chase TN. Opposite effects of NMDA and AMPA receptor blockade on catalepsy induced by dopamine receptor antagonists. *Eur J Pharmacol* 1993;232:247–53.
- [61] Schmidt WJ, Bubser M. Anticataleptic effects of the N-methyl-D-aspartate antagonist MK-801 in rats. *Pharmacol Biochem Behav* 1989;32:621–3.
- [62] Allen HL, Iversen LL. Phencyclidine, dizocilpine, and cerebrocortical neurons. *Science* 1990;247:221.
- [63] Hashimoto K, Fujimura M, Yamagami K. Dizocilpine-induced neuropathological changes in rat retrosplenial cortex are reversed by subsequent clozapine treatment. *Life Sci* 2000;66:1071–8.
- [64] Hucker HB, Hutt JE, White SD, Arison BH, Zacchei AG. Disposition and metabolism of (+)-5-methyl-10,11-dihydro-5H-dibenzo[a,d] cyclohepten-5,10-imine in rats, dogs, and monkeys. *Drug Metab Dispos* 1983;11:54–8.
- [65] Vezzani A, Serafini R, Stasi MA, Caccia S, Conti I, Tridico RV, et al. Kinetics of MK-801 and its effect on quinolinic acid-induced seizures and neurotoxicity in rats. *J Pharmacol Exp Ther* 1989;249:278–83.
- [66] O'Hearn E, Molliver ME. Administration of a non-NMDA antagonist, GYKI 52466, increases excitotoxic Purkinje cell degeneration caused by ibogaine. *Neuroscience* 2004;127:373–83.
- [67] Zadow B, Schmidt WJ. The AMPA antagonists NBQX and GYKI 52466 do not counteract neuroleptic-induced catalepsy. *Naunyn Schmiedeberg Arch Pharmacol* 1994;349:61–5.
- [68] Klockgether T, Turski L, Honore T, Zhang ZM, Gash DM, Kurlan R, et al. The AMPA receptor antagonist NBQX has antiparkinsonian effects in monoamine-depleted rats and MPTP-treated monkeys. *Ann Neurol* 1991;30:717–23.
- [69] Loschmann PA, Lange KW, Kunow M, Rettig KJ, Jahnig P, Honore T, et al. Synergism of the AMPA-antagonist NBQX and the NMDA-antagonist CPP with L-dopa in models of Parkinson's disease. *J Neural Transm Park Dis Dement Sect* 1991;3:203–13.
- [70] Cools R, Altamirano L, D'Esposito M. Reversal learning in Parkinson's disease depends on medication status and outcome valence. *Neuropsychologia* 2006;44:1663–73.
- [71] Frank MJ. Schizophrenia: a computational reinforcement learning perspective. *Schizophr Bull* 2008;34:1008–11.
- [72] Bowers MS, Chen BT, Bonci A. AMPA receptor synaptic plasticity induced by psychostimulants: the past, present, and therapeutic future. *Neuron* 2010;67:11–24.