

IDS 702: MODULE 4.3

MULTILEVEL/HIERARCHICAL LINEAR MODELS (ILLUSTRATION I)

DR. OLANREWaju MICHAEL AKANDE

THE RADON ANALYSIS

There are 919 total observations in the data. The data is in the file `Radon.txt` on Sakai.

Variable	Description
radon	radon levels for each house
log_radon	log(radon)
state	state
floor	lowest living area of each house: 0 for basement, 1 for first floor
countyname	county names
countyID	ID for the county names (1-85)
fips	state + county fips code
uranium	county-level soil uranium
log_uranium	log(uranium)

THE RADON ANALYSIS

```
Radon <- read.csv("data/Radon.txt", header = T, sep = ",")
Radon$floor <- factor(Radon$floor, levels = c(0, 1), labels = c("Basement", "First Floor"))
str(Radon)
```

```
## 'data.frame':    919 obs. of  9 variables:
## $ radon      : num  2.2 2.2 2.9 1 3.1 2.5 1.5 1 0.7 1.2 ...
## $ state      : Factor w/ 1 level "MN": 1 1 1 1 1 1 1 1 1 1 ...
## $ log_radon  : num  0.788 0.788 1.065 0 1.131 ...
## $ floor      : Factor w/ 2 levels "Basement","First Floor": 2 1 1 1 1 1 1 1 1 1 ...
## $ countyname : Factor w/ 85 levels "AITKIN","ANOKA",...: 1 1 1 1 2 2 2 2 2 2 ...
## $ countyID   : int   1 1 1 1 2 2 2 2 2 2 ...
## $ fips       : int  27001 27001 27001 27001 27003 27003 27003 27003 27003 27003 ...
## $ uranium    : num  0.502 0.502 0.502 0.502 0.429 ...
## $ log_uranium: num  -0.689 -0.689 -0.689 -0.689 -0.847 ...
```

```
head(Radon)
```

```
##   radon state log_radon      floor countyname countyID  fips  uranium
## 1   2.2    MN 0.7884574 First Floor    AITKIN         1 27001 0.502054
## 2   2.2    MN 0.7884574   Basement    AITKIN         1 27001 0.502054
## 3   2.9    MN 1.0647107   Basement    AITKIN         1 27001 0.502054
## 4   1.0    MN 0.0000000   Basement    AITKIN         1 27001 0.502054
## 5   3.1    MN 1.1314021   Basement    ANOKA          2 27003 0.428565
## 6   2.5    MN 0.9162907   Basement    ANOKA          2 27003 0.428565
##   log_uranium
## 1 -0.6890476
## 2 -0.6890476
## 3 -0.6890476
## 4 -0.6890476
## 5 -0.8473129
## 6 -0.8473129
```

THE RADON ANALYSIS


```
table(Radon$countyname) #we don't have enough data in some counties, so we should look to borrow information across counties.
```

```
##
##      AITKIN      ANOKA      BECKER      BELTRAMI
##      4          52         3          7
##      BENTON      BIG STONE      BLUE EARTH      BROWN
##      4          3          14         4
##      CARLTON      CARVER      CASS      CHIPPEWA
##      10         6          5          4
##      CHISAGO      CLAY      CLEARWATER      COOK
##      6          14         4          2
##      COTTONWOOD      CROW WING      DAKOTA      DODGE
##      4          12         63         3
##      DOUGLAS      FARIBAUT      FILLMORE      FREEBORN
##      9          6          2          9
##      GOODHUE      HENNEPIN      HOUSTON      HUBBARD
##      14         105         6          5
##      ISANTI      ITASCA      JACKSON      KANABEC
##      3          11         5          4
##      KANDIYOHI      KITTSON      KOOCHICHING      LAC QUI PARLE
##      4          3          7          2
##      LAKE LAKE OF THE WOODS      LE SUEUR      LINCOLN
##      9          4          5          4
##      LYON      MAHNOMEN      MARSHALL      MARTIN
##      8          1          9          7
##      MCLEOD      MEEKER      MILLE LACS      MORRISON
##      13         5          2          9
##      MOWER      MURRAY      NICOLLET      NOBLES
##      13         1          4          3
##      NORMAN      OLMSTED      OTTER TAIL      PENNINGTON
##      3          23         8          3
##      PINE      PIPESTONE      POLK      POPE
##      6          4          4          2
##      RAMSEY      REDWOOD      RENVILLE      RICE
##      32         5          3          11
##      ROCK      ROSEAU      SCOTT      SHERBURNE
##      2          14         13         8
##      SIBLEY      ST LOUIS      STEARNS      STEELE
##      4          116        25         10
##      STEVENS      SWIFT      TODD      TRAVERSE
##      2          4          3          4
##      WABASHA      WADENA      WASECA      WASHINGTON
##      7          5          4          46
##      WATONWAN      WILKIN      WINONA      WRIGHT
##      3          1          13         13
##      YELLOW MEDICINE
```

THE RADON ANALYSIS

The raw radon levels can only take on positive values.

```
ggplot(Radon,aes(radon)) +  
  geom_histogram(aes(y=..density..),color="black",linetype="dashed",  
                 fill=rainbow(15),bins=15) + theme(legend.position="none") +  
  geom_density(alpha=.25, fill="lightblue") + scale_fill_brewer(palette="Blues") +  
  labs(title="Distribution of Radon Levels",y="Radon") + theme_classic()
```




Obviously very skewed.

THE RADON ANALYSIS

Let's look at `log_radon` instead.

```
ggplot(Radon, aes(log_radon)) +  
  geom_histogram(aes(y=..density..), color="black", linetype="dashed",  
                fill=rainbow(15), bins=15) + theme(legend.position="none") +  
  geom_density(alpha=.25, fill="lightblue") + scale_fill_brewer(palette="Blues") +  
  labs(title="Distribution of Log Radon Levels", y="Log Radon") + theme_classic()
```




THE RADON ANALYSIS

Are there any variations of radon levels by county? There are too many counties, so, let's do it for a random sample of counties.

```
set.seed(1000)
sample_county <- sample(unique(Radon$countyname),25,replace=F)
ggplot(Radon[is.element(Radon$countyname,sample_county),],
       aes(x=countyname, y=log_radon, fill=countyname)) +
  geom_boxplot() +
  labs(title="Log radon levels by county",
       x="County",y="Log Radon") + theme_classic() +
  theme(legend.position="none",axis.text.x = element_text(angle = 90))
```

THE RADON ANALYSIS




Looks like the levels vary by county. However, there are many counties with very little data.

THE RADON ANALYSIS

Let's focus on counties with at least 10 houses.

```
sample_county <- which(table(Radon$countyID) > 10)
ggplot(Radon[is.element(Radon$countyID,sample_county),],
       aes(x=countyname, y=log_radon, fill=countyname)) +
  geom_boxplot() +
  labs(title="Log radon levels by county",
       x="County",y="Log Radon") + theme_classic() +
  theme(legend.position="none",axis.text.x = element_text(angle = 90))
```

THE RADON ANALYSIS




What can you conclude from this plot?

THE RADON ANALYSIS

Next, the relationship with `floor`, the only individual-level (different observation for each house) variable we have.

```
ggplot(Radon, aes(x=floor, y=log_radon, fill=floor)) +  
  geom_boxplot() + scale_fill_brewer(palette="Greens") +  
  labs(title="Log radon vs floor", x="Lowest living area of each house", y="Log Radon") +  
  theme_classic() + theme(legend.position="none")
```




THE RADON ANALYSIS

Let's look at the same relationship for a random sample of counties.

```
sample_county <- sample(unique(Radon$countyname),8,replace=F)
ggplot(Radon[is.element(Radon$countyname,sample_county),],
       aes(x=floor, y=log_radon, fill=floor)) +
  geom_boxplot() +
  scale_fill_brewer(palette="Greens") +
  labs(title="Log radon vs floor by county",
       x="Lowest living area of each house",y="Log Radon") +
  theme_classic() + theme(legend.position="none") +
  facet_wrap( ~ countyname,ncol=4)
```

THE RADON ANALYSIS




Again, not enough data for some counties.

THE RADON ANALYSIS

Let's focus on counties with at least 15 houses.

```
sample_county <- which(table(Radon$countyID) > 15)
ggplot(Radon[is.element(Radon$countyID,sample_county),],
       aes(x=floor, y=log_radon, fill=floor)) +
  geom_boxplot() +
  scale_fill_brewer(palette="Greens") +
  labs(title="Log radon vs floor by county",
       x="Lowest living area of each house",y="Log Radon") +
  theme_classic() + theme(legend.position="none") +
  facet_wrap( ~ countyname,ncol=4)
```

THE RADON ANALYSIS



Even though the overall direction is the same, it looks like the actual differences between floor = 0 and floor = 1 differs for some counties.

THE RADON ANALYSIS

- Let's start by only focusing on `floor`.
- We will try a varying-slope, varying-intercept linear model.
- Let y_{ij} and x_{1ij} be the log radon level and indicator variable `floor` respectively for house i in county j .
- Mathematically, we have

$$\begin{aligned}y_{ij} &= (\beta_0 + \gamma_{0j}) + (\beta_1 + \gamma_{1j})x_{1ij} + \epsilon_{ij}; \quad i = 1, \dots, n_j; \quad j = 1, \dots, 85 \\ \epsilon_{ij} &\sim N(0, \sigma^2) \\ (\gamma_{0j}, \gamma_{1j}) &\sim N_2(\mathbf{0}, \Sigma).\end{aligned}$$

- Alternative representation:

$$\begin{aligned}\log(\text{radon}_{ij}) &= (\beta_0 + \gamma_{0j}) + (\beta_1 + \gamma_{1j}) \text{floor}_{ij} + \epsilon_{ij}; \quad i = 1, \dots, n_j; \quad j = 1, \dots, 85 \\ \epsilon_{ij} &\sim N(0, \sigma^2) \\ (\gamma_{0j}, \gamma_{1j}) &\sim N_2(\mathbf{0}, \Sigma).\end{aligned}$$

THE RADON ANALYSIS

- We skipped this before but Σ actually takes the form

$$\Sigma = \begin{bmatrix} \tau_0^2 & \rho\tau_0\tau_1 \\ \rho\tau_0\tau_1 & \tau_1^2 \end{bmatrix}$$

where

- τ_0^2 describes the within county variation attributed to the random/varying intercept,
- τ_1^2 describes the within county variation attributed to the random/varying slope (that is, floor), and
- ρ describes the correlation between γ_{0j} and γ_{1j} .

THE RADON ANALYSIS

In R, we have

```
Model1 <- lmer(log_radon ~ floor + (floor | countyname), data = Radon)
summary(Model1)
```

```
## Linear mixed model fit by REML ['lmerMod']
## Formula: log_radon ~ floor + (floor | countyname)
##      Data: Radon
##
## REML criterion at convergence: 2168.3
##
## Scaled residuals:
##      Min       1Q   Median       3Q      Max
## -4.4044 -0.6224  0.0138  0.6123  3.5682
##
## Random effects:
##      Groups      Name                Variance Std.Dev. Corr
## countyname (Intercept)      0.1216   0.3487
##              floorFirst Floor 0.1180   0.3436  -0.34
## Residual                    0.5567   0.7462
## Number of obs: 919, groups:  countyname, 85
##
## Fixed effects:
##              Estimate Std. Error t value
## (Intercept)      1.46277   0.05387  27.155
## floorFirst Floor -0.68110   0.08758  -7.777
##
## Correlation of Fixed Effects:
##              (Intr)
## florFrstFlr -0.381
```

INTERPRETATION OF FIXED EFFECTS

- Intuitively, we have an overall "average" regression line for all houses across all counties in Minnesota which has slope -0.68 and intercept 1.46.
- That is, the general estimated line for any of the houses in Minnesota is:

$$\log(\widehat{\text{radon}}_i) = 1.46 - 0.68 \times \text{floor}_i$$

- For any house in Minnesota with a basement as the lowest living area, the baseline radon level is $e^{1.46} = 4.31$.
- Then, for any house in Minnesota, having a first floor as the lowest living area, instead of a basement, reduces the radon level by a multiplicative effect of $e^{-0.68} = 0.51$, that is, about a 49% reduction.
- However, if the house is in Dakota county for example, we also need to add on the random intercepts and slopes for that county.

INTERPRETATION OF FIXED EFFECTS

- For Dakota county, we have

```
(ranef(Model1)$countyname)["DAKOTA",]
```

```
##           (Intercept) floorFirst Floor  
## DAKOTA   -0.1099069      -0.08787551
```

so that the estimated regression line for Dakota county is actually

$$\log(\widehat{\text{radon}}_i) = (1.46 - 0.11) + (-0.68 - 0.09) \times \text{floor}_i = 1.35 - 0.77 \times \text{floor}_i$$

- Thus, for any house in Dakota county in Minnesota with a basement as the lowest living area, the baseline radon level is actually $e^{1.35} = 3.86$, which is **lower than the overall state wide average**.
- And for any house in Dakota county in Minnesota, having the first floor be the lowest living area then reduces the radon level by a multiplicative effect of $e^{-0.77} = 0.46$, that is about a 54% reduction, **more than the overall state wide effect**.

THE RADON ANALYSIS

Again,


```
summary(Model1)
```

```
## Linear mixed model fit by REML ['lmerMod']
## Formula: log_radon ~ floor + (floor | countyname)
##    Data: Radon
##
## REML criterion at convergence: 2168.3
##
## Scaled residuals:
##      Min       1Q   Median       3Q      Max
## -4.4044 -0.6224  0.0138  0.6123  3.5682
##
## Random effects:
##   Groups      Name                Variance Std.Dev. Corr
##   countyname (Intercept)          0.1216   0.3487
##             floorFirst Floor 0.1180   0.3436  -0.34
##   Residual                        0.5567   0.7462
## Number of obs: 919, groups:  countyname, 85
##
## Fixed effects:
##              Estimate Std. Error t value
## (Intercept)    1.46277    0.05387  27.155
## floorFirst Floor -0.68110    0.08758  -7.777
##
## Correlation of Fixed Effects:
##              (Intr)
## florFrstFlr -0.381
```

INTERPRETATION OF RANDOM EFFECTS

- The estimated standard error $\hat{\sigma} = 0.75$ describes the unexplained within-county variation.
- The estimated $\hat{\tau}_0 = 0.35$ describes the within county variation attributed to the random intercept.
- The estimated $\hat{\tau}_1 = 0.34$ describes the within-county variation attributed to the random slope (the predictor, floor).
- Those two sources of within county variation are actually quite similar.
- The estimated correlation between γ_{0j} and γ_{1j} is $\hat{\rho} = -0.34$.
- You can visualize the random effects by typing `dotplot(ranef(Model1, condVar=TRUE))$countname` in R.
- So many counties! So, you will need to zoom out on your computer.

INTERPRETATION OF RANDOM EFFECTS



WHAT'S NEXT?

MOVE ON TO THE READINGS FOR THE NEXT MODULE!