

# Guided Diffusion Quantum Monte Carlo for Calculating Zero Point Energies

Simon Neidhart

Department of Physics, University of Basel

November 28, 2021

# Table of Contents

1. Introduction
2. The DQMC Algorithm
3. The Guiding Wave Function
4. Input and Output
5. Results
  - 5.1  $\text{C}_2\text{H}_6$
  - 5.2  $\text{C}_{12}\text{H}_{10}\text{O}$
  - 5.3  $\text{H}_2\text{@C}_{60}$
6. Conclusions and Improvements

# Introduction

- ▶ DQMC solves the Schrödinger equation and gives the quantum-mechanical ground state energy to then calculate the zero point energy (ZPE).
- ▶ A Gaussian guiding wave function is used to improve the performance.
- ▶ DFTB+ is used for calculation the energy.[1]

## The Unguided DQMC Algorithm [2]

- ▶ The stationary solution of the diffusion equation satisfies the Schrödinger equation and has the propagator  $G(x, y; \Delta t) = \frac{1}{\sqrt{2\pi\Delta t}} e^{-(x-y)^2/(2\Delta t)} e^{-\Delta t(V(y)-E_T)}$ .
- ▶ We simulate an ensemble of walkers to reach this stationary solution and the Schrödinger equation is solved.
- ▶ The propagator consists of a diffusion term  $\frac{1}{\sqrt{2\pi\Delta t}} e^{-(x-y)^2/(2\Delta t)}$  and a branching term  $e^{-\Delta t(V(y)-E_T)}$ .
- ▶ The diffusion term is simulated by randomly displacing the walker from its previous position according to a Gaussian distribution.
- ▶ The branching term updates the weight of a walker according to which the walker survives, reproduces or dies.
- ▶ The trial energy  $E_T$  is then adjusted to keep the population of walkers stable.
- ▶ With time, the trial energy  $E_T$  converges.

# The Guiding Wave Function [3]

- ▶ Harmonic approximation of the potential energy surface with the Hessian matrix.
- ▶ Gaussian guiding wave functions along the normal modes  $\psi_T(\mathbf{x}) = \mathcal{N}e^{-\mathbf{x}^2/(2\sigma^2)}$ .
- ▶ The width of the Gaussian is given by  $\sigma^2 = \frac{1}{\omega_m^2}$ , where  $\omega^2$  is the corresponding eigenvalue of the Hessian matrix.
- ▶ This is incorporated to the algorithm with a drift velocity  $\mathbf{v}(\mathbf{x}) = \nabla_{\mathbf{x}}\psi_T(\mathbf{x})$  which is added to the pure diffusion and a kinetic energy term in the local energy  $E_L(\mathbf{x}) = (H\psi_T(\mathbf{x}))/\psi_T(\mathbf{x})$  which replaces the potential energy in the branching step.

# The Guiding Wave Function




Fig. 1. Typical dimension in the coordinate system of the normal modes.

# Input and Output

## Input:

- ▶ Equilibrium geometry of the atoms (must not be perfectly accurate as geometry will be optimized by DFTB+ in the beginning)
- ▶ Masses of the atoms

## Output:

- ▶ Zero point energy
- ▶ Walker positions give a sample of the nucleonic wave function

# Results

$\text{C}_2\text{H}_6$  (first test case)




Fig. 2. Input structure of Ethane

First test case to develop the algorithm and check the result with a literature value.



# Results

$\text{C}_2\text{H}_6$  (first test case)

Simulation with 1000 walkers for 2000 time steps




Fig. 3. Fluctuation of the ZPE after equilibration.

ZPE = 0.07356 hartree with a standard deviation of 0.00060 hartree. The literature value is 0.073927 hartree [4].

# Results

$\text{C}_2\text{H}_6$  (first test case)

Simulation with 1000 walkers for 2000 time steps




Fig. 4. Convergence of the ZPE after equilibration.

Simulation time: 14 hours.  $\approx 0.02$  seconds per energy calculation.

# Results

$C_{12}H_{10}O$




Fig. 5. Input structure of 2-Phenylphenol

Slightly larger molecule to test the performance of the algorithm.

# Results



Simulation with 1000 walkers for 2000 time steps




Fig. 6. Fluctuation of the ZPE after equilibration.

ZPE = 0.18184 hartree with a standard deviation of 0.00117 hartree.

# Results

$\text{C}_{12}\text{H}_{10}\text{O}$

Simulation with 1000 walkers for 2000 time steps




Fig. 7. Convergence of the ZPE after equilibration.

Simulation time: 57 hours.  $\approx 0.11$  seconds per energy calculation.

# Results

$\text{H}_2@\text{C}_{60}$




Fig. 8. Input structure of the dihydrogen endofullerene

The dihydrogen endofullerene is expected to show strong anharmonic effects on the zero point energy.

# Results

H<sub>2</sub>@C<sub>60</sub>

Simulation with 1000 walkers for 2000 time steps




Fig. 9. Fluctuation of the ZPE after equilibration.

ZPE = 0.05914 hartree with a standard deviation of 0.00569 hartree.

# Results

H<sub>2</sub>@C<sub>60</sub>

Simulation with 1000 walkers for 2000 time steps




Fig. 10. Convergence of the ZPE after equilibration.

Simulation time: 51 hours.  $\approx 0.10$  seconds per energy calculation.



# Conclusions and Improvements

## Conclusions:

- ▶ DQMC delivers moderately accurate results for the ZPE in reasonable time.
- ▶ DQMC is very versatile and can calculate the ZPE of most molecules.

## Improvements:

- ▶ Adjustments for periodic structures
- ▶ Parallelization
- ▶ Making it more user-friendly

# References



B. Hourahine, B. Aradi, V. Blum, F. Bonafé, A. Buccheri, C. Camacho, C. Cevallos, M. Y. Deshayé, T. Dumitrică, A. Dominguez, S. Ehlert, M. Elstner, T. van der Heide, J. Hermann, S. Irle, J. J. Kranz, C. Köhler, T. Kowalczyk, T. Kubař, I. S. Lee, V. Lutsker, R. J. Maurer, S. K. Min, I. Mitchell, C. Negre, T. A. Niehaus, A. M. N. Niklasson, A. J. Page, A. Pecchia, G. Penazzi, M. P. Persson, J. Řezáč, C. G. Sánchez, M. Sternberg, M. Stöhr, F. Stuckenberg, A. Tkatchenko, V. W.-z. Yu, and T. Frauenheim.

Dftb+, a software package for efficient approximate density functional theory based atomistic simulations.

*The Journal of Chemical Physics*, 152(12):124101, 2020.



Anne B. McCoy.

Diffusion monte carlo approaches for investigating the structure and vibrational spectra of fluxional systems.

*International Reviews in Physical Chemistry*, 25(1-2):77–107, 2006.



Julien Toulouse, Roland Assaraf, and Cyrus J. Umrigar.

Chapter fifteen - introduction to the variational and diffusion monte carlo methods.

In Philip E. Hoggan and Telhat Ozdogan, editors, *Electron Correlation in Molecules – ab initio Beyond Gaussian Quantum Chemistry*, volume 73 of *Advances in Quantum Chemistry*, pages 285 – 314. Academic Press, 2016.



Amir Karton, Branko Ruscic, and Jan M.L. Martin.

Benchmark atomization energy of ethane: Importance of accurate zero-point vibrational energies and diagonal born–oppenheimer corrections for a ‘simple’