

Insomnia and Emotional Memory

Van Someren, Eus¹

¹ Netherlands Institute for Neuroscience, Netherlands

TO CITE

Van Someren, E. (2023). Insomnia and Emotional Memory. *Proceedings of the Paris Institute for Advanced Study*, 14. <https://doi.org/10.5072/zenodo.1156023>

PUBLICATION DATE

07/06/2019

ABSTRACT

Sleep and Memory. Paris IAS, 6-7 June 2019 - Session 6 - In Search of Lost Sleep

Insomnia is the second-most prevalent mental disorder and the primary modifiable risk factor for depression, anxiety disorders and PTSD ([Blanken et al., 2019](#)). In search of brain mechanisms, circadian and a homeostatic processes seem surprisingly intact. The hyperalert insomniac brain differs strongly from the hardly awake sleep-deprived brain ([Wei et al., 2016](#), [Stoffers et al., 2014](#)). Big data psychometrics (sleepregistry.org) showed different insomnia types, all involving distributed deviations in brain circuits underlying stable characteristics of affect ([Blanken et al., 2019](#)).

HD-EEG, MRI and animal model studies converge to suggest that the core issue causing insomnia is a deficiency in overnight processing of memories of distressing experiences ([Blanken et al., 2019](#), [Wassing et al., 2019](#)). Rapid Eye Movement (REM) sleep plays an important role in the reorganization of emotional memory circuits. Uniquely during the transition to REM sleep, and throughout REM sleep, the Locus Coeruleus (LC) is inhibited ([Swift et al., 2018](#)). This time window of low noradrenaline release facilitates synaptic depotentiation ([Swift et al., 2018](#), [Vanderheyden et al., 2014](#)). Meanwhile, increased activity

occurs in (para)limbic regions including amygdalae and anterior cingulate cortex, which is selectively activated by the claustrum during REM sleep ([Luppi et al., 2017](#)).

However, in insomnia and related mental disorders, abundant EEG arousals during sleep indicate insufficient inhibition of LC activity. Abiding LC activity during sleep abolishes the only noradrenaline-free time window available to the brain including its unique balance between synaptic potentiation and depotentiation during nocturnal replay and neuronal network adaptation. We posit that this in particular interferes with overnight processing of emotional distress ([Wassing et al., 2019](#), [Wassing et al., 2019](#)), resulting in daytime hyperactivation of salience- and emotional- circuits. This new model of circuits involved in insomnia is supported by GWAS ([Jansen et al., 2019](#)): risk genes for insomnia turn out to be expressed in cell types and brain areas identified with neuroimaging, including the claustrum and anterior cingulate cortex that activate during REM sleep ([Stoffers et al., 2014](#), [Wassing et al., 2019](#), [Luppi et al., 2017](#), [Altena et al., 2008](#)).

People with insomnia have developed a brain optimally wired to stay alert, at the cost of suffering bad sleep.

Bibliography

Altena, E., Van Der Werf, Y. D., Sanz-Arigita, E. J., Voorn, T. A., Rombouts, S. A. R. B., Kuijter, J. P. A., & Van Someren, E. J. W. (2008). Prefrontal Hypoactivation and Recovery in Insomnia. *Sleep*, 31(9), 1271–1276. <https://doi.org/10.5665/sleep/31.9.1271>

Blanken, T. F., Benjamins, J. S., Borsboom, D., Vermunt, J. K., Paquola, C., Ramautar, J., Dekker, K., Stoffers, D., Wassing, R., Wei, Y., & Van Someren, E. J. W. (2019). Insomnia disorder subtypes derived from life history and traits of affect and personality. *The Lancet. Psychiatry*, 6(2), 151–163. [https://doi.org/10.1016/S2215-0366\(18\)30464-4](https://doi.org/10.1016/S2215-0366(18)30464-4)

Blanken, T. F., Van Der Zweerde, T., Van Straten, A., Van Someren, E. J. W., Borsboom, D., & Lancee, J. (2019). Introducing Network Intervention Analysis to Investigate Sequential, Symptom-Specific Treatment Effects: A Demonstration in Co-Occurring Insomnia and Depression. *Psychotherapy and Psychosomatics*, 88(1), 52–54. <https://doi.org/10.1159/000495045>

Jansen, P. R., Watanabe, K., Stringer, S., Skene, N., Bryois, J., Hammerschlag, A. R., de Leeuw, C. A., Benjamins, J. S., Muñoz-Manchado, A. B., Nagel, M., Savage, J. E., Tiemeier, H., White, T., Tung, J. Y., Hinds, D. A., Vacic, V., Wang, X., Sullivan, P. F., van der Sluis, S., ... Posthuma, D. (2019). Genome-wide analysis of insomnia in 1,331,010 individuals identifies new risk loci and functional pathways. *Nature Genetics*, 51(3), 394–403. <https://doi.org/10.1038/s41588-018-0333-3>

Luppi, P.-H., Billwiller, F., & Fort, P. (2017). Selective activation of a few limbic structures during paradoxical (REM) sleep by the claustrum and the supramammillary nucleus: evidence and function. *Current Opinion in Neurobiology*, 44, 59–64. <https://doi.org/10.1016/j.conb.2017.03.002>

Stoffers, D., Altena, E., van der Werf, Y. D., Sanz-Arigita, E. J., Voorn, T. A., Astill, R. G., Strijers, R. L. M., Waterman, D., & Van Someren, E. J. W. (2014). The caudate: a key node in the neuronal network imbalance of insomnia? *Brain*, 137(2), 610–620. <https://doi.org/10.1093/brain/awt329>

Swift, K. M., Gross, B. A., Frazer, M. A., Bauer, D. S., Clark, K. J. D., Vazey, E. M., Aston-Jones, G., Li, Y., Pickering, A. E., Sara, S. J., & Poe, G. R. (2018). Abnormal Locus Coeruleus Sleep Activity Alters Sleep Signatures of Memory Consolidation and Impairs Place Cell Stability and Spatial Memory. *Current Biology*, 28(22), 3599–3609.e4. <https://doi.org/10.1016/j.cub.2018.09.054>

Vanderheyden, W. M., Poe, G. R., & Liberzon, I. (2014). Trauma exposure and sleep: using a rodent model to understand sleep function in PTSD. *Experimental Brain Research*, 232(5), 1575–1584. <https://doi.org/10.1007/s00221-014-3890-4>

Wassing, R., Schalkwijk, F., Lakbila-Kamal, O., Ramautar, J. R., Stoffers, D., Mutsaerts, H. J. M. M., Talamini, L. M., & Van Someren, E. J. W. (2019). Haunted by the past: old emotions remain salient in insomnia disorder. *Brain*, 142(6), 1783–1796. <https://doi.org/10.1093/brain/awz089>

Wassing, R., Benjamins, J. S., Talamini, L. M., Schalkwijk, F., & Van Someren, E. J. W. (2019). Overnight worsening of emotional distress indicates maladaptive sleep in insomnia. *Sleep*, 42(4), zsy268. <https://doi.org/10.1093/sleep/zsy268>

Wassing, R., Benjamins, J. S., Dekker, K., Moens, S., Spiegelhalder, K., Feige, B., Riemann, D., van der Sluis, S., Van Der Werf, Y. D., Talamini, L. M., Walker, M. P., Schalkwijk, F., & Van Someren, E. J. W. (2016). Slow dissolving of emotional distress contributes to hyperarousal. *Proceedings of the National Academy of Sciences*, 113(9), 2538–2543. <https://doi.org/10.1073/pnas.1522520113>

Wei, Y., Ramautar, J. R., Colombo, M. A., Stoffers, D., Gómez-Herrero, G., van der Meijden, W. P., te Lindert, B. H. W., van der Werf, Y. D., & Van Someren, E. J. W. (2016). I Keep a Close Watch on This Heart of Mine: Increased Interoception in Insomnia. *Sleep*, 39(12), 2113–2124. <https://doi.org/10.5665/sleep.6308>