What measures are actually effective in fighting COVID-19?

Abstract

Virus eradication or control strategy, timing, toughness and type of measures: what works and why? On the World Pandemic Research Network, several studies provide preliminary answers. Memorandum 12: overview of the studies and projects registered on WPRN database

What measures are actually effective in fighting COVID-19?

"Those who argue that lockdown measures were not necessary and that, without them, the final number of deaths would not be much different from those of a strong seasonal flu are wrong", explains Marc Fleurbaey, who has developed, with a team from Princeton University, a tool simulating the evolution of the pandemic according to the implemented measures (downloadable on WPRN). This simulator indicates that in the United Kingdom, total mortality would differ by a factor of nearly five with or without lockdown. In France, this factor is between ten and twenty. Virus eradication or control strategy, timing, toughness and type of measures: what works and why? On the World Pandemic Research Network, several studies provide preliminary answers.

Lockdown is very effective in the early stages of the pandemic.

According to the <u>Princeton simulator</u>, if a virus eradication strategy is implemented before the 20th week of the pandemic, it saves many more lives than a control strategy1, usually at a lower economic cost. But for this strategy to work, it must be firm, come early (ideally, the tenth week) and last long enough. Most countries which have succeeded in eradicating the virus (such as Vietnam, New Zealand or South Korea) adopted very early a lockdown policy coupled with testing and contact tracing in order to break the transmission chain of the virus. Once the virus has spread widely, lockdown is no longer so effective and the eradication strategy requires longer and repeated lockdown episodes: the socio-economic costs are then higher for less effectiveness.

An eradication strategy which is difficult to implement in reality.

On WPRN, Princeton University researches point out that for near-total suppression of the virus, by beginning the 15th week of the outbreak, the required lockdown duration would have to be 4 months with a 70% reduction in interpersonal contact (or 3 months with an 80% reduction in contact). Very small variations in the intensity and duration of lockdown can have a huge impact on its effectiveness. If authorities return to a more modest control strategy too soon, they will only have pushed the wave of infection into the future, while lockdown will have come at a high cost. However, it is politically difficult to impose a long and strict confinement, and very strong support measures for the populations deprived of work are necessary. In the special COVID-19 issue of the European Journal of Risk Regulation (University of Cambridge - relayed on WPRN), researchers point out that the eradication strategy was successful in New Zealand because the country began testing before its first symptomatic case and quarantined anyone entering the territory. Its insular nature has proven to be decisive. In comparison, the extent of the European Union's land borders makes absolute control impossible.

Softer measures can be more effective

A <u>study</u> published in *Nature Human Behaviour* shows that to stop the spread of the virus, an appropriate combination (sequence and timing) of less disruptive and less costly measures than lockdown can be as or even more - effective than lockdown. These results are confirmed through 226 countries, three databases and four independent methods. In terms of effectiveness, the greatest consensus surrounds closing or restricting access to places where people congregate for long periods of time (shops, bars, schools, etc.). Cancelling small gatherings decreases the Δ Rt2 virus reproduction rate between -0.22 and

-0.35, closing educational institutions decreases it from -0.15 to -0.21. Several less intrusive measures are also highly effective, including land border restrictions (ΔRt between -0.057 and -0.23) or communication about risks (ΔRt between -0.18 and -0.28: communicating on the importance of social distancing is only marginally less effective than legislating it).

Economic support has a health impact

Government food assistance programs and other financial support to vulnerable populations are also very effective (Δ Rt between -0.17 and -0.18). These measures not only have an impact on the socio-economic sphere, but also a positive effect on public health. For example, as outlined by the Oxford University COVID-19 government response tracker, allowing people to self-isolate without fear of losing their job or part of their salary can help reduce the rate of transmission. Facilitating people's access to testing by making it free is also helpful. On WPRN, the Princeton University simulator indicates that early testing is essential to reduce the time people are contagious and in contact with other people: cutting this in half reduces final mortality by a factor of five.

Timing is crucial.

The <u>CoronaNet research project</u>, led by researchers from the Technical University of Munich, New York University Abu Dhabi and Yale, and listed on WPRN has compiled more than 30,000 policies in over 190 countries. It concludes that by implementing several measures in the right order and at the right time, the reproduction rate of the virus can be reduced to below 1. Timing is crucial: the same measure can have a dramatically different impact depending on when it is implemented, which affects both the number of deaths and the length of time it takes to implement. The <u>meta-analysis</u> conducted by researchers from Oxford University indicates that strict and early restrictions are generally the most effective in limiting the number of deaths. However, they point out that while the implementation of containment measures in the early stages of a future health crisis is strongly recommended, this recipe does not always work (counter-example of Peru).

Evaluating the effectiveness of a measure: a headache.

Determining the impact of a measure is particularly difficult: several measures are often taken simultaneously and the sequence of implementation is crucial previous measures taken, in what order, stage of the pandemic, etc.). The application of the same measure may, moreover, vary from one country to another. Both the United Kingdom and France imposed lockdowns, but on the French side a certificate was required to leave one's home, whereas no document was required in England. In Madrid, at the time of the curfew in April 2021, the streets were still crowded: 45 minutes of tolerance seemed to be allowed to go home; in Canada, guards could patrol up to two hours before the curfew to ask people to leave the parks. The degree of the population'compliance with a measure has a crucial impact on its effectiveness. However, this is a parameter that is rarely measured by researchers. Results may also be biased by variations in testing and reporting policies between countries. For example, most states include exclusively confirmed COVID-19- related deaths in their statistics, however some, such as Belgium, also include suspected cases.

The effectiveness of many measures also depends on the local context. The <u>study</u> published in Nature Human Behaviour tells us that social distancing measures and travel restrictions have high entropy (effectiveness varies considerably across countries), in contrary to testing and contact cases finding. The Princeton University <u>study</u> points out that the demographic structure of the population and its general health status may also influence the lethality of the virus. On WPRN, several <u>meta-analyses</u> conclude that a policy that is effective in one setting may be ineffective in another due to a variety of factors.

What effectiveness are we talking about?

Finally, can the effectiveness of a measure be assessed only by the decrease in the transmission rate or the number of deaths? Closing schools interrupts learning and can lead to poor nutrition, stress and social isolation for children. Lockdown has greatly increased the rate of domestic violence in many countries, and has had a negative impact on the mental health of populations. It has also limited access to long-term care such as chemotherapy, with substantial impacts on patients' health and chances of survival. In France, each month of the first lockdown cost three points of annual GDP. What about individual liberties in countries that have been governed under a state of emergency for months? The study published in Nature Human Behaviour concludes that, taken together, social distancing and movement restriction policies can be considered as the "nuclear option" of measures: very effective, but causing substantial collateral damages to society, economy, and human rights.

Dictatorship vs. democracy: who does better?

If we divide the world into countries where the number of deaths is higher or lower than the average (in proportion to the population), we find in both groups rich and poor states, democracies and dictatorships, nations ruled by populists as well as by technocrats. The Oxford University meta-analysis found that no one of these characteristics was proven to predetermine the quality (or robustness) of a response to the pandemic. Some countries with strong scientific and health facilities have been overwhelmed and have had many deaths. Others, less well equipped, such as Mongolia, Thailand and Senegal, have managed to keep the population healthy and the economy going.