

RESEARCH

Thirteen genomes

Corrinne E Grover^{1*†}, Mark A Arick II², Justin C Conover¹, Jodi A Scheffler³, William S Sanders², Daniel G Peterson², Brian E Scheffler³ and Jonathan F Wendel¹

*Correspondence:

corrinne@iastate.edu

¹Department of Ecology, Evolution, and Organismal Biology, Iowa State University, Ames, IA 50011, USA

Full list of author information is available at the end of the article

†Equal contributor

Abstract

Background: Text for this section.

Results: Text for this section.

Conclusions: Text for this section.

Keywords: genome sequence; cotton; *Gossypium*; molecular evolution

Background

The American diploid “D-genome” cottons (subgenus *Houzingenia*) comprise a monophyletic clade of cytogenetically and morphologically distinct species largely distributed from Southwest Mexico to Arizona, with additional disjunct species distributions in Peru and the Galapagos Islands [Corrinne](#). Among the 13 species currently included in the D-genome [Corrinne](#) are *G. harknessii* (D2-2), an important species for cytoplasmic male sterility in cotton, and *G. raimondii* (D5), the model diploid progenitor to wild and domesticated allopolyploid cotton [Corrinne](#). The close relationship of *Houzingenia* species to the agronomically important polyploids, combined with the relative ease of sampling this subgenus for early cotton taxonomists, facilitated much of the current understanding of the relationships among D-genome species.

These early taxonomists divided subgenus *Houzingenia* into two sections and six subsections, whose species alliances have largely been retained by subsequent phylogenetic studies [Corrinne](#). Several molecular datasets have been used to evaluate these relationships, including chloroplast restriction sites [Corrinne](#); simple sequence repeat (SSR) and expressed sequence tag (EST)-SSR markers [Corrinne](#); random amplified polymorphic DNA (RAPD) markers [Corrinne](#); internal transcribed sequences (ITS) [Corrinne](#); and few single-copy nuclear genes [Corrinne](#). Relationships among the six subsections, however, remain unclear despite numerous, and often conflicting, studies [Corrinne](#). Determining the closest living relative of the D-genome ancestor to the polyploid, however, has been met with greater success. Early morphological and cytogenetic comparisons using intergenomic hybrids quickly identified *G. raimondii* as the closest living relative to the D-genome ancestor of polyploid cotton species [Corrinne](#). Subsequent analyses have largely supported this observation (Abdalla et al., 2001; Cronn 1999, Liu 2001, Cronn et al., 1996 Seelanan et al., 1997 Small et al., 1998; Small and Wendel, 2000a,b), with few conflicts [Corrinne](#).

A secondary outcome of this research has been the elucidation of multiple instances of hybridization among D-genome (i.e., *Houzingenia*) species [Corrinne](#), and, in one remarkable case (i.e., *G. gossypoides*), between a *Houzingenia* species and

Corrinne: which citations here, Endri...

Corrinne: although see Ulloa 2013

Corrinne: Endri... 1985, others?

Corrinne: Cronn al., 1996; Seelanan et al., 1997; Small and Wendel, 2000; Wendel and Albe... 1992; Wendel et al. 1995b; alvarez 20...

Corrinne: citation

Corrinne: citation

Corrinne: citation

Corrinne: citation

Corrinne: citation

Corrinne: Cronn al., 1996; Liu et al. 2001b; Small and Wendel, 2000b m... citations

Corrinne: Stephe... 1944b Hutchinson et al., 1945, 1947; Gerstel and Phill... 1958; Phillips, 19... reviewed in Wenc... and Cronn 2003

Corrinne: see, h...

Corrinne: cite cryptic
Trysts again

another, geographically isolated subgenus from Africa (either A-, B-, or, F-genome [Corrinne](#)). Notably, *G. gossypoides* is multiply introgressant, with subsequent hybridization to a member of the *G. raimondii* lineage resulting in chloroplast, if not further (and cryptic), nuclear introgression (Cronn 2003, cryptic trysts). Cytoplasmic introgression, and possibly cryptic nuclear, is also present in some populations of *G. aridum*, i.e., the Mexican Colima populations; *G. aridum* accessions derived from this location possess a *G. davidsonii*- or *G. klotzschianum*-like cytoplasm.

Corrinne: Zhao 1998

Modest attempts at understanding the evolution of the repetitive fraction of this genus support the inference of African introgression in *G. gossypoides* [Corrinne](#); however, little else is understood with respect to the evolution of the non-genic fraction of *Houzingenia*. The D-genome cottons possess the smallest genome sizes in the genus, ranging only 1.11 fold, from 841 Mb – 934 Mb. Notably, the distribution of genome sizes among the subsections suggests that this subgenus has experienced differential growth and/or reduction in genome size among species [Corrinne](#); however, the patterns of sequence gain and loss have not been characterized for the subgenus.

Corrinne: need a figure

While the differences in genome size are not dramatic, there is evidence that the transposable element types which have accumulated in *G. raimondii* are different than those that have achieved higher copy numbers than the remainder of the genus

Corrinne: Jennifer, A domestication, kokia

[Corrinne](#). Furthermore, research comparing the two sister genera to cotton (i.e., *Kokia* and *Gossypoides*; [Corrinne](#)) reveals that their apparently static genome sizes belies

Corrinne: cite Kokia paper

both gain and loss of repetitive sequence, a result similar to that of the extant members of the A-genome (subgenus *Gossypium*), whose small change in genome size (1.05X) masks differences in element accumulation [Corrinne](#).

Corrinne: cite Agenome domestication

Modern sequencing techniques make it easy to produce a substantial amount of genomic sequencing suitable for addressing these basic questions in a more genomically comprehensive manner. Here we use modest coverage Illumina sequencing to present an in-depth view of the evolution subgenus *Houzingenia*, the cotton D-genome clade. We leverage newly generated genome and plastome sequences, representing the first for many species, to address questions surrounding genome evolution in a monophyletic group of closely related species. We characterize the patterns of molecular evolution of both genes and repetitive sequences to provide insight into the pace and pattern of evolution in this subgenus. For the first time, intergenic regions are evaluated to characterize the amount of divergence outside of genes, and due to indels or single-nucleotide polymorphisms (SNPs). Finally, we revisit the phylogeny of the D-genome, both adding additional insight into the relationships among species using hundreds of nuclear genes, as well as addressing questions regarding sequence gain and loss among closely related species. The genome characterized here not only provides insight into molecular evolution on a relatively recent timeframe, but it also provides resources for comparative research and the cotton community at large.

Results

Text for this section ...

Results subsection

Text for this sub-heading ...

Discussion

Conclusions

Methods

Sequence generation and initial processing

DNA was extracted from (LEAVES) using (WHAT KIT), and sent to (WHERE) for library construction and sequencing. Sequencing was completed on the Illumina (WHAT MACHINE) using (WHICH SEQUENCING). The data were trimmed and filtered with Trimmomatic v0.32 [Corrinne](#) with the following options : (1) sequence adapter removal, (2) removal of leading and/or trailing bases when the quality score (Q) <28, (3) removal of bases after average Q <28 (8 nt window) or single base quality <10, and (4) removal of reads < 85 nt. Detailed parameters can be found at https://github.com/williamssanders/D_Cottons_USDA. [Corrinne](#)

Corrinne: citation

Corrinne: Let's port this repo to a lab site after and give the new url

Genome assembly and annotation

The trimmed data was independently assembled for each species via ABySS v2.0.1 [Corrinne](#), using every 5th kmer value from 40 through 100. A single assembly with the highest E-size (an alternative statistic to N50; [Corrinne](#)) was selected for each species and subsequently annotated with MAKER v2.31.6 [Corrinne](#) using evidence from: (1) the NCBI *G. raimondii* EST database [Corrinne](#), (2) *G. raimondii* reference genome predicted proteins, as hosted by CottonGen.org [Corrinne](#), and (3) three *ab initio* gene prediction programs, i.e. Genemark v4.30 [Corrinne](#), SNAP v2013-11-29 [Corrinne](#), and Augustus v3.0.3 [Corrinne](#). Both the SNAP and Augustus models were trained using BUSCO v2.0 [Corrinne](#).

Corrinne: citation

Corrinne: citation
Salzberg 2011

Corrinne: citation

Corrinne: citation

Corrinne: citation

Corrinne: citation

Corrinne: citation

Gene stuff

Gene orthology and family designations were determined via OrthoFinder [Corrinne](#)...

Corrinne: citation

Corrinne: citation

Repetitive characterization

Corrinne: citation

Phylogenetics and ancestral state reconstruction

Competing interests

The authors declare that they have no competing interests.

Author's contributions

Text for this section ...

Acknowledgements

We would like to thank....

Author details

¹Department of Ecology, Evolution, and Organismal Biology, Iowa State University, Ames, IA 50011, USA. ²Institute for Genomics, Biocomputing, and Biotechnology, Mississippi State University, Mississippi State, MS 39762, USA. ³Jamie Whitten Delta States Research Center, USDA-ARS, Stoneville, MS 38776, USA.

References

Figures

Figure 1 Sample figure title. A short description of the figure content should go here.

Figure 2 Sample figure title. Figure legend text.

Table 1 Sample table title. This is where the description of the table should go.

	B1	B2	B3
A1	0.1	0.2	0.3
A2
A3

Tables

Additional Files

Additional file 1 — Sample additional file title

Additional file descriptions text (including details of how to view the file, if it is in a non-standard format or the file extension). This might refer to a multi-page table or a figure.

Additional file 2 — Sample additional file title

Additional file descriptions text.