

E393048_25

The equation $x^4 + 2x + y^3 - 2y + 3 = 0$ has no integer solutions

Size $h = 35$, length $l = 12.58$

Theorem 1. *The equation*

$$x^4 + 2x + y^3 - 2y + 3 = 0 \tag{1}$$

has no integer solutions.

The proof uses a polynomial identity of the form $HT = U^2 - 2V^2$ on the curve (1), and the fact that an odd prime p with $(2/p) = -1$ divides $U^2 - 2V^2$ only to an even power. Apart from polynomial identities, which can be checked by expanding both sides, and finite checks of congruences, the proof uses only basic properties of the Jacobi symbol and the law of quadratic reciprocity.

1 Jacobi symbols and the norm form $a^2 - 2b^2$

For a prime p and a non-zero integer N , we write $v_p(N)$ for the exponent of p in the prime factorization of N . For an integer a and an odd prime p , the Legendre symbol (a/p) is 0 if $p \mid a$, 1 if a is a non-zero square modulo p , and -1 otherwise. For an odd positive integer n , the Jacobi symbol is $(a/n) = \prod_p (a/p)^{v_p(n)}$, the product over the prime factors p of n . It is multiplicative in a and in n , it depends only on a modulo n , and it is non-zero exactly when a and n are coprime. We use the law of quadratic reciprocity for Jacobi symbols together with its supplements [1, Chapter 5, §2]: for odd positive integers m and n ,

$$\begin{aligned} \left(\frac{-1}{n}\right) &= (-1)^{(n-1)/2}, & \left(\frac{2}{n}\right) &= (-1)^{(n^2-1)/8}, \\ \left(\frac{m}{n}\right)\left(\frac{n}{m}\right) &= (-1)^{(m-1)(n-1)/4} \quad \text{if } \gcd(m, n) = 1. \end{aligned}$$

Lemma 2. *Let p be an odd prime with $(2/p) = -1$, and let $a, b \in \mathbb{Z}$. If $p \mid a^2 - 2b^2$, then $p \mid a$ and $p \mid b$.*

Proof. Suppose that $p \nmid b$. Choose $c \in \mathbb{Z}$ with $bc \equiv 1 \pmod{p}$. Then $(ac)^2 \equiv 2(bc)^2 \equiv 2 \pmod{p}$, so 2 is a square modulo p . As $(2/p) \neq 0$, it is a non-zero square, so $(2/p) = 1$, a contradiction. Hence $p \mid b$, and then $p \mid a^2$, so $p \mid a$. $\square$

Lemma 3. *Let $a, b \in \mathbb{Z}$ be not both zero. Then $a^2 - 2b^2 \neq 0$, and $v_p(a^2 - 2b^2)$ is even for every odd prime p with $(2/p) = -1$.*

Proof. Since 2 is not the square of an integer, $a^2 - 2b^2 \neq 0$. Let p be an odd prime with $(2/p) = -1$, let p^k be the largest power of p dividing both a and b , and write $a = p^k a'$ and $b = p^k b'$. Then p does not divide both a' and b' , so $p \nmid a'^2 - 2b'^2$ by Lemma 2. Since $a^2 - 2b^2 = p^{2k}(a'^2 - 2b'^2)$, we get $v_p(a^2 - 2b^2) = 2k$. $\square$

Lemma 4. *Let n and n' be odd positive integers with $n' \equiv n \pmod{8}$, or with $n' \equiv -n \pmod{8}$. Then $(2/n') = (2/n)$.*

Proof. Write $2 = \varepsilon 2^e q$ with $\varepsilon = 1$, $e = 1$ and $q = 1$. By multiplicativity and the reciprocity laws,

$$\left(\frac{2}{n}\right) = \left(\frac{\varepsilon}{n}\right) \left(\frac{2}{n}\right)^e (-1)^{(n-1)(q-1)/4} \left(\frac{n}{q}\right).$$

Here (ε/n) and $(-1)^{(n-1)(q-1)/4}$ depend only on n modulo 4, $(2/n)$ depends only on n modulo 8, and (n/q) depends only on n modulo q . Since 8 is divisible by $4q$, and by $8q$, the first claim follows. If $n' \equiv -n \pmod{8}$, then, as $\varepsilon = 1$, the factor (ε/n) does not change, $(2/n)$ does not change because $n'^2 \equiv n^2 \pmod{16}$, and each of $(-1)^{(n-1)(q-1)/4}$ and (n/q) is multiplied by $(-1)^{(q-1)/2}$, because $(n' - 1)/2 \equiv (n - 1)/2 + 1 \pmod{2}$ and $(n'/q) = (-1/q)(n/q) = (-1)^{(q-1)/2}(n/q)$. This proves the second claim. $\square$

Lemma 5. *Let n be a non-zero integer, and let s , u and m be positive integers such that every prime factor of s divides 4, both $2s$ and $8s$ divide m , u is odd, $(2/u) = -1$, and $n \equiv su \pmod{m}$. Then there is an odd prime p with $(2/p) = -1$ such that $v_p(n)$ is odd.*

Proof. Since $s \mid m$, we have $s \mid n$, and $n' = n/s$ satisfies $n' \equiv u \pmod{m/s}$. Since m/s is even and divisible by 8, n' is odd and $n' \equiv u \pmod{8}$. Since $|n'| \equiv \pm u \pmod{8}$, Lemma 4 gives $(2/|n'|) = (2/u) = -1$. By multiplicativity, the product of $(2/p)^{v_p(n')}$ over the prime factors p of $|n'|$ is -1 , so some prime factor p of n' satisfies $(2/p) = -1$ and $v_p(n')$ is odd. Then p is odd, and $p \nmid 4$, so $p \nmid s$ and $v_p(n) = v_p(n')$ is odd. $\square$

2 The auxiliary polynomials

Put $F(x, y) = x^4 + 2x + y^3 - 2y + 3$, so that (1) is the equation $F(x, y) = 0$. Define

$$\begin{aligned} H &= x^2 + 2x + 4y + 1, \\ T &= x^4 + 4x^3 - 4x^2y + 6x^2 - 8xy + 4x + 16y^2 - 4y - 31, \\ U &= x^3 + 3x^2 - 13x + 17, \\ V &= 4x^2 - 4x + 16, \\ G &= 64, \\ A &= 52x - 48, \\ B &= -13x^2 - 40x + 217. \end{aligned}$$

Expanding both sides, one checks the identities

$$HT - U^2 + 2V^2 = GF, \tag{2}$$

$$AU + BV = 2^5 \cdot 83. \tag{3}$$

3 Proof of Theorem 1

Suppose that $(x, y) \in \mathbb{Z}^2$ is a solution of (1). Then $F(x, y) = 0$, and (2) gives

$$HT = U^2 - 2V^2, \tag{4}$$

where H , T , U and V now denote the values of these polynomials at (x, y) .

Step 1: congruences. Checking the 8^2 residue pairs modulo 8, we find that (1) has exactly 4 solutions modulo 8, and that $H \equiv 5 \pmod{8}$ at each of them. This residue is odd, and $(2/5) = -1$.

Step 2: H is non-zero. By (3), U and V are not both zero, so $U^2 - 2V^2 \neq 0$ by Lemma 3, and (4) gives $H \neq 0$.

Step 3: primes p with $(2/p) = -1$. Let p be an odd prime with $(2/p) = -1$. We show that $v_p(H)$ is even. By (3), U and V are not both zero, so (4) and Lemma 3 show that $v_p(HT) = v_p(U^2 - 2V^2)$ is even. Suppose that p divides both H and T . Then $p \mid U^2 - 2V^2$ by (4), so $p \mid U$ and $p \mid V$ by Lemma 2, and $p \mid 2^5 \cdot 83$ by (3). As p is odd, we get $p = 83$. Checking the 83^2 residue pairs modulo 83, we find that F , H and T have no common zero modulo 83, a contradiction. Hence p does not divide both H and T . If $p \nmid T$, then $v_p(H) = v_p(HT)$ is even, and if $p \nmid H$, then $v_p(H) = 0$.

Step 4: contradiction. By Step 1, $H \equiv su \pmod{8}$ with s and u as in Lemma 5, and $H \neq 0$ by Step 2. Hence Lemma 5 applies to $n = H$, so there is an odd prime p with $(2/p) = -1$ such that $v_p(H)$ is odd. This contradicts Step 3, and the theorem is proved. $\square$

References

- [1] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, 2nd ed., Graduate Texts in Mathematics 84, Springer, New York, 1990.