

E128544_53

The equation $-x^4 + xy + x + y^3 + 5 = 0$ has no integer solutions

Size $h = 35$, length $l = 12.32$

Theorem 1. *The equation*

$$-x^4 + xy + x + y^3 + 5 = 0 \tag{1}$$

has no integer solutions.

The proof uses a polynomial identity of the form $HT = U^2 - 7V^2$ on the curve (1), and the fact that an odd prime p with $(7/p) = -1$ divides $U^2 - 7V^2$ only to an even power. Apart from polynomial identities, which can be checked by expanding both sides, and finite checks of congruences, the proof uses only basic properties of the Jacobi symbol and the law of quadratic reciprocity.

1 Jacobi symbols and the norm form $a^2 - 7b^2$

For a prime p and a non-zero integer N , we write $v_p(N)$ for the exponent of p in the prime factorization of N . For an integer a and an odd prime p , the Legendre symbol (a/p) is 0 if $p \mid a$, 1 if a is a non-zero square modulo p , and -1 otherwise. For an odd positive integer n , the Jacobi symbol is $(a/n) = \prod_p (a/p)^{v_p(n)}$, the product over the prime factors p of n . It is multiplicative in a and in n , it depends only on a modulo n , and it is non-zero exactly when a and n are coprime. We use the law of quadratic reciprocity for Jacobi symbols together with its supplements [1, Chapter 5, §2]: for odd positive integers m and n ,

$$\begin{aligned} \left(\frac{-1}{n}\right) &= (-1)^{(n-1)/2}, & \left(\frac{2}{n}\right) &= (-1)^{(n^2-1)/8}, \\ \left(\frac{m}{n}\right)\left(\frac{n}{m}\right) &= (-1)^{(m-1)(n-1)/4} \quad \text{if } \gcd(m, n) = 1. \end{aligned}$$

Lemma 2. *Let p be an odd prime with $(7/p) = -1$, and let $a, b \in \mathbb{Z}$. If $p \mid a^2 - 7b^2$, then $p \mid a$ and $p \mid b$.*

Proof. Suppose that $p \nmid b$. Choose $c \in \mathbb{Z}$ with $bc \equiv 1 \pmod{p}$. Then $(ac)^2 \equiv 7(bc)^2 \equiv 7 \pmod{p}$, so 7 is a square modulo p . As $(7/p) \neq 0$, it is a non-zero square, so $(7/p) = 1$, a contradiction. Hence $p \mid b$, and then $p \mid a^2$, so $p \mid a$. $\square$

Lemma 3. *Let $a, b \in \mathbb{Z}$ be not both zero. Then $a^2 - 7b^2 \neq 0$, and $v_p(a^2 - 7b^2)$ is even for every odd prime p with $(7/p) = -1$.*

Proof. Since 7 is not the square of an integer, $a^2 - 7b^2 \neq 0$. Let p be an odd prime with $(7/p) = -1$, let p^k be the largest power of p dividing both a and b , and write $a = p^k a'$ and $b = p^k b'$. Then p does not divide both a' and b' , so $p \nmid a'^2 - 7b'^2$ by Lemma 2. Since $a^2 - 7b^2 = p^{2k}(a'^2 - 7b'^2)$, we get $v_p(a^2 - 7b^2) = 2k$. $\square$

Lemma 4. *Let n and n' be odd positive integers with $n' \equiv n \pmod{28}$, or with $n' \equiv -n \pmod{28}$. Then $(7/n') = (7/n)$.*

Proof. Write $7 = \varepsilon 2^e q$ with $\varepsilon = 1$, $e = 0$ and $q = 7$. By multiplicativity and the reciprocity laws,

$$\left(\frac{7}{n}\right) = \left(\frac{\varepsilon}{n}\right) \left(\frac{2}{n}\right)^e (-1)^{(n-1)(q-1)/4} \left(\frac{n}{q}\right).$$

Here (ε/n) and $(-1)^{(n-1)(q-1)/4}$ depend only on n modulo 4, $(2/n)$ depends only on n modulo 8, and (n/q) depends only on n modulo q . Since 28 is divisible by $4q$, the first claim follows. If $n' \equiv -n \pmod{28}$, then, as $\varepsilon = 1$, the factor (ε/n) does not change, $(2/n)$ does not change because $n'^2 \equiv n^2 \pmod{16}$, and each of $(-1)^{(n-1)(q-1)/4}$ and (n/q) is multiplied by $(-1)^{(q-1)/2}$, because $(n' - 1)/2 \equiv (n - 1)/2 + 1 \pmod{2}$ and $(n'/q) = (-1/q)(n/q) = (-1)^{(q-1)/2}(n/q)$. This proves the second claim. $\square$

Lemma 5. *Let n be a non-zero integer, and let s , u and m be positive integers such that every prime factor of s divides 14, both $2s$ and $28s$ divide m , u is odd, $(7/u) = -1$, and $n \equiv su \pmod{m}$. Then there is an odd prime p with $(7/p) = -1$ such that $v_p(n)$ is odd.*

Proof. Since $s \mid m$, we have $s \mid n$, and $n' = n/s$ satisfies $n' \equiv u \pmod{m/s}$. Since m/s is even and divisible by 28, n' is odd and $n' \equiv u \pmod{28}$. Since $|n'| \equiv \pm u \pmod{28}$, Lemma 4 gives $(7/|n'|) = (7/u) = -1$. By multiplicativity, the product of $(7/p)^{v_p(n')}$ over the prime factors p of $|n'|$ is -1 , so some prime factor p of n' satisfies $(7/p) = -1$ and $v_p(n')$ is odd. Then p is odd, and $p \nmid 14$, so $p \nmid s$ and $v_p(n) = v_p(n')$ is odd. $\square$

2 The auxiliary polynomials

Put $F(x, y) = -x^4 + xy + x + y^3 + 5$, so that (1) is the equation $F(x, y) = 0$. Define

$$H = x^2 - 16x - 14y + 1,$$

$$T = x^4 - 32x^3 + 14x^2y + 258x^2 - 224xy + 164x + 196y^2 + 14y + 1,$$

$$U = x^3 - 24x^2 - 11x + 132,$$

$$V = -19x^2 - 18x + 23,$$

$$G = -2744,$$

$$A = -323x + 2069,$$

$$B = -17x^2 + 533x - 2952.$$

Expanding both sides, one checks the identities

$$HT - U^2 + 7V^2 = GF, \tag{2}$$

$$AU + BV = 2^2 \cdot 3 \cdot 7^2 \cdot 349. \tag{3}$$

3 Proof of Theorem 1

Suppose that $(x, y) \in \mathbb{Z}^2$ is a solution of (1). Then $F(x, y) = 0$, and (2) gives

$$HT = U^2 - 7V^2, \tag{4}$$

where H , T , U and V now denote the values of these polynomials at (x, y) .

Step 1: congruences. Checking the 196^2 residue pairs modulo 196, we find that (1) has exactly 112 solutions modulo 196, and that at each of them H is congruent modulo 196 to one of the 22 residues r listed in Table 1. Each r is listed there as $r = su$ with s and u positive and u odd, or as r if $s = 1$, and in each case $(7/u) = -1$. Here for $s = 1$, $2s = 2$ and $28s = 28$ divide 196, and for $s = 7$, every prime factor of s divides 14, and $2s = 14$ and $28s = 196$ divide 196.

Step 2: H is non-zero. By (3), U and V are not both zero, so $U^2 - 7V^2 \neq 0$ by Lemma 3, and (4) gives $H \neq 0$.

Step 3: primes p with $(7/p) = -1$. Let p be an odd prime with $(7/p) = -1$. We show that $v_p(H)$ is even. By (3), U and V are not both zero, so (4) and Lemma 3 show that $v_p(HT) = v_p(U^2 - 7V^2)$ is even. Suppose that p divides both H and T . Then $p \mid U^2 - 7V^2$ by (4), so $p \mid U$ and $p \mid V$ by Lemma 2, and $p \mid 2^2 \cdot 3 \cdot 7^2 \cdot 349$ by (3). As p is odd and $p \neq 3$ and $p \neq 7$ because $(7/3) = 1$ and $(7/7) = 0$, we get $p = 349$. Checking the 349^2 residue pairs modulo 349, we find that F , H and T have no common zero modulo 349, a contradiction. Hence p does not divide both H and T . If $p \nmid T$, then $v_p(H) = v_p(HT)$ is even, and if $p \nmid H$, then $v_p(H) = 0$.

Step 4: contradiction. By Step 1, $H \equiv su \pmod{196}$ with s and u as in Lemma 5, and $H \neq 0$ by Step 2. Hence Lemma 5 applies to $n = H$, so there is an odd prime p with $(7/p) = -1$ such that $v_p(H)$ is odd. This contradicts Step 3, and the theorem is proved. $\square$

References

- [1] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, 2nd ed., Graduate Texts in Mathematics 84, Springer, New York, 1990.

Table 1: The residues $r = su$ of H modulo 196 in Step 1.

11	43	79	119 = 7 · 17	151	183
15	51	95	123	155	191
23	67	99	127	163	
39	71	107	135	179	