

E212901_8

The equation $x^4 + xy + x + y^3 - 3 = 0$ has no integer solutions

Size $h = 33$, length $l = 11.58$

Theorem 1. *The equation*

$$x^4 + xy + x + y^3 - 3 = 0 \tag{1}$$

has no integer solutions.

The proof uses a polynomial identity of the form $HT = U^2 + 2V^2$ on the curve (1), and the fact that an odd prime p with $(-2/p) = -1$ divides $U^2 + 2V^2$ only to an even power. Apart from polynomial identities, which can be checked by expanding both sides, and finite checks of congruences, the proof uses only basic properties of the Jacobi symbol and the law of quadratic reciprocity.

1 Jacobi symbols and the norm form $a^2 + 2b^2$

For a prime p and a non-zero integer N , we write $v_p(N)$ for the exponent of p in the prime factorization of N . For an integer a and an odd prime p , the Legendre symbol (a/p) is 0 if $p \mid a$, 1 if a is a non-zero square modulo p , and -1 otherwise. For an odd positive integer n , the Jacobi symbol is $(a/n) = \prod_p (a/p)^{v_p(n)}$, the product over the prime factors p of n . It is multiplicative in a and in n , it depends only on a modulo n , and it is non-zero exactly when a and n are coprime. We use the law of quadratic reciprocity for Jacobi symbols together with its supplements [1, Chapter 5, §2]: for odd positive integers m and n ,

$$\begin{aligned} \left(\frac{-1}{n}\right) &= (-1)^{(n-1)/2}, & \left(\frac{2}{n}\right) &= (-1)^{(n^2-1)/8}, \\ \left(\frac{m}{n}\right)\left(\frac{n}{m}\right) &= (-1)^{(m-1)(n-1)/4} \quad \text{if } \gcd(m, n) = 1. \end{aligned}$$

Lemma 2. *Let p be an odd prime with $(-2/p) = -1$, and let $a, b \in \mathbb{Z}$. If $p \mid a^2 + 2b^2$, then $p \mid a$ and $p \mid b$.*

Proof. Suppose that $p \nmid b$. Choose $c \in \mathbb{Z}$ with $bc \equiv 1 \pmod{p}$. Then $(ac)^2 \equiv -2(bc)^2 \equiv -2 \pmod{p}$, so -2 is a square modulo p . As $(-2/p) \neq 0$, it is a non-zero square, so $(-2/p) = 1$, a contradiction. Hence $p \mid b$, and then $p \mid a^2$, so $p \mid a$. $\square$

Lemma 3. *Let $a, b \in \mathbb{Z}$ be not both zero. Then $a^2 + 2b^2 \neq 0$, and $v_p(a^2 + 2b^2)$ is even for every odd prime p with $(-2/p) = -1$.*

Proof. Clearly $a^2 + 2b^2 > 0$. Let p be an odd prime with $(-2/p) = -1$, let p^k be the largest power of p dividing both a and b , and write $a = p^k a'$ and $b = p^k b'$. Then p does not divide both a' and b' , so $p \nmid a'^2 + 2b'^2$ by Lemma 2. Since $a^2 + 2b^2 = p^{2k}(a'^2 + 2b'^2)$, we get $v_p(a^2 + 2b^2) = 2k$. $\square$

Lemma 4. *Let n and n' be odd positive integers with $n' \equiv n \pmod{8}$. Then $(-2/n') = (-2/n)$.*

Proof. Write $-2 = \varepsilon 2^e q$ with $\varepsilon = -1$, $e = 1$ and $q = 1$. By multiplicativity and the reciprocity laws,

$$\left(\frac{-2}{n}\right) = \left(\frac{\varepsilon}{n}\right) \left(\frac{2}{n}\right)^e (-1)^{(n-1)(q-1)/4} \left(\frac{n}{q}\right).$$

Here (ε/n) and $(-1)^{(n-1)(q-1)/4}$ depend only on n modulo 4, $(2/n)$ depends only on n modulo 8, and (n/q) depends only on n modulo q . Since 8 is divisible by $4q$, and by $8q$, the claim follows. $\square$

Lemma 5. *Let n be a non-zero integer, and let s, u and m be positive integers such that every prime factor of s divides 4, both $2s$ and $8s$ divide m , u is odd, $(-2/u) = -1$, and $n \equiv su \pmod{m}$. Suppose also that $n > 0$. Then there is an odd prime p with $(-2/p) = -1$ such that $v_p(n)$ is odd.*

Proof. Since $s \mid m$, we have $s \mid n$, and $n' = n/s$ satisfies $n' \equiv u \pmod{m/s}$. Since m/s is even and divisible by 8, n' is odd and $n' \equiv u \pmod{8}$. Since $n > 0$, Lemma 4 gives $(-2/n') = (-2/u) = -1$. By multiplicativity, the product of $(-2/p)^{v_p(n')}$ over the prime factors p of n' is -1 , so some prime factor p of n' satisfies $(-2/p) = -1$ and $v_p(n')$ is odd. Then p is odd, and $p \nmid 4$, so $p \nmid s$ and $v_p(n) = v_p(n')$ is odd. $\square$

2 The auxiliary polynomials

Put $F(x, y) = x^4 + xy + x + y^3 - 3$, so that (1) is the equation $F(x, y) = 0$. Define

$$\begin{aligned} H &= 8x^2 - 2xy - 2x + y^2 - 4y + 11, \\ T &= -2x^5 - x^4y + 60x^4 + 16x^3y - 44x^3 - 4x^2y^2 + 28x^2y + 199x^2 \\ &\quad + 19xy^2 + 13xy - 48x + 5y^2 + 47y + 133, \\ U &= CU_0, \\ V &= CV_0, \\ C &= x^2 + 2, \\ U_0 &= x^2 - 20x + 12, \\ V_0 &= 5x + 11, \\ G &= -x^4 + 24x^3 - 4x^2y + 6x^2 + 19xy - 73x + 5y + 27, \\ A &= 25, \\ B &= -5x + 111, \\ \alpha &= -611456x^9 + 28095904x^8 - 440969594x^7 + 1630206891x^6 \\ &\quad - 5815091414x^5 + 8094185180x^4 - 4434445574x^3 + 4179415149x^2 \\ &\quad + 1104823504x + 6552300176, \\ \beta &= 38216x^9 - 1292625x^8 + 152864x^7y + 19657312x^7 - 2227868x^6y \\ &\quad - 94539800x^6 + 1529915x^5y + 395509210x^5 + 26395093x^4y \\ &\quad - 500325634x^4 + 29786886x^3y + 333388309x^3 - 62309431x^2y \\ &\quad - 447571337x^2 - 110450224xy - 262437557x - 23353855y - 192338167, \\ \gamma &= -152864x^6 + 38216x^5y + 4253316x^5 - 375441x^4y - 10001704x^4 \\ &\quad - 1353096x^3y + 29439360x^3 - 297734x^2y - 33731479x^2 + 4341115xy \\ &\quad - 6506043x + 4670771y - 24120698. \end{aligned}$$

Expanding both sides, one checks the identities

$$HT - U^2 - 2V^2 = GF, \tag{2}$$

$$AU_0 + BV_0 = 3^2 \cdot 13^2, \quad (3)$$

$$\alpha C + \beta H + \gamma T = 3^{12} \cdot 11^4, \quad (4)$$

$$4C = (2x)^2 + 8, \quad (5)$$

$$7H = 7(x - y + 2)^2 + (7x - 3)^2 + 40. \quad (6)$$

3 Proof of Theorem 1

Suppose that $(x, y) \in \mathbb{Z}^2$ is a solution of (1). Then $F(x, y) = 0$, and (2) gives

$$HT = U^2 + 2V^2, \quad (7)$$

where H, T, U and V now denote the values of these polynomials at (x, y) .

Step 1: congruences. Checking the 64^2 residue pairs modulo 64, we find that (1) has exactly 32 solutions modulo 64, and that $H \equiv 56 \pmod{64}$ at each of them. We write this residue as su with s and u positive and u odd: $56 = 8 \cdot 7$ with $(-2/7) = -1$. Here for $s = 8$, every prime factor of s divides 4, and $2s = 16$ and $8s = 64$ divide 64.

Step 2: signs. By (6), $H > 0$.

Step 3: primes p with $(-2/p) = -1$. Let p be an odd prime with $(-2/p) = -1$. We show that $v_p(H)$ is even. By (5), $C > 0$, and by (3), U_0 and V_0 are not both zero. Since $U^2 + 2V^2 = C^2(U_0^2 + 2V_0^2)$, (7) and Lemma 3 show that $HT \neq 0$ and that $v_p(HT) = 2v_p(C) + v_p(U_0^2 + 2V_0^2)$ is even. Suppose that p divides both H and T . If $p \mid C$, then $p \mid 3^{12} \cdot 11^4$ by (4). But p is odd, and $p \neq 3$ and $p \neq 11$ because $(-2/3) = 1$ and $(-2/11) = 1$, a contradiction. Otherwise $p \mid U_0^2 + 2V_0^2$ by (7), so $p \mid U_0$ and $p \mid V_0$ by Lemma 2, and $p \mid 3^2 \cdot 13^2$ by (3). As p is odd and $p \neq 3$ because $(-2/3) = 1$, we get $p = 13$. Checking the 13^2 residue pairs modulo 13, we find that the only common zero of F, H and T modulo 13 is $(3, 5)$. If $(x, y) \equiv (3, 5) \pmod{13}$, write $x = 3 + 13X$ and $y = 5 + 13Y$ with $X, Y \in \mathbb{Z}$. Then $F(x, y) = 13L(X, Y)$ and $H(x, y) - F(x, y) = 13^2K(X, Y)$, where

$$\begin{aligned} L(X, Y) &= 2197X^4 + 2028X^3 + 702X^2 + 13XY + 114X + 169Y^3 + 195Y^2 + 78Y + 17, \\ K(X, Y) &= -169X^4 - 156X^3 - 46X^2 - 3XY - 6X - 13Y^3 - 14Y^2 - 6Y - 1 \end{aligned}$$

have integer coefficients and no common zero modulo 13. Since $F(x, y) = 0$, we get $L(X, Y) = 0$, so $13 \nmid K(X, Y)$ and $v_{13}(H) = 2$, which is even. Hence, apart from the case treated above, where $v_p(H) = 2$, p does not divide both H and T . If $p \nmid T$, then $v_p(H) = v_p(HT)$ is even, and if $p \nmid H$, then $v_p(H) = 0$.

Step 4: contradiction. By Step 1, $H \equiv su \pmod{64}$ with s and u as in Lemma 5, and $H > 0$ by Step 2. Hence Lemma 5 applies to $n = H$, so there is an odd prime p with $(-2/p) = -1$ such that $v_p(H)$ is odd. This contradicts Step 3, and the theorem is proved. $\square$

References

- [1] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, 2nd ed., Graduate Texts in Mathematics 84, Springer, New York, 1990.